

Effect of Different Water Stress on Growth Index and Yield of Semilate Rice [†]

Yixin Zhang ¹, Xinyu Zhao ^{1,*}, Fangping Liu ², Lvdan Zhu ¹ and Honglei Yu ²

¹ School of Hydraulic and Ecological Engineering, Nanchang Institute of Technology; 2021301026@nit.edu.cn (Y.Z.); 515244985@qq.com (L.Z.)

² Jiangxi Provincial Irrigation Experimental Station; shong62@163.com (F.L.); 1016974093@qq.com (H.Y.)

* Correspondence: spxyz@nit.edu.cn; Tel.: +86-18970091896

[†] Presented at the 7th International Electronic Conference on Water Sciences, 15–30 March 2023; Available online: <https://ecws-7.sciforum.net>.

Abstract: In order to find out the effects of different degrees of drought stress on the growth index and yield of semilate rice. In this experiment, a pot experiment was conducted with hybrid rice Tianyouhuazhan as material. Intermittent irrigation was used as control (CK) at tillering stage, booting stage, heading stage, and milk stage. Three water stress gradients of light drought (70% saturated moisture content), medium drought (60% saturated moisture content), and severe drought (50% saturated moisture content) were set to monitor soil moisture content. The plant height, number of tillers, leaf area, and chlorophyll content under different water gradients at different growth stages were measured, and the effects of different water stress treatments at different growth stages on rice growth index, yield, and its components were analyzed. The results showed that the inhibition of plant height at the jointing and booting stage was the most significant, the number of tillers and leaf area decreased most significantly at tillering stage, and the heading and flowering stage had the greatest effect on chlorophyll synthesis. As far as yield is concerned, the most sensitive stage is the heading and flowering stage. Different stress treatments reduced the effective panicle number, 1000-grain weight, and seed setting rate of rice, and reduced the yield by 27.57%, 44.23%, and 46.32% respectively compared with the normal control. The correlation analysis showed that the correlation degree of affecting yield from large to small was 1000-grain weight, seed setting rate, and effective panicle. Therefore, ensuring water supply at the heading and flowering stage can effectively improve 1000-grain weight, effective panicle, and seed setting rate, and increase rice yield and water use efficiency.

Citation: ZHANG, Y.; ZHAO, X.; LIU, F.; ZHU, L.; YU, H. Effect of Different Water Stress on Growth Index and Yield of Semilate Rice. *Environ. Sci. Proc.* **2023**, *5*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Published: 15 March 2023



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: water stress; growth index; 1000 grain weight; yield

1. Introduction

There is a great shortage of water resources in China, and the biological characteristics of rice have great water-saving potential [1]. therefore, in the process of rice planting, it is of great significance to study the tolerance of water stress at different growth stages and its effects on the physiological index and yield of rice [2–5]. There were differences in the effects of water stress and the degree of water stress on rice growth and yield at different growth stages; compared with rice under severe water shortage, rice recovered better and had a higher compensation effect under mild water stress. Water stress at the booting stage decreased the number of effective panicles, grains per panicle, and seed setting rate, resulting in a decrease in yield [6,7]. Water stress at the flowering stage resulted in a significant decrease in 1000-grain weight and seed setting rate, resulting in a decrease in yield per plant [8,9]. There have been many studies on the effects of water stress on rice yield, but the conclusions are different due to different planting sites, rice

varieties, and experimental methods. there are relatively few studies on the effects of water stress on rice yield and growth index in a reproductive growth stage. In this experiment, according to the actual situation of seasonal drought of mid-season rice in Jiangxi, the effects of water stress on rice yield and growth index were studied, which provided theoretical support for the study of rice water requirement and water-saving irrigation in Jiangxi.

2. Materials and Methods

2.1. Experimental Design

The experiment was carried out in the irrigation experiment center of Jiangxi Province, using pot experiment, pot length 38 cm, pot width 26 cm, carried out in a mobile rain shelter to avoid the effect of rainfall on pot culture. The middle rice variety Yangliangyou was selected, and four middle rice varieties were selected at tillering stage (T1), jointing and booting stage (T2), heading and flowering stage (T3) and milk stage (T4). The degree of drought was controlled by the soil water content after the water layer was dried, and three levels were set, which were 50% of the saturated moisture content (W1, severe drought), 60% (W2, moderate drought) and 70% (W3, light drought) respectively. The two factors were combined into 12 treatments, and another reference treatment (CK) was set up. There were 13 treatments for late rice. Reference treatment (CK) intermittent irrigation, always maintain 2–3 cm thin water layer in the green period, 7–10 days in the late tillering period, and naturally dry in the yellow mature stage; the rest of the growth period irrigates 2–3 cm each time and dries naturally for 3–5 days, that is, the soil moisture content reaches more than 70% of the saturated moisture content, and then irrigates again and again. Each process sets 3 repeats, with a total of 39 cells (see Table 1).

Table 1. Experimental Design of semi-late Rice subjected to drought.

Serial Number	Treatment Mode	Water Condition of Paddy Field at Different Stages						Yellow Ripening Stage
		Rejuventi on Stage	Tillering Stage	Booting Stage	Heading Stage	Milk Stage		
1	CK	Normal	Normal	Normal	Normal	Normal	Dorp dry	
2	W1T1	Normal	70%	Normal	Normal	Normal	Dorp dry	
3	W2T1	Normal	60%	Normal	Normal	Normal	Dorp dry	
4	W3T1	Normal	50%	Normal	Normal	Normal	Dorp dry	
5	W1T2	Normal	Normal	70%	Normal	Normal	Dorp dry	
6	W2T2	Normal	Normal	60%	Normal	Normal	Dorp dry	
7	W3T2	Normal	Normal	50%	Normal	Normal	Dorp dry	
8	W1T3	Normal	Normal	Normal	70%	Normal	Dorp dry	
9	W2T3	Normal	Normal	Normal	60%	Normal	Dorp dry	
10	W3T3	Normal	Normal	Normal	50%	Normal	Dorp dry	
11	W1T4	Normal	Normal	Normal	Normal	70%	Dorp dry	
12	W2T4	Normal	Normal	Normal	Normal	60%	Dorp dry	
							Dorp dry	
							Dorp dry	
13	W3T4	Normal	Normal	Normal	Normal	50%	Dorp dry	
							Dorp dry	
							Dorp dry	

2.2. Observation Items and Methods

The control methods of soil moisture are as follows: (1) before rice planting, weigh the test chamber, measure the quality of wet soil and test box, measure the soil moisture content by drying method, and then calculate the dry soil mass in each soil box; (2) in the

process of drought test after rice planting, weigh the weight in the test box, calculate the soil water content in the test box, and control the soil water content.

Plant height: measure the distance from the surface of the potted soil layer to the top of the longest leaf before the heading and flowering stage, and the distance from the surface of the potted soil layer to the longest ear from the heading stage to the later stage.

The number of tillers: from the tillering stage, the number of tillers per pot was recorded and measured every 5 days

Chlorophyll: using hand-held SPAD-502 chlorophyll tester, 6 plants were selected for each plant, one leaf was completely opened for each plant, and the SPAD values at the middle part of the leaf and at the upper and lower part of the leaf were measured, and the average SPAD values were taken.

Leaf area: the first three plants of each pot were measured directly with a ruler, and three inverted sword leaves were measured in each plant, and the midvein length and maximum leaf width of each leaf were measured respectively.

Yield and quality: according to the results of effective panicle investigation, 3 plants were selected in each pot area for seed testing; at the same time, the yield of all rice plants in each pot was measured. The contents of seed examination include effective panicle length, effective panicle number, total grains per panicle, filled grain weight per panicle, 1000-grain weight and actual yield.

3. Results

3.1. Analysis of Growth Indexes of Rice under Different Water Stress

It can be seen from Figure 1a that drought stress has an inhibitory effect on the plant height of rice during the whole growth stage, and the order of plant height from high to low under each water stress is as follows: CK > W3T4 > W2T4 > W1T4 > W3T3 > W2T3 > W1T3 > W3T2 > W2T2 > W1T2 > W2T1 > W1T1 > W1T2. It can be seen from the figure that drought stress has the most significant inhibitory effect on plant height at jointing and booting stage, and the effect of drought stress on plant height under drought stress is less than that in the previous two stages, which proves that water has a greater effect on plant height growth in vegetative growth stage.




Figure 1. (a) Rice plant height under different water stress; (b) Rice tillering under different drought treatments.

It can be seen from Figure 1b that water stress inhibited the tillering of rice, which decreased most obviously at tillering stage (T1) and booting stage (T2), followed by heading stage (T3) and milk stage (T4). It was proved that drought stress had a strong inhibitory effect on rice tillering at vegetative growth stage.

It can be seen from Figure 2a that the SPAD value of rice during the overall growth period is in a downward trend, and the chlorophyll content of rice will increase briefly during the same growth period, and some treatments are higher than the normal control. The treatment in T1 stage is as follows: W2 > W1 > W3 in T2 and W3 > W2 > W1 in T3, but it is basically lower than that in normal control. During the period of T4, W3 > W2 > W1 and W3 increased briefly during drought and was higher than that of normal control. the gradient change of W2 treatment was smaller than that of other treatments, and the decrease of W1 treatment was the most obvious.




Figure 2. (a) Rice SPAD under different drought treatments; (b) Leaf area of Rice under different drought treatments.

It can be seen from Figure 2b that water stress can inhibit the growth of rice leaf area, in which tillering stage and jointing stage are the most obvious, followed by heading stage, and the effect on milky rice is less, the overall performance is W3 > W2 > W1, that is, the more serious the water deficit, the smaller the leaf area.

3.2. Analysis of Yield and Its Components under Drought Treatment

It can be seen from Table 2 that the effect of drought stress on rice yield is the greatest at heading stage, and the highest reduction in rice yield is 46.32%. In the same period, it is W3 < W2 < W1, and the effect at tillering stage is the least significant.

Table 2. Examination results of yield indexes in growth period of each treatment.

Treatment	Number of Effective Ears (Piece/Pot)	1000 Grain Weight (g)	Spike Length (cm)	Number of Real Grains	Number of Real Grains	Total Grains	Seed Setting Rate (%)	Actual Yield (g/Pot)	Increase or Decrease Compared with CK (%)
CK	62	19.86	19.54	3916	245	4161	94%	77.54	0
W3T1	61	19.7	19.3	3869	341	4210	92%	76.22	-1.7
W2T1	56	19.31	18.9	3795	368	4163	91%	73.28	-5.49
W1T1	52	18.81	17.7	3640	402	4042	90%	68.47	-11.7
W3T2	58	19.2	19.3	3649	450	4099	89%	70.06	-9.64
W2T2	54	18.1	18.6	3377	484	3861	87%	61.12	-21.17
W1T2	49	17.4	18.4	3027	610	3637	83%	56.15	-27.58
W3T3	57	17.7	16.9	3173	733	3906	81%	56.16	-27.57
W2T3	53	16.3	16.8	2453	979	3432	71%	43.24	-44.23
W1T3	47	16.1	16.7	2385	1076	3461	69%	41.62	-46.32
W3T4	57	18.58	18.51	2354	610	2964	79%	58.6	-24.42
W2T4	54	17.87	18.27	2337	795	3132	75%	54.27	-30.01
W1T4	49	16.96	17.84	2264	866	3130	72%	50.27	-35.17

Table 2. Water consumption and actual yield at each growth stage.

Variable Treatment Mode	Water Consumption/mm				Yield/(g/Pot)
	Tillering Stage	Booting Stage	Heading Stage	Milk Stage	
CK	275.86	367.35	121.86	140.37	77.54
W1T1	274.07	366.35	120.44	117.51	76.22
W2T1	262.54	355.00	116.28	110.54	73.28
W3T1	257.15	351.78	115.40	103.32	68.47
W1T2	233.92	350.88	111.68	138.84	70.06
W2T2	212.11	311.93	101.54	131.57	61.12
W3T2	204.14	312.45	97.73	127.59	56.15
W1T3	261.87	331.18	83.70	135.93	56.16
W2T3	259.09	323.00	66.24	137.25	43.24
W3T3	248.60	299.33	58.27	132.61	41.62
W1T4	254.49	243.80	107.36	129.50	58.60
W2T4	262.01	198.53	107.84	134.21	54.27
W3T4	252.82	177.78	101.58	129.50	50.27

The yield components include 1000-grain weight, grain number per spike, seed setting rate and effective panicle number. From the point of view of effective panicle, in the same growth period, the order is CK > W3 > W2 > W1. In terms of seed setting rate, severe drought stress at heading stage had the greatest effect on rice seed setting rate, which was 25% lower than that of CK. In the same growth period, it was CK > W3 > W2 > W1. In terms of effective panicle length, severe drought stress at heading stage had the most significant effect on effective panicle length, with an average decrease of 2.84 cm per plant, and the order was CK > W3 > W2 > W1 in the same period. In terms of grain number per ear, light drought stress at milk stage had the greatest effect on grain number per ear, which was 28.77% less than that of CK. In the same growth period, the milk stage is CK > W2 > W1 > W3. In terms of 1000-grain weight, the effect of severe drought at heading stage on 1000-grain weight was the most obvious, which was 18.93% lower than that of CK, and the order was CK > W3 > W2 > W1 in the same period.

3.3. Correlation Analysis between Growth Indexes and Stage Water Consumption of Rice

The following conclusions can be drawn from Table 3: Under the normal irrigation level, the correlation between water consumption and plant height at the jointing stage is the largest. The correlation between water consumption at heavy drought levels and plant height at the tillering stage is the strongest, 0.947. The weak correlation between light drought and moderate drought levels at different growth stages shows that severe drought at tillering stage will significantly impact rice plant height. Under the conventional irrigation mode, most of the water consumption at the jointing stage is used to increase the plant height of rice. Water consumption at light drought level is negatively correlated with rice tillering at all stages, with the strongest correlation at the jointing stage being 0.587. Water consumption at heavy drought levels strongly correlates with rice tillering at all stages, with the most significant correlation at tillering stage being 0.926. Normal irrigation level and moderate drought level are weakly correlated with other treatments, indicating that under light drought level, Most of the water consumed by rice at the jointing stage was not used for the increase of tiller number, while under the heavy drought level, most of the water consumed at each growth stage was used for the increase of tiller number of rice, especially for rice at the tillering stage, which had a significant positive correlation.

Table 3. Correlation analysis of water consumption with plant height and tiller number.

Index Water Gradient	Child-Bearing Period							
	Plant Height/cm				Tiller Number/Tiller			
	T1	T2	T3	T4	T1	T2	T3	T4
CK	-0.217	0.805 **	0.653 *	0.531	-0.023	0.415	0.435	0.400
W3	-0.379	-0.332	-0.523	-0.520	-0.467	-0.587 *	-0.524	-0.576 *
W2	-0.363	-0.242	0.232	0.336	-0.209	-0.014	-0.152	-0.112
W1	0.947 **	0.292	0.291	0.386	0.926 **	0.637 *	0.701 **	0.689 **

T1 represents the tillering stage of rice, T2 represents the booting stage of rice, T3 represents the heading stage of rice, T4 represents the milky stage of rice. **. At 0.01 level (double tail), the correlation was significant. *. At 0.05 level (double tail), the correlation was significant.

The following conclusions can be drawn from Table 4: Water consumption of rice under moderate and severe drought levels has a significant positive correlation with SPAD at tillering, jointing, and milking stages, which are 0.919, 0.569, and 0.835, respectively. In contrast, water consumption under normal irrigation and light drought levels have no significant impact on SPAD value, proving that under severe stress at tillering stage, most water consumption is used to synthesize chlorophyll. Water consumption is the main factor affecting SPAD value change under moderate stress at the jointing stage and heading stage. Under normal irrigation, water consumption and leaf area at the milky stage are negatively correlated, with a coefficient of 0.765. Therefore, water consumption does not promote leaf area growth at the milky stage. In contrast, water consumption and leaf area at the jointing stage are positively correlated, with a coefficient of 0.59, indicating that water consumption promotes leaf area growth. Under moderate and severe drought levels, rice leaf area and water consumption are positively correlated, with coefficients of 0.718 and 0.719, respectively. It was proved that water stress had a significant impact on leaf area.

Table 4. Correlation analysis of water consumption with SPAD and leaf area.

Index Water Gradient	Child-Bearing Period							
	SPAD Value				Leaf Area/cm ²			
	T1	T2	T3	T4	T1	T2	T3	T4
CK	-0.418	0.548	0.190	-0.765 **	0.416	0.590 *	0.321	-0.765 **
W3	-0.389	0.427	-0.063	0.425	-0.384	0.043	0.493	0.425
W2	-0.476	0.569 *	0.835 **	0.107	-0.229	0.719 **	0.027	0.107
W1	0.919 **	-0.286	-0.627 *	-0.198	0.718 **	-0.317	0.534	-0.198

** At 0.01 level (double tail), the correlation was significant. * At 0.05 level (double tail), the correlation was significant.

3.4. Correlation Analysis of Water Consumption and Yield at Each Growth Stage

From the results of Pearson correlation analysis in Appendix A Table A2, the following conclusions can be drawn: the water consumption at the booting stage and heading stage is significantly correlated with the yield of rice, that is, the water deficit at the booting stage and heading stage will have a severe impact on the yield of rice, among which the correlation at heading stage is the strongest, which is 0.88. Therefore, it is inappropriate to lack water at this growth stage excessively. In contrast, the water consumption at the tillering and milky stages are weakly correlated and negatively correlated, respectively, indicating that appropriate water stress at a tillering stage has little effect on the final yield of rice. The same is accurate at the milky stage.

4. Discussion

4.1. Effect of Water Stress on the Growth of Semilate Rice

In this study, it was found that with different water gradients, the overall plant height of rice under heavy drought was lower than that under light and moderate drought, and under the same water condition, the plant height was CK > T4 > T3 > T1 > T2. It can be inferred that water stress has a greater inhibitory effect on rice plant height at jointing stage and tillering stage, which is consistent with the research results of some scholars [10,11].

Tillering shows that under the same water stress, T1 < T2 < T3 < T4 < CK, this is because when rice is in reproductive growth stage, mild and moderate water stress has little effect on rice tillering in reproductive growth stage, and is beneficial to reduce ineffective tiller. Severe water stress leads to damage of late filling and dry matter distribution, resulting in death of effective tiller, which is similar to the conclusion of Yang X L [12].

The overall photosynthesis showed a downward trend, the SPAD value of rice at booting stage increased during the short-term drought period, and most of the rice was higher than CK. Short-term water deficit could promote the photosynthetic potential of chlorophyll. When the drought time is longer than 10 days, the SPAD value of rice decreases continuously, which is basically consistent with the results of previous studies, this is because at the early stage of reproductive growth, water can affect the synthesis of chlorophyll in rice, and proper water stress can increase the concentration of chlorophyll, thus promoting the photosynthetic effect of rice.

Water stress can inhibit the growth of rice leaves, thus inhibit the growth of leaf area, the overall performance is W3 > W2 > W1, that is, the more serious the water deficit, the smaller the leaf area.

4.2. Effect of Water Stress on Yield of Semilate Rice

In this study, from the yield and its composition, it was found that under different water treatments, the effective panicles were T1 > T2 > T4 > T3. Drought stress inhibited the differentiation of effective panicles, resulting in a significant decrease in the number of effective panicles [13,14]. It was found that the number of effective panicles of rice was

very low under the irrigation mode of severe water deficit, which was similar to the results of this experiment [15]. Proper water addition was needed to improve the differentiation of effective panicles in the middle and later stages of reproductive growth.

Under the same water treatment, the panicle length of light drought was $T1 \geq T2 > T4 > T3$, the moderate drought was $T1 > T2 > T4 > T3$, and the severe drought was $T2 > T4 > T1 > T3$. In the late growth stage, the inhibitory effect of water stress on rice panicle length is very obvious, this is because the flag leaf of rice is the main organ for photosynthesis, water deficiency will lead to the curl of rice flag leaves and affect the opening and closing of stomata and the synthesis of chlorophyll. So that the ear can not grow normally [16,17].

The effect of light drought on the number of grains per panicle was the least at tillering stage and the greatest at milking stage. Water played a key role in millet filling in the later stage of rice reproductive growth, and water deficit inhibited grain filling and starch accumulation in rice. studies have shown that water stress significantly reduces the total amount of dry matter and limits the quantity and rate of stem and sheath carbohydrate transport to grains at heading and flowering stage. At the same time, it accelerated the senescence of rice leaves, led to insufficient grain filling time and reduced yield [18].

Under the same water treatment, 1000-grain weight was $T1 > T2 > T4 > T3$, which proved that water stress had more significant inhibitory effect on rice grain filling at heading stage. some scholars found that drought stress at heading stage led to a significant decrease in rice yield. the main factor leading to yield reduction is the decrease of seed setting rate and 1000-grain weight due to insufficient water supply under drought stress [12], which is similar to the results of this experiment.

5. Conclusions

Based on the above analysis results, the effects of different water stress on rice growth index and yield were concluded:

- (1) Water stress at different growth stages had different effects on rice growth indexes. The plant height was most significantly inhibited at the jointing and booting stage, and its water consumption was significantly positively correlated with plant height, which was 0.805. The tillering stage most obviously affected the tiller number and leaf area. The correlation between water consumption, tiller number, and leaf area under heavy drought levels was very significant, which was 0.926 and 0.718. The heading and flowering stages had the most significant impact on chlorophyll. The correlation between water consumption and chlorophyll under medium drought levels was significant, which was 0.835.
- (2) Different stress treatments reduced the adequate panicle number, 1000-grain weight, and seed-setting rate of rice. The most sensitive period was the heading and flowering stage. Compared with the standard control, the yield of the three stress levels decreased by 27.57%, 44.23%, and 46.32%, respectively. The correlation analysis showed that the correlation degree of yield was 1000-grain weight, seed setting rate, and adequate panicle number (see Appendix A Table A1). The 1000-grain weight can have a more significant impact on the yield, which is 0.980.
- (3) There was a specific relationship between water consumption and rice yield at different growth stages. The water consumption at the booting and heading stages was significantly correlated with rice yield. The correlation between water consumption and yield at the heading stage was the strongest, which was 0.880.

Therefore, increasing water input at the tillering and jointing stages is beneficial to tillering of rice and can effectively promote stem and leaf growth and panicle differentiation and improve plant height and leaf area of rice. It is unsuitable for saving water excessively at the heading and flowering stage. Increasing the water supply can improve 1000-grain weight, effective panicle, and seed setting rate of rice, and improve rice yield and water use efficiency.

Funding: This research was funded by Jiangxi Natural Science Foundation Project, grant number 20202BABL204068.

Appendix A

Table A1. Correlation Analysis of yield and its components.

Dimension	Effective Panicle Number	1000-Grain Weight	Number of Grains per Spike	Seed Setting Rate	Actual Yield
Effective panicle number	1				
1000-grain weight	0.820 **	1			
Number of grains per spike	0.517	0.643 *	1		
Seed setting rate	0.684**	0.912 **	0.835 **	1	
Actual yield	0.762**	0.980 **	0.755 **	0.961 **	1

** . At 0.01 level (double tail), the correlation was significant. * . At 0.05 level (double tail), the correlation was significant.

Table A2. Correlation analysis of water consumption and yield at each growth stage.

Dimension	Tillering Stage	Booting Stage	Heading Stage	Milk Stage	Actual Yield
Tillering stage	1				
Booting stage	0.078	1			
Heading stage	0.175	0.159	1		
Milk stage	-0.125	-0.236	-0.389	1	
Actual yield	0.247	0.585 *	0.880 **	-0.377	1

** . At 0.01 level (double tail), the correlation was significant. * . At 0.05 level (double tail), the correlation was significant.

References

- Shao, X.W.; Liu, H.D.; Du, Z.Y.; et al. Effects of water treatments at different stages on growth and yield of rice. *J. Soil Water Conserv.* **2007**, *193*, 196.
- Zheng, G.P.; Li, J.F.; Qian, Y.D.; et al. Effects of soil moisture on rice yield and quality. *Acta Crops* **2006**, 1261–1264.
- Guo, H.; Ma, J.; Li, S.X.; et al. Studies on the compensation effects of water stress at booting stage on some physiological characteristics and yield of rice. *J. South. Agric.* **2013**, *44*, 1448–1454.
- Guo, H.; Ma, J.; Li, S.X.; et al. Effects of Water stress at booting stage on partial physiological characteristics and yield compensation of Rice. *Agric. Sci. Technol.* **2013**, *14*, 1750–1755.
- Feng, X.L.; Zhong, K.Y.; Tang, X.R.; et al. Effects of water stress at booting stage on stomatal traits and CAT activity in flag leaves of rice. *J. Irrig. Drain.* **2014**, *33*, 135–137.
- Hu MM Gong, J.; Lan, Y.; et al. Effects of water stress at different growth stages on yield and quality of rice. *J. Jiangxi Agric. Univ.* **2021**, *43*, 971–982.
- Zheng, C.J.; Li, S. Effect of water stress at flowering stage on rice growth and rice quality. *China Rice* **2017**, *23*, 43–45.
- Sun, Z.Y.; Xiao, M.G.; Zhao, B.P.; et al. High yield cultivation techniques of rice at reproductive growth stage in cold regions *Anhui Agric. Bull.* **2021**, *27*, 42–44.
- Li, X.Y.; He, Y.X.; Li, S.W.; Wang, C.T. Study on agronomic regulation of rice to drought stress. *J. Southwest Agric.* **2005**, 244–249.
- Wang, Y.L.; Wang, J.; Du, J.Z.; et al. Effects of drought stress in different periods on agronomic characters of millet. *North China J. Agric.* **2012**, *27*, 125–129.

11. Duan, S.M.; Yang, A.Z.; Huang, Y.D.; et al. Effects of drought stress on growth, physiological characteristics and yield of rice. *J. Nucl. Agric.* **2014**, *28*, 1124–1132.
12. Yang, X.L.; Wang, B.F.; Chen, L.; et al. Effects of drought at heading stage on physiological characters and yield of rice. *China Rice* **2015**, *21*, 138–141.
13. Zhu, H.P.; Li, G.Y.; Xia, Q.M.; et al. Effects of drought stress at different stages on rice yield and growth characteristics. *China Rice* **2017**, *23*, 135–138.
14. Zhang, L.Q. Research on the relationship between leaf area index and its composition at the early growth stage of rice and yield. *J. Huaiyin Inst. Technol.* **2004**, 71–76.
15. Yang, X.L.; Wang, B.F.; Li, Y.; et al. A review of effects of drought stress on agronomic characteristics and yield of rice. *Hubei Agric. Sci.* **2020**, *59*, 39–43.
16. Xu, W.; Cui, K.; Xu, A.; Nie, L.; Huang, J.; Peng, S. Drought stress condition increases root to shoot ratio via alteration of carbohydrate partitioning and enzymatic activity in rice seedlings. *Acta Physiol. Plant.* **2015**, *37*, 9.
17. Yang, X.; Wang, B.; Chen, L.; Li, P.; Cao, C. The different influences of drought stress at the flowering stage on rice physiological traits, grain yield, and quality. *Sci. Rep.* **2019**, *9*, 3742.
18. Shao, X.W.; Zhang, R.Z.; Qi, C.Y.; Tong, S.Y.; Yang, M. Effect of water stress at jointing and booting stage on rice growth and yield. *J. Jilin Agric. Univ.* **2004**, 237–241.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.