

Proceedings

Photometric determination of iron(III) with 2-naphthylcarboxymethylene citrate

R.S. Jurayev^{1,*}, A.U. Choriev² and N.T.Qaxxorov¹

¹ Department of "Chemical Engineering and Quality Management", Shakhrisabz Branch of Tashkent Institute of Chemical Technology, 20, Shahrisabz str., Shakhrisabz 181306, Uzbekistan; jurayevorganikqdu-1992@mail.ru(J.S.R.); navrozbekqahhorov70@gmail.com(Q.T.N.);

² Department of "Organic chemistry", Karshi State University, 17, Kuchabog str., Karshi 180103, Uzbekistan; azimjon-organik@mail.ru(A.U.C.)

* Correspondence: jurayevorganikqdu-1992@mail.ru)

Abstract: The most prevalent element in nature is iron, which is also one of the essential, active macroelements. Today, photometric techniques are frequently employed to identify harmful and extremely poisonous heavy metals. Due to its sensitivity, simplicity, and short analysis time, this approach is extremely important. The technique of photometrically determining Fe(III) was devised, and the ideal conditions for the formation of a complex of Fe(III) with 2-naphthylcarboxymethylene citrate reagent were explored. The following steps were taken during the photometric determination of Fe (III) using 2-naphthylcarboxymethylcitrate: selection of a light filter, dependence of complex formation on environmental acidity, dependence on the composition of the buffer solution, dependence on the composition of the reagent, field of obedience to Ber's law, the ratio of the component moles of the complex, the Sendal sensitivity, the lowest detection limit of Fe(III), the molar extinction coefficient, and the technique of photometrically determining Fe(III).

Keywords: 2-naphthylcarboxymethylene citrate; Ber's law; buffer solution; sensitivity according to Sendal; molar extinction coefficient; photometric determination; complex formation

Citation: R.S. Jurayev; A.U. Choriev; N.T.Qaxxorov1 Photometric determination of iron(III) with 2-naphthylcarboxymethylene citrate. 2023, volume number, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Received: date

Accepted: date

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

A literature review shows that many organic reagents have been proposed for the determination of iron and other metals, but only a few are highly sensitive and selective. Basically, the sensitive ones have low selectivity, the selective ones have low sensitivity. Therefore, the development of new detection methods for iron detection with high sensitivity and selectivity is an urgent problem and the main goal of our scientific research [1,2].

Currently, photometric and spectrophotometric methods are widely used to determine the amount of a substance. In this work, the works of photometric and spectrophotometric determination of iron(III) are considered.

Photometric methods are widely used in the determination of iron, as well as small amounts of various metals. The advantage of this method is that it is very simple and fast.

This method is widely used to determine the reaction of iron ions in organic reagents. Usually, for the reaction, colored compounds of cationic and anionic ions of iron in solution enter into salt-forming and complex-forming reactions.

Not all organic reagents used for photometric determination of iron have selectivity. Therefore, it is very important to create reagents with high selectivity and analytical ability [1].

Complex compounds of metals with hydroxy acids are of considerable interest for various fields of chemistry and chemical technology. Various hydroxy acids are able to form such complex compounds with more than 60 elements of the periodic system. At present, the study of the properties of this group of compounds is still far from complete. Nevertheless, there is already enough experimental material for some generalizations about the regularities of the complex formation process. In this article, only complex compounds of hydroxy acids of the fatty series are considered. The most typical and widespread representative of such hydroxy acids is tartaric acid; therefore, the main attention is paid to metal tartrate complexes. In some cases, in order to elucidate the influence of the structure of hydroxy acids on the chemical properties of complex compounds, literature data on complexes of citric, trihydroxyglutaric, and some other hydroxy acids are also used.

Even in the last century, it was noticed that the cations of many heavy metals in the presence of tartaric acid lost the ability to precipitate with an alkali in the form of insoluble hydroxides and changed other characteristic analytical properties. Qualitative studies have shown the connection of these phenomena with the hydroxy groups of tartaric acid.

It turned out, in addition, that other organic substances containing hydroxy groups also exhibited a similar masking effect, however, the replacement of hydroxy groups with basic or acidic residues, such as HS⁻, CN⁻, -NH₂, Cl⁻, etc., led to a significant loss of complexing ability. Thus, the special role of hydroxy groups in the process of complex formation was elucidated.

Photometric and spectrophotometric methods, mainly the formation of colored compounds that absorb light in the visible region of the spectrum, are not very specific to fatty hydroxy acids. Photometric measurements were used to study copper, vanadium, chromium, uranium, iron, cobalt, nickel tartrate and nitrate complexes. These works in some cases made it possible not only to determine the composition of complex compounds in solutions, but also to determine the corresponding values of instability constants [3].

Typically, the optical density of a series of solutions with a constant concentration of the central ion and a variable concentration of the hydroxy acid is measured to determine the composition of the compound. The isomolar series method was also used. Both of these options were mainly suitable for determining the composition of complexes under conditions where the acidity of the solution was sufficient to prevent the hydrolysis of the metal salt. An auxiliary complexing agent is sometimes added to the solution to prevent hydrolysis at high pH. As a last resort, for example, sodium salicylate solution was used in the study of tartrate and citrate complexes of iron by the method of isomolar series.

As a result of review of literature materials, it became known that compounds containing oxycarbonic acid and aromatic rings have high analytical ability. Taking this into account, in this work, naphthalene-2-yl 2-chloroacetate was formed as a result of reaction of naphthol-2 with chloroacetyl chloride. The resulting naphthalene-2-yl 2-chloroacetate was subjected to a nucleophilic exchange reaction with the monopotassium salt of citric acid. As a result of the above processes, the reagent 2-naphthylcarboxymethylene citrate (according to systematic nomenclature it is named as follows: 3-hydroxy-3-((2-naphthalen-2-yloxy)-2-oxoethoxy) pentanedioic acid (ChemOffice12)) was formed, and the analytical capabilities of this substance were used for the purpose of photometric determination of Fe(III) [4,5].

2. Experimental

2.1. Materials and measurements.

It is known that organic reagents used in analytical chemistry contain electron-donating groups (-CH₃; -C₂H₅;-OH; -OCH₃, etc.) such reagents are very sensitive, if electron-accepting groups (-SO₃;-NO₂;-COOH; CN⁻; CHO⁻, etc.), then the description of selective influence is improved. If both electron-donor (ED) and electron-acceptor (EA)

groups are introduced into the analytical reagent molecule at the same time, the color of the reagent is clearly visible and the analytical and metrological characteristics of the complex are improved. Such reagents increase in sensitivity and selectivity. In this scientific work, when preparing a new 0.1% solution of 2-naphthylcarboxymethylene citrate, 0.1 g was taken from it, put into a 100 ml measuring flask and prepared by dissolving it in distilled water up to the mark.




Figure 1. Chemical formula of 2-naphthylcarboxymethylene citrate (3-hydroxy-3-((2-naphthalen-2-yl)oxy)-2-oxoethoxy) pentanedioic acid).

Initially, a standard solution of iron(III) was prepared as a working solution. For this $\text{Fe}_2(\text{SO}_4)_3 \cdot 6\text{H}_2\text{O}$ k.t. branded salt was used. 4.5478 g of the calculated amount of salt was weighed on an analytical balance and put into a 1000 ml measuring flask. A 1 mg/ml (1000 $\mu\text{g}/\text{ml}$) solution was prepared by dissolving it in distilled water. Working solutions were prepared by diluting an aliquot of 1mg/ml standard solution before each work.

2.2. Selection of optimal conditions for iron(III) complexation with 2-naphthylcarboxymethylene citrate reagent.

Spectrophotometric determinations are performed under optimal conditions that ensure complete formation of the analytical form in solution and no or minimal deviation from the Bouguer-Lambert-Beer law. The most important of them are: the optimal value of pH, the sufficient amount of the reagent, the selectivity of the analytical (photometric) reaction, and the most favorable conditions for light absorption.

In order to choose the optimal value of pH, the effect of pH on the intensity of the color of the solution at a certain wavelength was studied when the concentrations of the tested substance and reagent were unchanged. In this case, when the reagent is colorless, the area where the absorbance is the largest is considered. In colored solutions, the most favorable conditions correspond to the largest difference between the absorbances of the analytical form and the starting reagents. Under optimal conditions, small changes in pH have virtually no effect on the absorbance of the solution when the absorbance is at its maximum. Photometrically, the pH of the solution being analyzed is kept constant by using appropriate buffer solutions or sufficient amounts of acid or alkali.

The amount of analytical reagent to be added should be sufficient to convert all of the analyte within a given concentration range into analytical form. Adding more reagent does not increase the yield of the reaction product and does not increase the light absorption of the solution.

In spectrophotometric analysis, the solution must remain completely soluble in the entire range of detectable concentrations. If this condition is not met, it is necessary to use lower concentrations or to use protective substances that prevent the formation of a solid phase. Sometimes it is necessary to change the entire spectrophotometric detection scheme.

2.3. Selection of optimal light filter for Fe(III) complexation with 2-naphthylcarboxymethylene citrate reagent.

It is known that each substance by its nature absorbs light of a certain wavelength, taking into account that the highest absorption area of the complex of Iron(III) with 2-naphthylcarboxymethylene citrate was determined as follows: 5.0 ml of buffer solution, 1.0 ml of 0.1% 2-naphthylcarboxymethylene citrate solution, and 1.0 ml of 20 µg/ml Fe(III) solution were added to a 25 ml volumetric flask and filled with distilled water up to the mark of the flask. . The optical density of the resulting complex was measured in a concentration photoelectrocolorimeter CPhC-3, in different light filters with a light absorption thickness of 1=3.0 cm. A solution containing all components except iron (III) was used as a reference solution. The measurement results are presented in Table 1, Figure 2.

Table 1. Dependence of the optical density of the iron(III) complex with 2-naphthylcarboxymethylene citrate on the light filter (n=3).

№ Nur filtri	Color	λ , nm	\bar{A}
1	Purple	315	0.000
2	Blue	364	0.000
3	Green-blue	400	0.000
4	Blue green	440	0.028
5	Green	490	0.033
6	Yellow-green	540	0.037
7	Yellow	590	0.042
8	Red	670	0.153
9	Dark red	750	0.115
10	Infrared	870	0.035
11	Infrared	980	-




Figure 2. Dependence of the optical density of the FeR complex on the light filter .

As can be seen from the obtained results, the complex compound 8-light filter showed high optical density at $\lambda_{max} = 670nm$. Further work was carried out at $\lambda_{max} = 670nm$.

2.4. Dependence of optical density value of Fe(III) complex with 2-naphthylcarboxymethylene citrate reagent on solution medium (pH).

Taking into account that one of the important conditions for the reaction is its environment, universal buffer solutions with different pH values were prepared for choosing optimal conditions for the complex combination of Iron(III) with 2-naphthylcarboxymethylene citrate.

Detection method:

To do this, add 5.0 ml of a universal buffer solution with an pH of 2.0 to -11.10, 1.0 ml of a 0.1% 2-naphthylcarboxymethylene citrate reagent solution, 20 µg/ml of Fe(III) to a 25 ml volumetric flask.) solution was added to 1.0 ml and the flask was filled with distilled water up to the mark. The optical densities of complex compound solutions were measured in CPhC-3 cuvette at $\lambda_{max} = 670 \text{ nm}$ and absorption thickness $l = 3.0 \text{ cm}$. Table 2 shows the obtained results.

Table 2. Dependence of the optical density of the complex compound (FeR) on the solution medium pH. $(T_{Fe^{3+}} = 20 \text{ µg/ml}, n=3)$.

pH	2.18	2.80	3.46	4.02	4.52	5.48	6.54	7.21	8.01	9.04	10.55	11.27	11.80	12.49
\bar{A}	0	0	0.012	0.013	0.015	0.017	0.032	0.035	0.073	0.088	0.11	0.175	0.183	0.163

As can be seen from Table 2, the highest optical density of the complex compound was observed in the range of pH=10.55-12.0, and pH=11.80 was chosen as the optimal environment, because the optical density at this pH has the maximum analytical signal. A buffer solution with a pH of 11.80 was used in further research.

2.5. Dependence of the optical density of the complex compound on the composition of the buffer solution.

Universal and with sodium citrate buffer solutions with pH=11.80 were prepared to study the dependence of buffer solution composition on the components of the main reaction ($Fe^{3+} + 2\text{-naphthylcarboxymethylene citrate}$).

Detection method:

for the preparation of photometric solutions, 5.0 ml of various buffer solutions with pH=11.80, 1.0 ml of 0.1% aqueous solution of 2-naphthylcarboxymethylene citrate, 20 µg/ml of Add 1.0 ml of Fe(III) solution and fill the flask up to the mark with distilled water. Optical densities were measured against reference solution in CPhC-3, $l = 3.0 \text{ cm}$ cuvette. The obtained results are presented in Table 3.

Table 3. Choosing the optimal buffer solution (n=3).

The name of the buffer solution	The composition of the buffer solution	pH	\bar{A}_{total}
Universal	$(H_3PO_4 + CH_3COOH + H_3BO_3 + NaOH)$	11.80	0.183
Na- citrate	$(HCl + NaH_2C_6H_5O_7)$	11.80	0.046

As can be seen from the obtained experimental results, when the universal buffer solution was used, the complex compound solution had the maximum optical density. A universal buffer solution with pH=11.80 was used in further research.

2.6. Study of the stability of Fe(III) complex with 2-naphthylcarboxymethylene citrate reagent over time.

To determine the stability of the complex, the stability of the optical density of the solution with respect to time was studied.

Detection method:

Add 5.0 ml of universal buffer solution (pH=11,80), 1.0 ml of 0.1% 2-naphthylcarboxymethylene citrate reagent, 1.0 ml of 20 µg/ml Fe(III) solution to a 25 ml volumetric flask. , the flask was filled to the mark with distilled water. The optical density of the resulting complex was measured against a reference solution at certain time intervals. The obtained results are presented in Table 4.

Table 4. Stability of optical density of complex (FeR) compound with respect to time ($\lambda_{\max}=670$ nm, $l=3,0$ cm, $n=3$).

t_{\min}	1	5	10	15	20	30	40	50
$\bar{A}_{урт}$	0.275	0.275	0.274	0.274	0.273	0.273	0.273	0.272
t_{\min}	60	70	90	100	120	140	160	180
$\bar{A}_{урт}$	0.272	0.271	0.271	0.270	0.270	0.270	0.269	0.269

The experimental results show that (Table 4) the optical density value of the complex compound did not change much for 2 hours, and then a slight decrease was observed. It can be concluded that this time interval is sufficient to perform the analysis.

2.7. Study of the pouring order of the structural components of the complex compound (FeR).

Taking into account that the yield of the reaction also depends on the order of pouring the components, the solutions were prepared by the above-mentioned method, and several experiments were carried out by changing the order of pouring the components. The measurement results are presented in Table 5.

Table 5. The results of studying the order of pouring components ($n=3$).

№	Casting Order	\bar{A}_{total}
1	buffer-reagent- Iron -dist. water	0.455
2	buffer - reagent - dist. water - Iron	0.421
3	buffer – Iron - dist. water – reagent	0.406
4	buffer - Iron – reagent - dist. water	0.482
5	reagent - Iron - buffer - dist. water	0.517
6	reagent - buffer - Iron - dist. water	0.468
7	reagent - dist. water - buffer - Iron	0.479
8	Iron - buffer - reagent - dist. Water	0.561
9	Iron - reagent - buffer - dist. Water	0.572
10	Iron - dist. water - buffer – reagent	0.559
11	Iron - reagent - dist. water – buffer	0.545

It can be concluded from the obtained results that the order of pouring of components is quite important. In further research work, the 9th casting order was chosen. Carrying out the reaction in this order ensures that the complex exhibits maximum optical density.

2.8. Dependence of the optical density of the complex compound on the amount of added reagent.

In practical studies, an excess of reagent is usually taken to fully bind the metal to the complex. For this purpose, to study the dependence of the optical density of the complex compound on the amount of added reagent, photometric solutions were prepared in 25 ml measuring flasks.

Detection method:

1.0 ml of a 20 µg/ml Fe(III) solution in 25 ml volumetric flasks, variable amounts from 0.20 to 4.00 ml of 0.1% aqueous solution of 2-naphthylcarboxymethylene citrate, 5.0 ml (pH=11,80) and mixed with distilled water up to the mark of the flask. The optical density of the complex compound was measured in CPhC-3, Nf №4, $l = 3.0$ cm against reference solution. The obtained results are presented in Table 6 and Figure 3.

Table 6. Dependence of the optical density of the color complex of iron (III) with 2-naphthylcarboxymethylene citrate reagent on the amount of added reagent (n=3).

V_R, ml	0.25	0.5	0.75	1.00	1.25	1.50	1.75	2.00	2.25	2.50
A	0.095	0.150	0.184	0.247	0.293	0.338	0.369	0.404	0.410	0.421

V_R, ml	2.75	3.00
A	0.419	0.419




Figure 3. Graph of dependence of the optical density of the complex compound (FeR) on the amount of added reagent.

The obtained results show that 2.5 ml of 0.1% reagent, i.e. 2.5 ml of aqueous solution of 2-naphthylcarboxymethylene citrate, is sufficient for complete binding of 20 $\mu\text{g/ml}$ of Iron(III) to the complex. In subsequent work, a 0.1% solution of the reagent was prepared and used in a volume of 2.5 ml.

2.9. Dependence of the complex of iron(III) with 2-naphthylcarboxymethylene citrate on the amount of elements (subordination to the Bouguer-Lambert-Beer law).

The behavior of the solution of the complex solution of iron(III) with 2-naphthylcarboxymethylene citrate was studied under selected optimal conditions.

Detection method:

25 mL volumetric flasks were filled with varying amounts of a 20 $\mu\text{g/mL}$ solution of Fe(III), 2.5 mL of a 0.1% 2-naphthylcarboxymethylene citrate solution, and 5.0 mL of a universal buffer solution of $rN=4.52$ up to the mark of the flask. diluted with distilled water. By mixing the solutions, their optical density ($\lambda_{\text{max}}=670 \text{ nm}$, $l=3.0\text{cm}$) was measured against the reference solution. The obtained results are presented in Table 7 and Figure 4.

Table 7. Dependence of the optical density of the complex compound (FeR) on the amount of added Iron(III) (Obedience to the law of Ber). ($C_{Fe^{3+}}=20,0 \mu\text{g/ml}$, $l=3,0 \text{ cm}$, $n=3$).

№	$V_{Fe^{3+}} \text{ mL}$	The amount of iron (III). μg	\bar{A}_{total}
1	0.10	2.0	0.060
2	0.25	5.0	0.114
3	0.50	10.0	0.155
4	0.75	15.0	0.203
5	1.00	20.0	0.286
6	1.25	25.0	0.335
7	1.50	30.0	0.400
8	1.75	35.0	0.448
9	2.00	40.0	0.506
10	2.25	45.0	0.592
11	2.50	50.0	0.597
12	2.75	55.0	0.604
13	3.00	60.0	0.616




Figure 4. The graph of optical density depending on the amount of added Iron(III).

As can be seen from the obtained results, the field of obedience to Bouguer-Lambert-Beer law was observed in the range of 2-45 $\mu\text{g}/25\text{ml}$ in 25ml solution. At higher concentrations, there was a deviation from the linear relationship, that is, the range from 2 μg to 45 $\mu\text{g}/25 \text{ ml}$ allows the quantitative determination of Fe(III).

3. Reaction results and discussion of results

3.1. Spectral description of iron(III) complex with 2-naphthylcarboxymethylene citrate reagent.

Absorption spectra of the complex formed with 2-naphthylcarboxymethylene citrate reagent and Iron(III) were obtained under selected optimal conditions.

Detection method:

1.0 ml of 45.0 $\mu\text{g/ml}$ Fe(III) solution, 2.5 ml of 0.1% aqueous solution of 2-naphthylcarboxymethylene citrate reagent and 5.0 ml of pH 4.52 buffer were added to a

25 ml volumetric flask. from the solution, diluted with distilled water up to the mark of the flask and mixed. The absorption spectrum of the produced complex compound was measured in the spectrophotometer "SF-46" in a quartz cuvette with an absorption thickness of $l = 1.0 \text{ cm}$ compared to the reference solution. The absorption spectrum of the reagent was obtained against distilled water. The results are presented in Figure 6. According to the absorption spectrum presented in Figure 6, the maximum light absorption area of the complex of 2-naphthylcarboxymethylene citrate reagent with Iron(III) is located at $\lambda_{\text{comp}}=675 \text{ nm}$, the maximum light absorption area of the 2-naphthylcarboxymethylene citrate reagent was observed in the field of shorter spectral waves, i.e. at $\lambda_{\text{reagent}}= 380 \text{ nm}$.

Using the value of the maximum optical density of the complex compound (in the area of $\lambda=675 \text{ nm}$), the apparent molar extinction coefficient (ε) was determined by the following formula:

$$\varepsilon = A/C \cdot l$$

Here: ε - molar extinction coefficient of light; C- Iron(III) concentration (mol/l); l - thickness of absorbent layer (cm); A- the optical density of a complex compound measured relative to a reference solution.

The spectral characterization of the complex and reagent is depicted in Table 8 and Figure 6.

The Cendel sensitivity of the method is calculated by the formula:

$$C.S. = \frac{Q \cdot l \cdot 0,001}{A \cdot 25} = \frac{45,0 \cdot 1,0 \cdot 0,001}{0,345 \cdot 25,0} = 0,0052 \mu\text{g}/\text{cm}^2$$

Cendel sensitivity was found to be $0.0052 \mu\text{g}/\text{cm}^2$.

Table 8. Spectral description of the complex of iron(III) with 2-naphthylcarboxymethylene citrate reagent and the reagent ($l \approx 3.0 \text{ cm}$, $C_{\text{Fe}^{3+}} = 45 \mu\text{g}/\text{ml}$).

Complex color	pH	$\lambda_{\text{MeR}}, \text{nm}$	$\lambda_{\text{HR}}, \text{nm}$	$\Delta\lambda$	$C_{\text{Fe}^{3+}}, \mu\text{g}$	$C_{\text{Fe}^{3+}}, \text{mol/L}$	\bar{A}	Sensitivity according to Sendel is $\mu\text{g}/\text{cm}^2$
Green	4.52	675	380	295	45.0	$3.22 \cdot 10^{-5}$	0.345	0.0052




Figure 5. Absorption spectrum of iron(III) complex with 2-naphthylcarboxymethylene citrate reagent.

The obtained results show that the reaction has high contrast ($\Delta\lambda = 295 \text{ nm}$) and sensitivity ($C.S.=5.2 \cdot 10^{-3}$).

3.2. Determination of the ratio of component moles of the complex formed by iron(III) with 2-naphthylcarboxymethylene citrate reagent.

In the complex formed by iron(III) with 2-naphthylcarboxymethylene citrate reagent, the molar ratios of components were determined by the method of isomolar series and Asmus' straight line method.

3.2.1. Determination of the composition of the complex using the method of isomolar series.

Equal concentration solutions of Iron(III) and 2-naphthylcarboxymethylene citrate reagent were used to determine the molar ratio of iron(III) in the complex formed by 2-naphthylcarboxymethylene citrate reagent by isomolar series method:

$$C_{\text{Fe}^{3+}} = C_{\text{HR}} = 3,22 \cdot 10^{-5} \text{ mol/l}$$

Detection method:

A series of solutions was prepared in 25 ml volumetric flasks. To do this, add variable amounts of iron(III) solutions (from 9.0 to 1.0 ml) to each of them, then add variable amounts of 2-naphthylcarboxymethylene citrate reagent solution (from 1.0 ml to 9.0 ml) and pH Add 5.0 mL of 11,80 universal buffer solution to each, dilute to the mark with distilled water, and mix. The optical density of the prepared solutions was measured in CPhC-3 in Nf № 8 in cuvettes with an absorption thickness of $l = 3.0 \text{ cm}$ compared to a reference solution. The obtained results are presented in Table 9 and Figure 6.

Table 9. The results of the study of the composition of the complex compound (FeR) using the isomolar series method. ($C_{\text{Fe}^{3+}} = C_{\text{HR}} = 3,22 \cdot 10^{-5} \text{ M}$, $l = 3,0 \text{ cm}$, $n=3$).

№	Obtained Fe(III), mL	Obtained V_{HR} , mL	Buffer solution, mL	\bar{A}
1	9.00	1.00	5.00	0.115
2	8.00	2.00	5.00	0.202
3	7.00	3.00	5.00	0.286
4	6.00	4.00	5.00	0.365
5	5.00	5.00	5.00	0.440
6	4.00	6.00	5.00	0.545
7	3.00	7.00	5.00	0.612
8	2.50	7.50	5.00	0.635
9	2.00	8.00	5.00	0.568
10	1.00	9.00	5.00	0.455




Figure 6. The curve for determining the composition of the complex compound (FeR) using the isomolar series method.

As can be seen from this graph, the composition of the complex formed by Iron (III) and 2-naphthylcarboxymethylene citrate reagent corresponded to Fe:R=1:3 molar ratio.

3.2.2. Determining the composition of the complex using the linear Asmus method.

In order to graphically determine the stoichiometric ratios of the interacting components, a $(1/V)n = l/A$ dependence graph was created. This relationship only represents a straight line for the true value of n.

Detection method:

Add 1.0 ml of an equal concentration of iron(III) solution to 25 ml volumetric flasks, add variable amounts of an aqueous solution of 2-naphthylcarboxymethylene citrate reagent and 5.0 ml each of a universal buffer with pH = 4.52, and add distilled water to the mark of the flask was diluted and mixed with. Optical densities of prepared photometric solutions in CPhC-3 N.f. in №8, light absorption was measured against a reference solution in cuvettes with a thickness of $l = 3.0$ cm. The obtained results are presented in Table 10 and Figure 7.

Table 10. Determination of the composition of iron (III) complex formed with 2-naphthylcarboxymethylene citrate reagent using Asmus' straight line method. ($C_{Fe^{3+}}=C_{HR}=3,22 \cdot 10^{-5}$ M, $l = 3,0$ cm, $n=3$).

№	$V_{HR, ml}$	\bar{A}	$\frac{l}{\bar{A}}$	$\frac{1}{V}$	$(\frac{1}{V})^2$	$(\frac{1}{V})^3$
1	0.2	0.060	50.00	5.0	25.0	125.0
2	0.4	0.121	24.79	2.5	6.25	15.63
3	0.6	0.190	15.80	1.67	2.78	4.53
4	0.8	0.274	10.95	1.25	1.56	1.95
5	1.0	0.369	8.13	1.00	1.00	1.00
6	1.2	0.400	7.50	0.83	0.69	0.57
7	1.4	0.410	7.32	0.71	0.51	0.36
8	1.6	0.494	6.07	0.63	0.39	0.25
9	1.8	0.535	5.60	0.56	0.31	0.17
10	2.0	0.570	5.26	0.5	0.25	0.125




Figure 7. Graph of determination of the composition of the complex of iron (III) with 2-naphthylcarboxymethylene citrate using Asmus' straight line method.

The obtained results show that the straight line corresponded to the ratio of 1:3 moles. Therefore, both methods confirmed each other and showed that the composition was Fe:HR=1:3.

Based on the conclusions obtained from the above methods, the complex formation reaction occurs based on the reaction equation shown in Figure 8.




Figure 8. Fe:HR=1:3 ratio complex formation reaction equation.

4. Conclusions

This article proves that organic chemistry and analytical chemistry are inextricably linked. Studies on the photometric determination of a very small amount of Fe(III) ion with one of the new organic synthesis products A were carried out. It was proved that A can be used in the determination of Fe(III) ion. We are continuing our research to identify other d-metal ions similar to the Fe(III) ion by this method.

Acknowledgments: The authors wish to acknowledge Karshi State University, Karshi, Uzbekistan, the Ministry of Higher and Secondary Specialized Education of the Republic of Uzbekistan and the Ministry of Innovative Development of the Republic of Uzbekistan, Tashkent, Uzbekistan.

References

1. Segoviano-Garfias, J.J.N.; Zanon, G.A.; Ávila-Ramos, F.; Bivián-Castro, E.Y.; Rubio-Jiménez, C.A. Spectrophotometric Determination of Formation Constants of Iron(III) Complexes with Several Ligands. *Chemistry* **2022**, *4*, 701–716.

2. Ueno, T.; Miyano, K.; Hamada, D.; Ono, H.; Fujinami, T.; Matsumoto, N.; Sunatsuki, Y. Abrupt Spin Transition and Chiral Hydrogen-Bonded One-Dimensional Structure of Iron(III) Complex [FeIII(Him)2(hapen)]SbF6 (Him = imidazole, H2hapen = N,N'-bis(2-hydroxyacetophenylidene)ethylenediamine). *Magnetochemistry* **2015**, *1*, 72–82,
3. L. Dobb in, *J. Am. Chem.Soc*, *41*, 934 (1919); *C*, 1919, III, 702
4. Sadikova S.B.; Abdushukurov, A.K.; Choriyeu, A.U. Chloroacetylation of hydroquinone and its Esters with Lewis acids. *Univers. Chem. Biol.* **2019**, *5*, 52–56.
5. Choriev, A.U.; Jurayev, R.S.; Abdushukurov, A.K.; Abdullayev, M.G. Synthesis of 2-Izopropyl-5-methylphenylcarboxymethylen Tartrate. *2023*, *37*, 57.
6. M. Kass, A. Ivaska / *Talanta* *58* (2002) 1131-1137
7. Miranda, J.C.; Kamogawa, M.Y.; Reis, B.F. Development of a portable setup and a multicommutated flow analysis procedure for the photometric determination of Fe(III) and Fe(II) in fresh water. *Sensors Actuators B: Chem.* *2015*, *207*, 811–818, <https://doi.org/10.1016/j.snb.2014.10.122>.
8. Karthikeyan, S.; Hashigaya, S.; Kajiya, T.; Hirata, S. Determination of trace amounts of phosphate by flow-injection photometry. *Anal. Bioanal. Chem.* *2004*, *378*, 1842–1846.
9. Hanrahan, G.; Gledhill, M.; Fletcher, P.J.; Worsfold, P.J. High temporal resolution field monitoring of phosphate in the River Frome using flow injection with diode array detection. *Anal. Chim. Acta* *2001*, *440*, 55–62, [https://doi.org/10.1016/s0003-2670\(00\)01395-7](https://doi.org/10.1016/s0003-2670(00)01395-7).
10. Mendham J, Denney R C, Barnes J D & Thomas M, *Statistics:Introduction to Chemometrics*, in *Vogel's Textbook of Quantitative Chemical Analysis*, 6th edn (Pearson Education, Singapore) 2002, 137.
11. G.D.Christian, *Analytical Chemistry*, 4th edn, Wiley, Chichester,1986
12. D.C.Harris, *Quantitative Chemical Analysis*, Freeman, San Fransisco, 1982