

Proceeding Paper

Grassland Crops as Drivers for the Improvement of Soil Fertility [†]

Tomás Cavaco ^{1,2,*}, Maria Odete Torres ² and Jorge M. S. Faria ^{1,3}

¹ INIAV, I.P., National Institute for Agrarian and Veterinary Research, Quinta do Marquês, 2780-159 Oeiras, Portugal; fariajms@gmail.com

² Instituto Superior de Agronomia (ISA), Universidade de Lisboa, 1349-107 Lisboa, Portugal; motorres@isa.ulisboa.pt

³ MED, Mediterranean Institute for Agriculture, Environment and Development & CHANGE—Global Change and Sustainability Institute, Institute for Advanced Studies and Research, Évora University, Pólo da Mitra, Ap. 94, 7006-554 Évora, Portugal

* Correspondence: tomasfcavaco@gmail.com

[†] Presented at the 3rd International Electronic Conference on Agronomy, 15–30 October 2023; Available online: <https://iecag2023.sciforum.net/>.

Abstract: Managing soil fertility is vital for agriculture. However, modern farming excessively relies on mineral fertilizers, which lessens profit and endangers ecosystem health. Grasslands made up of Poaceae and Fabaceae, including woody species, offer feed for livestock, lowers farmers' economic risks, and conserve resources. Grassland crops can enhance soil fertility in a more sustainable way than mineral fertilization. To counter fertilizer-driven soil decline, permanent grasslands or crop rotations are effective. Also, grassland soils generally contain more nitrogen, potassium, organic matter, and less phosphorus than cropland soils. They additionally enhance soil's physical and biological parameters, limiting erosion while elevating biodiversity. This work focuses on the benefits of grasslands towards crop production, reviewing their influence on soil fertility parameters that boost soil health.

Keywords: forages; grassland crops; pastures; soil fertility; soil health; sustainable agriculture

1. Introduction

Modern agriculture faces several challenges in producing food to support an increasing world population, while in need of rapidly adapting to climate change. Soil fertility is one of the key aspects of agricultural management. Nevertheless, enhancing crop yield has driven farmers to excessive utilization of inorganic fertilizers, with heavy damage to biodiversity and posing environmental and human health risks [1].

Agroecological practices can increase the sustainability of modern agriculture by adapting ecological principles and traditional practices, which minimizes the impact on the environment and boosts soil fertility and contributes to the management of crop protection [2,3].

Soil fertility emerges from the interaction of biological, chemical, and physical processes [4]. Soil microbial and mesofauna communities are extremely relevant for plant nutrition and defense [5]. Arbuscular mycorrhizal fungi (AMF), plant growth-promoting rhizobacteria (PGPR), and beneficial nematodes are examples of important contributors to the enhancement of soil biological fertility [6–8]. Soil chemical fertility influences and is influenced by the other components and is related to the concentration of inorganic elements and their bioavailability. Key parameters in this context include the cation exchange capacity (CEC), soil organic matter (SOM) or content (SOC), ratios of macro- and micronutrients, and the pH that can influence their availability to plants [9]. Lastly, soil

Citation: Cavaco, T.; Torres, M.O.; Faria, J.M.S. Grassland Crops as Drivers for the Improvement of Soil Fertility. *Biol. Life Sci. Forum* **2023**, *27*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

physical fertility relates to porosity, structure, and drainage that influence water availability and aeration [10].

Grasslands, encompassing both natural expanses and cultivated areas, predominantly comprise members of the Poaceae and Fabaceae families. These ecosystems stand as primary sources of sustenance for numerous livestock, underpinning global food security [11]. The use of pastures and forages has been associated with several ecosystem services that enhance the farm's long-term sustainability. These benefits include the promotion of soil fertility, carbon storage, water regulation, biodiversity, pollination, and pest control [12,13]. Hence, the use of these crops can be considered an agroecological measure that ensures sustainable food production.

In this study, a review of the beneficial role of grasslands on soil chemical fertility is presented comparing the effects of implementing croplands versus grasslands.

2. Bibliographic Sources

The research was performed using the Web of Science search engine (<https://www.webofknowledge.com>, last accessed July 2023), in all available databases, on published works from 2018 to 2023, using the topics "grassland crops" and "soil fertility". A total of 17 works were retrieved concerning soil chemical fertility parameters measured in croplands in comparison to grasslands (Figure 1). Works were mainly published in journals specialized in agriculture, environmental sciences, biodiversity conservation, plant sciences, and chemistry. These selected articles were cited 108 times, thus the average number of citations per article is approximately 6.35.

To assess the beneficial influence of grassland crops, when compared to croplands, on soil chemical fertility, a total of 10 parameters were considered. For this purpose, percentual relative differences of mean values were calculated following this formula:

$$\text{Relative Differences (\%)} = [(GL-CL)/GL] \times 100 \quad (1)$$

where GL is the grassland mean value and CL is the cropland mean value.




Figure 1. The cumulative number of published articles (a) and respective cumulative number of citations (b), between January of 2018 and July of 2023.

3. The Influence of Grasslands on Soil Chemical Fertility

Differences between conventional crop fields and grasslands were reported for the selected soil chemical parameters. Several reports suggest a tendency for increased levels of soil organic carbon (SOC) or soil organic matter (SOM) through forage or pasture cultivation. Likewise, grassland soils frequently exhibited elevated nitrogen and potassium content, indicating improved nutrient conditions. Similarly, the C/N ratio tended to be slightly higher in pastures and forage soils, contributing to a slower and sustained organic matter mineralization.

vs.							
Potato (<i>Solanum tuberosum</i>)							
Guinea grass (<i>Megathyrsus maximus</i>), Sedge (<i>Cyperus</i> spp.), Waterleaf (<i>Talinum fruticosum</i>), and Paspalum (<i>Paspalum decumbens</i>)							
0–30		+6	+2			–107	–2 [24]
vs.							
Cassava (<i>Manihot esculenta</i>), Peanut (<i>Arachis hypogea</i>), Maize (<i>Z. mays</i>), and Cowpea (<i>Vigna unguiculata</i>)							
Alfalfa (<i>M. sativa</i>), Sorghum (<i>S. bicolor</i>), Winter wheat (<i>Triticum aestivum</i>), Chickpea (<i>Cicer arietinum</i>), and Spring barley (<i>Hordeum vulgare</i>)							
not re- ferred	+60		+64 ³	+62	+76		[25]
vs.							
Winter wheat (<i>T. aestivum</i>), Chickpea (<i>C. arietinum</i>), and Spring barley (<i>H. vulgare</i>)							
Perennial ryegrass (<i>L. perenne</i>) and Clover (<i>Trifolium</i> spp.)							
vs.							
0–30		+45		+43			[26]
Potato (<i>S. tuberosum</i>), Maize (<i>Z. mays</i>), and Winter wheat (<i>Triticum aestivum</i>)							
Grassland (sp. not referred)							
0–15	+46						
vs.							
15–30	+42						[27]
Rice (<i>Oryza sativa</i>), Maize (<i>Z. mays</i>), and Okra (<i>Abelmoschus esculentus</i>)							
30–45	+13						
Grassland (sp. not referred)							
vs.							
0–50						–3 ⁵	[28]
Cropland (sp. not referred)							
Blue grama (<i>Bouteloua gracilis</i>), Buffalo grass (<i>B. dactyloides</i>), and Little barley (<i>H. pusillum</i>)							
0–8	+67					+2	
8–15	+37					+6	[29]
16–23	+37					–1	
vs.							
23–30	+28					0	
Wheat (<i>T. aestivum</i>)							
Grassland (sp. not referred)							
vs.							
0–30	+40		+32			–233	+8 [30]
Cropland (sp. not referred)							

¹ SOM—soil organic matter; SOC—soil organic carbon; C/N—Carbon/Nitrogen ratio; TN—total nitrogen (N); TP—total phosphorus (P); TK—total potassium (K); AN—available nitrogen; AP—available phosphorus; AK—available potassium. ² The values concern average plant derived C_{org} inputs per year. ³ The values concern average total inputs of N, P, and K. ⁴ This land use concerns an area with very sparse vegetation with less than 5% ground cover. ⁵ The authors mention that the pH of both land-use types showed a bimodal distribution, with peaks at pH 6.1 and 7.3 (cropland) and pH 5.0 and 7.1 (grassland).

The beneficial role of grasslands on soil fertility is strongly connected to the agricultural practices used and the crops grown. In fact, the precise net impact of grasslands on the overall improvement of soil fertility may be undervalued due to the difficulty in adequately implement control plots for diverse land uses with varying levels of mineral fertilizer application.

Generally, great focus has been given to the topsoil layer, given its crucial role in plant nutrition [9]. This soil layer is the most bioactive part of agricultural soils since beneficial microbial communities, that can fix atmospheric N₂ and solubilize phosphorus, are deeply dependent on the proximity to the plant's roots [31,32]. Furthermore, the impact

of land use is prone to diminish with the increase of soil depth, putatively related to the volume explored by the root system of the crops. On the other hand, root architecture can substantially alter the depth of the soil layers that can be influenced [33]. Therefore, the traditional combination of grasses and legumes is a proficient approach for more comprehensive soil volume exploration, with the added benefit of enhancing rhizosphere microbial activity [34].

4. Conclusions

The study of the beneficial role of grasslands for soil fertility can reveal novel means to improve food sustainably. Several fertility parameters were found to be greater in pastures and forages than in croplands. Switching to an eco-friendlier crop production system supported by the integration of grasslands into sustainable agricultural practices may improve crop yields, soil health, and biodiversity and provide other benefits in the long term.

Author Contributions: Conceptualization, T.C., M.O.T. and J.M.S.F.; methodology, T.C. and J.M.S.F.; formal analysis, T.C.; investigation, T.C.; data curation, T.C.; writing—original draft preparation, T.C.; writing—review and editing, T.C., M.O.T. and J.M.S.F. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement:

Informed Consent Statement:

Data Availability Statement:

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Cordell, D.; Drangert, J.-O.; White, S. The Story of Phosphorus: Global Food Security and Food for Thought. *Glob. Environ. Change* **2009**, *19*, 292–305.
2. Gargano, G.; Licciardo, F.; Verrascina, M.; Zanetti, B. The Agroecological Approach as a Model for Multifunctional Agriculture and Farming towards the European Green Deal 2030—Some Evidence from the Italian Experience. *Sustainability* **2021**, *13*, 2215.
3. Ros, G.H.; Verweij, S.E.; Janssen, S.J.C.; De Haan, J.; Fujita, Y. An Open Soil Health Assessment Framework Facilitating Sustainable Soil Management. *Environ. Sci. Technol.* **2022**, *56*, 17375–17384.
4. Abbott, L.K.; Murphy, D. V What Is Soil Biological Fertility? In *Soil Biological Fertility: A Key to Sustainable Land Use in Agriculture*; Springer: Berlin/Heidelberg, Germany, 2007; pp. 1–15.
5. Larkin, R.P. Soil Health Paradigms and Implications for Disease Management. *Annu. Rev. Phytopathol.* **2015**, *53*, 199–221.
6. Faria, J.M.S.; Barrulas, P.; Pinto, A.P.; Brito, I.; Teixeira, D.M. Mycorrhizal Colonization of Wheat by Intact Extraradical Mycelium of Mn-Tolerant Native Plants Induces Different Biochemical Mechanisms of Protection. *Plants* **2023**, *12*, 2091.
7. Jha, C.K.; Saraf, M. Plant Growth Promoting Rhizobacteria (PGPR): A Review. *J. Agric. Res. Dev.* **2015**, *5*, 108–119.
8. Pires, D.; Orlando, V.; Collett, R.L.; Moreira, D.; Costa, S.R.; Inácio, M.L. Linking Nematode Communities and Soil Health under Climate Change. *Sustainability* **2023**, *15*, 11747.
9. Weil, R.R.; Brady, N.C. *The Nature and Properties of Soils*, ed.; Pearson: Columbus, OH, USA, 2016; Volume 910.
10. Rasool, R.; Kukal, S.S.; Hira, G.S. Soil Physical Fertility and Crop Performance as Affected by Long Term Application of FYM and Inorganic Fertilizers in Rice–Wheat System. *Soil Tillage Res.* **2007**, *96*, 64–72. <https://doi.org/10.1016/j.still.2007.02.011>.
11. O'Mara, F.P. The Role of Grasslands in Food Security and Climate Change. *Ann. Bot.* **2012**, *110*, 1263–1270.
12. Tamburini, G.; Aguilera, G.; Öckinger, E. Grasslands Enhance Ecosystem Service Multifunctionality above and Below-Ground in Agricultural Landscapes. *J. Appl. Ecol.* **2022**, *59*, 3061–3071. <https://doi.org/10.1111/1365-2664.14302>.
13. Dondini, M.; Martin, M.; De Camillis, C.; Uwizeye, A.; Soussana, J.-F.; Robinson, T.; Steinfeld, H. Global Assessment of Soil Carbon in Grasslands. In *From Current Stock Estimates to Sequestration Potential*; FAO Animal Production and Health Paper, 187; FAO: Rome, Italy, 2023.
14. Cantú Silva, I.; Yáñez Díaz, M.I. Effect of the Change of Land Use on the Contents of Soil Organic Carbon and Nitrogen. *Rev. Mex. Cienc. For.* **2018**, *9*, 122–151.
15. Han, D.; Yu, P.; Zhu, X.; Sun, T.; Jia, H. Relationships between soil physiochemical properties and soil respiration under different land uses in an alpine grassland during freeze-thaw period. *Fresenius Environ. Bull.* **2018**, *27*, 7364–7373.
16. Kharal, S.; Khanal, B.R.; Panday, D. Assessment of Soil Fertility under Different Land-Use Systems in Dhading District of Nepal. *Soil Syst.* **2018**, *2*, 57. <https://doi.org/10.3390/soilsystems2040057>.

17. Rolando, J.L.; Dubeux, J.C.B.; Ramirez, D.A.; Ruiz-Moreno, M.; Turin, C.; Mares, V.; Sollenberger, L.E.; Quiroz, R. Land Use Effects on Soil Fertility and Nutrient Cycling in the Peruvian High-Andean Puna Grasslands. *Soil Sci. Soc. Am. J.* **2018**, *82*, 463–474. <https://doi.org/10.2136/sssaj2017.09.0309>.
18. Zhang, Q.; Shao, M. Soil Fertility Increases Rapidly during the 6-10 Yr Following Conversion of Cropland to Grassland in China's Loess Plateau Region. *Can. J. Soil Sci.* **2018**, *98*, 531–541. <https://doi.org/10.1139/cjss-2017-0107>.
19. Hu, T.; Taghizadeh-Toosi, A.; Olesen, J.E.; Jensen, M.L.; Sørensen, P.; Christensen, B.T. Converting Temperate Long-Term Arable Land into Semi-Natural Grassland: Decadal-Scale Changes in Topsoil C, N, 13 C and 15 N Contents. *Eur. J. Soil Sci.* **2019**, *70*, 350–360. <https://doi.org/10.1111/ejss.12738>.
20. Pantoja, J.C.M.; Campos, M.C.C.; Lima, A.F.L. de; Cunha, J.M. da; Simões, E.L.; Oliveira, I.A. de; Silva, L.S. Multivariate Analysis in the Evaluation of Soil Attributes in Areas under Different Uses in the Region of Humaitá, AM. *Rev. Ambiente Água* **2019**, *14*.
21. Damene, S.; Bahir, A.; Villamor, G.B. The Role of Chomo Grass (*Brachiaria humidicola*) and Exlosures in Restoring Soil Organic Matter, Total Nitrogen, and Associated Functions in Degraded Lands in Ethiopia. *Reg. Environ. Change* **2020**, *20*, 1–13.
22. Fernández, R.; Ezequiel Furch, N.; Bissolino, M.; Frasier, I.; Daniel Scherger, E.; Raul Quiroga, A. Effect of perennial pastures in physical and biological fertility in Mollisols of the semiarid Pampas region. *Cienc. Del Suelo* **2020**, *38*.
23. Jacobs, A.; Poeplau, C.; Weiser, C.; Fahrion-Nitschke, A.; Don, A. Exports and Inputs of Organic Carbon on Agricultural Soils in Germany. *Nutr. Cycl. Agroecosyst.* **2020**, *118*, 249–271. <https://doi.org/10.1007/s10705-020-10087-5>.
24. Nyameasem, J.K.; Reinsch, T.; Taube, F.; Yaw Fosu Domozoro, C.; Marfo-Ahenkora, E.; Emadodin, I.; Malisch, C.S. Nitrogen Availability Determines the Long-Term Impact of Land Use Change on Soil Carbon Stocks in Grasslands of Southern Ghana. *SOIL* **2020**, *6*, 523–539. <https://doi.org/10.5194/soil-6-523-2020>.
25. Shatokhin, A.A.; Chamurliev, O.G.; Zelenev, A.V.; Chamurliev, G.O.; Vorontsova, E.S. Field Crop Rotations in Organic Agriculture of the Volgograd Region. *BIO Web Conf.* **2020**, *27*, 00152. <https://doi.org/10.1051/bioconf/20202700152>.
26. Bankó, L.; Tóth, G.; Marton, C.L.; Hoffmann, S. Hot-Water Extractable C and N as Indicators for 4p1000 Goals in a Temperate-Climate Long-Term Field Experiment: A Case Study from Hungary. *Ecol. Indic.* **2021**, *126*, 107364. <https://doi.org/10.1016/j.ecolind.2021.107364>.
27. Saha, M.; Das, M.; Sarkar, A. Distinct Nature of Soil Organic Carbon Pools and Indices under Nineteen Years of Rice Based Crop Diversification Switched over from Uncultivated Land in Eastern Plateau Region of India. *Soil Tillage Res.* **2021**, *207*, 104856. <https://doi.org/10.1016/j.still.2020.104856>.
28. Müller, T.S.; Dechow, R.; Flessa, H. Inventory and Assessment of PH in Cropland and Grassland Soils in Germany. *J. Plant Nutr. Soil Sci.* **2022**, *185*, 145–158. <https://doi.org/10.1002/jpln.202100063>.
29. Waldrip, H.; Schwartz, R.C.; He, Z.; Todd, R.W.; Baumhardt, R.L.; Zhang, M.; Parker, D.; Brauer, D.; Min, B.R. Soil Water Extractable Organic Matter under Long-Term Dryland Cropping Systems on the Texas High Plains. *Soil Sci. Soc. Am. J.* **2022**, *86*, 1249–1263.
30. Buraka, T.; Elias, E.; Lelago, A. Effects of Land-Use-Cover-Changes on Selected Soil Physicochemical Properties along Slope Position, Coka Watershed, Southern Ethiopia. *Heliyon* **2023**, *9*, e16142. <https://doi.org/10.1016/j.heliyon.2023.e16142>.
31. Bargaz, A.; Elhaissofi, W.; Khourchi, S.; Benmrid, B.; Borden, K.A.; Rchiad, Z. Benefits of Phosphate Solubilizing Bacteria on Belowground Crop Performance for Improved Crop Acquisition of Phosphorus. *Microbiol. Res.* **2021**, *252*, 126842.
32. Reis, V.M.; Teixeira, K.R. dos S. Nitrogen Fixing Bacteria in the Family Acetobacteraceae and Their Role in Agriculture. *J. Basic Microbiol.* **2015**, *55*, 931–949.
33. Villordon, A.Q.; Ginzberg, I.; Firon, N. Root Architecture and Root and Tuber Crop Productivity. *Trends Plant Sci.* **2014**, *19*, 419–425.
34. DuPont, S.T.; Beniston, J.; Glover, J.D.; Hodson, A.; Culman, S.W.; Lal, R.; Ferris, H. Root Traits and Soil Properties in Harvested Perennial Grassland, Annual Wheat, and Never-Tilled Annual Wheat. *Plant Soil* **2014**, *381*, 405–420.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.