

Housing Conditions and Its Impact on Health of Residents [†]

Mohd. Zuber ^{1,*}, Charu Khosla ² and Nargis Begum Javed ³

¹ Department of Allied Health Sciences, Chitkara School of Health Sciences, Chitkara University, Punjab, India

² Department of Applied Sciences, Chitkara University Institute of Engineering and Technology, Chitkara University, Punjab, India; charu.khosla@chitkara.edu.in

³ Department of Public Health, College of Health Sciences, Saudi Electronic University, Dammam, Saudi Arabia; n.javed@seu.edu.sa

* Correspondence: mohdzuber.phd2020@chitkara.edu.in; Tel.: +91-9415964211

[†] Presented at the 4th International Electronic Conference on Applied Sciences, 27 October–10 November 2023, Available online: <https://asec2023.sciforum.net/>.

Abstract: Housing attributes to the physical structure that provides shelter, social services with hygienic neighborhood to fulfill the essential needs of the people. Housing factors have shown to have an effect on an individual's status of physical, mental, social and economic well-being. Indoor environmental factors such as crowding, environmental tobacco smoke, biofuel, dampness, house dust mites, temperature, age of building, pets, indoor plants affect the wellbeing and productivity of the occupants. A literature review was done on studies of housing conditions and health outcomes conducted in India and abroad from 1999 to 2020. The studies assessed the housing quality by self-reported questionnaire administered through postal, face-to-face or internet. Visual signs and non-volumetric method were used to assess indoor air quality and housing conditions, while health of residents were assessed by self-reported questionnaire, or SF-36 questionnaire. Studies conducted in United States of America, Europe, United Kingdom, Middle East, Africa and Australasia revealed that factors effecting health conditions were ventilation, dampness, presence of molds, overcrowding, house dust mite allergens, age and renovation of buildings and these factors showed association with respiratory illness, cold, cough, asthma, conjunctivitis, atopic dermatitis and ear infection. However, studies in India revealed that lack of proper ventilation, use of traditional fuels, crowding and poor hygienic conditions are the main factors associated with acute respiratory infection, asthma, tuberculosis, cardiovascular disease and lung cancer. Thus, the review highlights that there is a need to improve housing conditions in India to enable the people to lead a healthy and productive life.

Keywords: housing condition; health outcomes; overcrowding; ventilation; dampness

Citation: Zuber, M.; Khosla, C.; Javed, N.B. Housing Conditions and Its Impact on Health of Residents. *Eng. Proc.* **2023**, *52*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor: Simeone Chianese

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Housing attributes to the physical structure that provides shelter, social services with hygienic neighbourhood to fulfill the essential needs of the people [1]. The World Health Organization (WHO) states that “housing should provide protection against communicable diseases, protection against injury, poisoning and chronic diseases and reduce psychological and social stress to a minimum” [2].

Housing factors have shown to have an effect on an individual's status of physical, mental, social and economic well-being [3]. Housing is a place in which a group of individuals use up more than 80% (3/5) of their time for daily activities, which can be somewhere between 15.5 to 15.7 h per day [4,5]. Housing is seen as a “key determinant of health” as it is stated “good housing and good health go together” [6]. Breyse et al. (2004) reported housing, as a neglected site for public health action [7].

Indoor environments are the fundamental environmental factors that affect the health, wellbeing and productivity of the occupants [8]. The consequence on health progresses with an increase in the exposure time and level of air pollution [9,10]. Indoor air is composed of a heterogeneous combination of bio-aerosols (fungi, bacteria, and irritants) and the non-biological particles such as dust, smoke and particle emissions produced during cooking activities along with organic and inorganic gases [11]. A high concentration of airborne microorganisms when present in indoor environments result in development of various health problems [12]. Indoor air pollution is being linked in ongoing research literature to a greater probability leading to upper and lower respiratory tract infections, aggravation of the underlying inflammatory lung diseases, leading to chronic obstructive pulmonary disease (COPD), cardiac events, stroke, eye disorders, tuberculosis, cancer, and hospital admissions that are particular in females and children those are most prone to developing such infections because of longer stay indoors [13–18]. In developing countries, around one-third of hospital in patients and 20–30% of fatalities among children below the age of five years occur due to Acute Respiratory Infections (ARIs) [19,20].

As limited studies have been conducted in India in respect to housing conditions and health of the people, this review was done to find out the housing conditions of residents and their health status that will help us to identify risk factors that are associated with health conditions. This study shall provide an evidence so that policy makers can take it into consideration regarding framing of common housing standards for the population.

2. Methods

The literature on the topic was searched on websites base-search.net, academic.microsoft.com, dimensions.ai, PubMed, Google scholar using keywords “housing quality” “indoor air quality” “ventilation” “dampness” “health outcomes” “health status” “ill health”, “India” and Boolean operator “AND” was used to make the search more precise. The literature review was classified based on research papers of studies conducted in India and other countries.

3. Results and Discussion

3.1. Studies Conducted in India

- Pokhrel et al. [14] aimed to associate indoor cooking with cataract. This was an institutional case-control study that was designed to include women with cataract as cases and women attending refractive error clinic as controls. The study utilized a standardized questionnaire to collect the data. The lack of proper ventilation in kitchen was identified as an independent risk factor for developing cataract.
- Firdaus G and Ahmad A [21] aimed to analyze the indoor air quality (IAQ) and its determinants with its contribution to possible health outcomes at the household level. A questionnaire was used for data collection across homes from 35 wards in Delhi. The highest risk to the health of individuals was posed by the utilization of traditional fuels, absence of a proper separate cooking space, vulnerability to ETS, and insufficient ventilation. Residents’ most common medical problem was found to be acute respiratory infections, which were closely linked to the usage of traditional fuels. Asthma was strongly associated to the use of conventional energy sources, exposure to ETS and poor ventilation. Exposure to ETS was found to have a strong correlation with lung cancer and cardiovascular diseases.
- In another study, Firdaus G and Ahmad A [22] aimed to assess accommodation standards with the possible health outcomes across three selected districts of the National Capital Territory (NCT) of Delhi, India. Door-to-door survey was conducted to collect the data. In the analysis, household level data was controlled for individual socio-economic and demographic covariates. Overcrowded housing conditions were observed in the selected districts. Respiratory infections were the main health outcomes attached to overcrowding that included ARI, tuberculosis and asthma.

- Vaid U and Evans GW [23] conducted a study to assess whether improved housing standards improve occupants health and well-being. Trained raters assessed the housing quality at a walk-through amidst females residing in public dwellings and those currently living in slums on waiting lists to relocate to the public housings. This is the only study from India that has assessed the slum rehabilitation program. Limitation of the study was that all the health outcomes reported in this study were self-reported through a standard questionnaire, which so may result in over- or under-reporting of health conditions. A better physical and mental health was depicted by those females shifted from slum locality to public housing but they had lesser social connections compared to those females who were staying in slum dwellings.

3.2. Studies Conducted in Other Countries

- Koskinen O et al. [24]. aimed to assess the health of adults based on the presence of moisture or mould in their houses. This study was conducted in Finland. The strengths were the survey used a random sampling for studying the outcome. The researcher collected data about presence of moisture and mould by visualizing it at the identification reported by the occupant. Secondly the health data was collected by sending postal questionnaire which helped in controlling the interviewer bias. Limitation of the study was that it had not adjusted the findings for other confounding factors which can be associated with the outcomes observed. In this study it was concluded that the exposure of occupants to moisture was related to the developing of sinusitis, acute bronchitis, nocturnal cough and dyspnoea and sore throat. Whereas, mould exposure of the occupants was associated with the development of common cold, cough without phlegm, nocturnal cough, sore throat, rhinitis, tiredness, and attention problems.
- Oie L et al. [25] conducted a study to assess the effect of ventilation rate in the development of bronchial obstruction during the first two years of life. The study was conducted in Oslo, Norway. The strength of the study was the selection of cases and controls which were matched for date of birth and the ventilation rate was measured so the results are reliable. In the households included in this study, the ventilation rate was not directly linked with the risk of developing bronchial obstruction.
- Evans J et al. [26] conducted a study with the aim to examine the association between damp housing and adult health including other factors that may influence health and can be a confounder. The study was conducted on adults, aged between 18–64 years, from Oxfordshire, Buckinghamshire, Berkshire and Northamptonshire. The strength of the study was the confounders which may influence health were acknowledged and use of SF-36 questionnaire a standardized health questionnaire. However, it has limitations like it is a secondary analysis and housing dampness was assessed by self-reporting which may have led to over- or under- reporting of dampness. Those occupants who were unable to keep their homes warm “very frequently” and “most of the time” had a worse health status than those who were unable to keep their homes warm “just sometimes.”
- Baker M et al. [27] conducted a case-control study to assess the risk factors related to meningococcal disease in Auckland children. The inclusion criterion was children less than 8 years of age with or without meningococcal disease as case and control respectively. The strength of the study was that the confounders were adjusted and overcrowding was measured by the number of adolescent and adult (10 years or older) household members residing in a room. In this study, a multivariate model was used to control the age, ethnicity, season and socioeconomic factors. There was a strong association between the possibility of developing meningococcal disease and overcrowding. Measurement was done by the number of adolescents and adults (10 years or older) residing in one room of a household. When two more adolescents or adults were added to a six-room residence, the risk was doubled.

- Kilpelainen M et al. [28] conducted a study to assess the relationship between housing dampness and respiratory symptoms in adults. The study was carried out on university students in Finland. The strength of this study was a large sample size, and analysis was performed controlling for confounders. Limitation of the study was dampness assessment and health condition assessment was based on self-report survey questionnaire, so may result in over- or under- reporting of dampness and health conditions. A positive association was observed with home dampness and conditions like recent asthma, allergic rhinitis, atopic dermatitis and common cold occurring more than or equal to four times a year along with other respiratory illnesses. There was a strong association between exposure to visible mould, asthma and common cold.
- Zock JP et al. [29] in his study aimed to assess the distribution and determinants of house dust allergens in 10 European countries. The strength of the study was large sample size, multi-centric assessment of allergens conducted on mattress dust sample, the details of housing features were obtained by observation and interview as well as geographical distribution and determinants were assessed using a common protocol. There was a large difference in allergen levels between study centres was located at different geographic locations. The Der p 1 allergens were reduced by low winter temperatures in comparison to Der f 1 allergens.
- Al-Khatib IA and Tabakhna H [30] in their study aimed to assess the impact of housing condition in camp with refugees' health especially respiratory symptoms and disease. The strength of the study was randomised sampling from the housing units covering one-fifth of the camp's population. Limitation of the analysis does not control for the confounders. Respiratory illnesses such as common cold, cough, tonsillitis, and ear infection were significantly linked to overcrowding and large population density, as well as poor living conditions such as dampness and mould.
- Habib R et al. [31] in his study aimed to examine the association between housing quality and chronic illness among the household members. This was a population based cross-sectional study conducted in Lebanon. The strength of the study was that the researcher conducted face-to-face interview to obtain the details of health and for assessing housing conditions. Moreover, a multivariate regression model was applied to examine the alliance of poor health and housing quality alongside controlling for other covariates. Housing conditions were found to have a substantial positive linkage with chronic illness.
- Turunen M et al. [3] in their study aimed to create a data collection and response system to help in the assessment of the Finnish housing stock on quality, health and safety. The strength of the study was a large sample size and random selection of the sample population. Limitation of this study was that all the health outcomes were self-reported through a questionnaire by the respondents, which may have resulted in over- or under- reporting of health conditions. Most respondents had mechanical ventilation or mechanical exhaust in their homes. The primary heating system used in the households was direct heating Among other heating systems used were electric heating and oil heating. The most common symptoms reported by the respondents were upper respiratory tract infections and/or arthralgia followed by general symptoms including eye and skin symptoms and muscular pain. Few respondents reported of having regularly sleeping problems, daily lower respiratory tract symptoms and doctor-diagnosed asthma. The common allergies reported by the respondents were hay fever and pet allergy.
- Pudpong N et al. [32] conducted a study with the aim to find out the indoor concentrations of particulate matter with less than 10µm in diameter (PM10). The strength lies in quantification of the PM10 as it provide a result which which will be reliable. An important finding was reported in the study. In a non-smoking setting, the most significant source of indoor PM10 concentrations was cooking. A positive correlation

was observed between the PM10 concentrations and number of children who complained of cough in the high polluted areas of Bangkok city.

- Ana GR et al. [33] in their study aimed to determine the amount of airborne microorganisms in the interior dwelling environment that predispose children under the age of five years to acute respiratory illnesses. In this study, a prospective case-control methodology was used. The study's strength was that it used a non-volumetric method to determine the load of indoor airborne microorganisms. The total counts per cubic metre were compared using the parameters provided by the American Industrial Hygiene Association. In comparison to controls, for cases the mean indoor total bacterial count was markedly elevated than the acceptable limit of (≤ 500 cfu/m³) recommended by American Industrial Hygiene Association. It was advocated to educate the mothers of below five years of children regarding the importance of improved ventilation, following of proper sanitary practices so as to reduce the indoor microbial load.
- Heijnen M et al. [34] conducted a systematic review and meta-analysis to compare the effects of shared sanitation with individual household latrines (IHLs) on health outcomes. Only 22 titles and abstracts out of 25,477 were found to meet the inclusion requirements. The majority of studies established a correlation between shared sanitation and poor health outcomes. However, the quality of these studies varied, and the data was insufficient, so they should be treated with caution.
- Tusting S L et al. [35] conducted a study with the aim to characterise all published and unpublished data, as well as assess the strength and quality of these data, in order to conduct a systematic evaluation of the evidence regarding the influence of housing renovations on malaria. African, Asian, and South American studies were included. Despite a scarcity of high-quality evidence, the direction and consistency of effects suggest that housing is a substantial malaria risk factor. Future research should look into the protective effects of certain dwelling features as well as incremental housing improvements linked to socioeconomic advancement.
- Palacios J et al. [36] in their study examined the association between housing conditions and health outcomes. A good sample size of 25,000 German families was used in the study, which spanned 25 years. Individual, housing, and temporal fixed factors were adjusted for, and the study was resilient to changes in socioeconomic position, lifestyle choices, and neighbourhood conditions. Individuals exposed to poor housing conditions were reported to having worst mental and physical health. A significant reduction in doctor visits was observed that was related to home renovations. This corroborated the findings on home conditions and health outcomes.

The information obtained from this review provides an evidence that changes in housing conditions are triggered by a number of factors including environment and climate. Though the prevalence of respiratory symptoms and diseases are common in developing countries, there is scarcity of data on housing condition and its association with health from developing countries including India.

4. Conclusions

Most of the research work focusing on impact of housing condition on occupants' health has been carried out in developed countries and have reported the indoor environment of house as detrimental to the occupants' health especially respiratory related symptoms and diseases. The studies which have been conducted in India have used self-reported method to assess the house quality or trained raters assessed it and indoor air quality was addressed. Therefore, there is a need to address the housing quality using quantitative measures to correctly recognize the indoor environmental factors detrimental to the health of the occupants. Further, based on the results of the study awareness is to be carried out amongst the residents, government officials and construction agencies in the

country regarding housing conditions, maintenance of the buildings and the health of individuals.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Yu, C.W.F.; Kim, J.T. Holistic healthy building for human habitat. *Indoor Built Environ.* **2011**, *20*, 3–4.
2. World Health Organization. *Health Principles of Housing*; World Health Organization: Geneva, Switzerland, 1989. Available online: <http://www.who.int/iris/handle/10665/39847> (accessed on).
3. Turunen, M.; Panala, A.; Villmen, J.; Nevalainen, A.; Haverinen-Shaughnessy, I. Evaluating housing quality, health and safety using an internet based data collection response: A cross-sectional study. *Environ. Health* **2010**, *9*, 69.
4. Bonnefoy, X.; Braubach, M.; Krapavickaite, D.; Ormandy, D.; Zurlyte, I. Housing conditions and self-reported health status: a study in panel block buildings in three cities of Eastern Europe. *J. Hous. Built Environ.* **2003**, *18*, 329–352.
5. Brasche, S.; Bischof, W. Daily time spent indoors in German homes—Baseline data for the assessment of indoor exposure of German occupants. *Int. J. Hyg. Environ. Health* **2005**, *208*, 247–253.
6. Breysee, S.J.; Alexander, A.; Easterlow, D. Rehousing as a health intervention: miracle or mirage? *Health Place* **1997**, *11*, 271–286.
7. Breyse, P.; Farr, N.; Galke, W.; Lanphear, B.; Morley, R.; Bergofsky, L. The relationship between housing and health children at risk. *Environ. Health Perspect.* **2004**, *112*, 1583–1588.
8. Gocgeldi, E.; Berdan, M.E.; Ucar, M.; Turker, T.; Istanbuluoglu, H.; Gulec, M.; Hasde, M. Analysis of children's rooms in terms of microbiological air quality. *J. Exp. Integr. Med.* **2011**, *1*, 51–58.
9. Hoskins, J.A. Health effects due to indoor air pollution. *Indoor Built Environ.* **2007**, *12*, 427–433.
10. Li, Y.; Leung, G.M.; Tang, J.W.; Yang, X.; Chao, C.Y.H.; Lin, J.Z.; Lu, J.W.; Nielsen, P.V.; Niu, J.; Qian, H.; et al. Role of ventilation in airborne transmission of infectious agents in the built environment—a multidisciplinary systematic review. *Indoor Air* **2007**, *17*, 2–18.
11. Hargreaves, M.; Parappukkaran, S.; Morawska, L.; Hitchins, J.; Congrong, H.; Gilbert, D. A pilot investigation into associations between indoor airborne fungal and non-biological particle concentrations in residential houses in Brisbane. *Sci. Total Environ.* **2003**, *312*, 89–101.
12. Stetzenbach, L.D.; Buttner, M.P.; Cruz, P. Detection and enumeration of airborne contaminants. *Curr. Biotechnol.* **2004**, *15*, 170–174.
13. Lin, H.H.; Murray, M.; Cohen, T.; Colijn, C.; Ezzati, M. Effects of smoking and solid-fuel use on COPD, lung cancer, and tuberculosis in China: a time-based, multiple risk factor, modelling study. *Lancet* **2008**, *372*, 1473–1483.
14. Pokhrel, A.K.; Smith, K.R.; Khalakdina, A.; Deuja, A.; Bates, M.N.. Case—control study of indoor cooking smoke exposure and cataract in Nepal and India. *Int. J. Epidemiol.* **2005**, *34*, 702–708.
15. Saha, A.; Rao, N.M.; Kulkarni, P.K.; Majumdar, P.K.; Saiyed, H.N. Pulmonary function and fuel use: a population survey. *Respir. Res.* **2005**, *6*, 127.
16. Smith, K.R.; Mehta, S.; et al. Indoor smoke from household use of solid fuels. In *Comparative Quantification of Health Risks: The Global Burden of Disease Due to Selected Risk Factors*; Ezzati, M., Lopez, AD., Rodgers, A., Murray, C.J.L., Eds.; World Health Organization: Geneva, Switzerland, 2004; Volume 2, pp. 1435–1493.
17. Kumar, P.; Kumar, M.; Bedi, O.; Gupta, M.; Kumar, S.; Jaiswal, G.; Rahi, V.; Yedke, N.G.; Bijalwan, A.; Sharma, S.; et al. Role of vitamins and minerals as immunity boosters in COVID-19. *Inflammopharmacology* **2021**, *29*, 1001–1016.
18. Mahajan, P.; Kaushal, J. Epidemic Trend of COVID-19 Transmission in India During Lockdown-1 Phase. *J. Community Health* **2020**, *45*, 1291–1300.
19. Demers, A.M.; Morency, P.; Mberyo-Yaah, F.; Jaffer, S.; Blais, C.; Somse, P. Risk factor for mortality among children hospitalized because of acute respiratory infections in Bangui, Central African Republic. *J. Pediatr. Infect Dis.* **2000**, *19*, 424–432.
20. Peppin, J.; Demers, A.M.; Mberyo-yaah, F.; Jaffer, S.; Blais, C.; Somse, P. Acute lower respiratory tract infection among children hospitalized in Bangui, Central African Republic: Towards a new case management algorithm. *Trans. R. Soc. Trop. Med. Hyg.* **2001**, *95*, 410–417.
21. Firdaus, G.; Ahmad, A. Indoor air pollution and self-reported diseases—A case study of NCT of Delhi. *Indoor Air.* **2011**, *21*, 410–416.
22. Firdaus, G.; Ahmad, A. Relationship between Housing and Health: A Cross-Sectional Study of an Urban Centre of India. *Indoor Built Environ.* **2012**, *22*, 498–507.
23. Vaid, U.; Evans, G.W. Housing Quality and Health: An Evaluation of Slum Rehabilitation in India. *Environ. Behav.* **2016**, *49*, 771–790.
24. Koskinen, O.M.; Husman, T.M.; Meklin, T.M.; Nevalainen, A.I. The relationship between moisture or mould observations in houses and the state of health of their occupants. *Eur. Respir. J.* **1999**, *14*, 1363–1367.
25. Oie, L.; Nafstad, P.; Botten, G.; Magnus, P.; Jaakkola, J.K. Ventilation in homes and bronchial obstruction in young children. *Epidemiology* **1999**, *10*, 294–299.
26. Evans, J.; Hyndman, S.; Stewart-Brown, S.; Smith, D.; Petersen, S. An epidemiological study of the relative importance of damp housing in relation to adult health. *J. Epidemiol. Community Health* **2000**, *54*, 677–686.

27. Baker, M.; McNicholas, A.; Garrett, N.; Jones, N.; Stewart, J.; Koberstein, V.; Lennon, D. Household crowding a major risk factor for meningococcal disease in Auckland children. *Pediatr. Infect Dis. J.* **2000**, *19*, 983–990.
28. Kilpelainen, M.; Terho, E.O.; Helenius, H.; Koskenvuo, M. Home dampness, current allergic diseases, and respiratory infections among young adults. *Thorax* **2001**, *56*, 462–467. <https://doi.org/10.1136/thorax.56.6.462>.
29. Zock, J.P.; Heinrich, J.; Jarvis, D.; Verlato, G.; Norback, D.; Plana, E.; Sunyer, J.; Chinn, S.; Olivieri, M.; Soon, A.; et al. Distribution and determinants of house dust mite allergens in Europe: the European Community Respiratory Health Survey II. *J. Allergy Clin Immunol.* **2006**, *118*, 682–690.
30. Al-Khatib, I.A.; Tabakhna, H. Housing conditions and health in Jalazone Refugee Camp in Palestine. *East. Mediterr. Health J.* **2006**, *12*, 144–152.
31. Habib, R.R.; Mahfoud, Z.; Fawaz, M.; Basma, S.H.; Yeretizian, J.S. Housing quality and ill health in a disadvantaged urban community. *Public Health* **2009**, *123*, 174–181.
32. Pudpong, N.; Krassi, R.; Nipapun, K. Indoor concentrations of PM10 and factors influencing its concentrations in day care centres in Bangkok, Thailand. *Asia J. Public Health J.* **2011**, *2*.
33. Ana, G.R.; Fakunle, G.A.; Ogunjobi, A.A. Indoor airborne microbial burden and risk of acute respiratory infections among children under five in Ibadan, Nigeria. *Indoor Built Environ.* **2013**, *0*, 1–7.
34. Heijnen, M.; Cumming, O.; Peletz, R.; Chan, G.K.-S.; Brown, J.; Baker, K.; Clasen, T. Shared Sanitation versus Individual Household Latrines: A Systematic Review of Health Outcomes. *PLoS ONE.* **2014**, *9*, e93300.
35. Tusting, S.L.; Ippolito, M.M.; Willey, B.A.; Kleinschmidt, I.; Dorsey, G.; Gosling, R.D.; Lindsay, S.W. The evidence for improving housing to reduce malaria: a systematic review and meta-analysis. *Malar. J.* **2015**, *14*, 209.
36. Palacios, J.; Eichholtz, P.; Kok, N.; Aydin, E. The impact of housing conditions on health outcomes. *Real Estate Econ.* **2020**, *49*, 1172–1200. <https://doi.org/10.1111/1540-6229.12317>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.