



The 9th International Electronic Conference on Medicinal Chemistry (ECMC 2023)

01–30 November 2023 | Online

Aptamer-based molecular probes: a multidisciplinary pipeline

Chaired by **Dr. Alfredo Berzal-Herranz**
and **Prof. Dr. Maria Emília Sousa**



pharmaceuticals



Victoria Calzada^{1,*}, Mauro de Castro¹, Santiago Pintos¹ Pablo Dans² and Hugo Cerecetto¹


¹ Aptamer Group, Centro de Investigaciones Nucleares. Facultad de Ciencias, Universidad de la República, Montevideo-Uruguay.;

² Computational Biophysics Group, Department of Biological Sciences, CENUR Litoral Norte (UdelaR), Montevideo-Uruguay.

* Corresponding author: vcalzada@fcien.edu.uy



Aptamer-based molecular probes: a multidisciplinary pipeline





Abstract:

Aptamers are small oligonucleotides that specifically recognize a molecular target with high affinity and specificity. Based on their versatility and physicochemical properties, several techniques were developed to exploit the advantages of aptamers as probes for molecular imaging. In our previous work, a DNA aptamer which exhibits specific binding for PTK7 receptor, was used as a molecular imaging probe. This aptamer was modified at 5'-position with several chelator agents and radiolabeled with technetium-99m or gallium-67. An Alexa647 derivative was also used as near infrared dye. All probes were physicochemical and biologically evaluated in BALB/c mice bearing lymphoma-tumor. In order to increase the specificity, the RP-HPLC purification procedure of probes was very strict. The chemical modifications were confirmed by mass spectrometry and RP-HPLC-gamma detection. Confocal microscopy and flow cytometry were performed in several tumor cells with the fluorescent probe. *In vivo* images were obtained in mice using X-ray, gamma and optical imaging strategies. After 24 h of injection, the probes were retained in the tumor and ratios tumor/non-target organ with values over 20 were obtained. In addition, *in vivo* specificity was evaluated in competition assays. The retention over time of the probe in the tumor and its quick excretion support the utility of labeled aptamers as molecular probes for diverse image strategies.

Keywords: Aptamer, Molecular Imaging, probes, drug-delivery



The 9th International Electronic Conference on Medicinal Chemistry

01–30 November 2023 | Online



Calzada, V.; Zhang, X.; Fernandez, M.; Diaz-Miqueli, A.; Iznaga-Escobar, N.; Deutscher, S.; Balter, H.; Quin, T.; Cabral, P. A potencial theranostic agent for EGF-R expression tumors: (177)Lu-DOTA-nimotuzumab. Curr. Radiopharm. 2012 Oct;5(4):318-24.



The 9th International Electronic Conference on Medicinal Chemistry

01–30 November 2023 | Online



Molecular Imaging

SPECT, γ emitters, ^{99m}Tc
PET, β^+ emitters, ^{18}F
FMT, fluorophores, $> 640\text{ nm}$

Reproduced from Ref. **Chem. Commun.**, 2013, **49**,
3812–3812 . with permission from the Royal Society
of Chemistry.


<http://pubs.rsc.org/en/Content/ArticleLanding/2013/CC/C3CC90110F#!divAbstract>

ChemComm

Chemical Communications

www.rsc.org/chemcomm

Volume 49 | Number 37 | 10 May 2013 | Pages 3811–3910



ISSN 1359-7345

RSC Publishing

COMMUNICATION

Rakesh N. Veedu, Kristofer J. Thurecht *et al.*

Aptamer-targeted hyperbranched polymers: towards greater specificity for tumours *in vivo*




Applications

- Tumour stratification,
- Metastasis detection,
- Guided surgery,
- Quantification of an injury,
- THERANOSTIC



APTAMERS



Aptamers are small oligonucleotides that specifically recognize a molecular target with high affinity and specificity.



Probe Design


Ref.: Shanguan, D.; Tang, Z.; Mallikaratchy, P.; Xiao, Z.; Tan, W. Optimization and modifications of aptamers selected from live cancer cell lines. *Chembiochem* 2007 Apr 16;8(6):603-6.

Name: Sgc8-c (DNA)

Size = 41 nb,

MW = 12634 Da

Target: PTK7 receptor



Sicco E, Báez J, Margenat J, García F, Ibarra M, Cabral P, Moreno M, Cerecetto H, Calzada V. Derivatizations of Sgc8-c aptamer to prepare metallic radiopharmaceuticals as imaging diagnostic agents: Syntheses, isolations, and physicochemical characterizations. *Chem Biol Drug Des*; 2018 Mar;91(3):747-755. doi: 10.1111/cbdd.13135. Epub 2017 Nov 24. PMID: 29080264.





The 9th International Electronic Conference on Medicinal Chemistry

01-30 November 2023 | Online





After 24 h of injection, *in vivo* and *ex vivo* images showed high tumor uptake.



Sicco E, Baez J, Ibarra M, Fernández M, Cabral P, Moreno M, Cerecetto H, Calzada V. Sgc8-c Aptamer as a Potential Theranostic Agent for Hemato-Oncological Malignancies. *Cancer Biother Radiopharm.* 2020 May;35(4):262-270. doi: 10.1089/cbr.2019.3402. PMID: 32407201.




Melanoma Sgc8-c-NOTA-⁶⁷Ga 24h



Sicco, Estefanía et al. "Metastatic and non-metastatic melanoma imaging using Sgc8-c aptamer PTK7-recognizer." *Scientific reports* vol. 11,1 19942. 7 Oct. 2021, doi:10.1038/s41598-021-98828-6




Biodistribution



Calzada V, Moreno M, Newton J, González J, Fernández M, Gambini JP, Ibarra M, Chabalgoity A, Deutscher S, Quinn T, Cabral P, Cerecetto H. Development of new PTK7-targeting aptamer-fluorescent and -radiolabelled probes for evaluation as molecular imaging agents: Lymphoma and melanoma in vivo proof of concept. *Bioorg Med Chem.* 2017 Feb 1;25(3):1163-1171. doi: 10.1016/j.bmc.2016.12.026. Epub 2016 Dec 24. PMID: 28089349.




Tumor/no tumor RATIOS




Sicco, Estefanía et al. "Metastatic and non-metastatic melanoma imaging using Sgc8-c aptamer PTK7-recognizer." *Scientific reports* vol. 11,1 19942. 7 Oct. 2021, doi:10.1038/s41598-021-98828-6



Chemistry modifications effect




Arévalo AP, Castelli R, Ibarra M, Crispo M, Calzada V. In Vivo Evaluation of Sgc8-c Aptamer as a Molecular Imaging Probe for Colon Cancer in a Mouse Xenograft Model. *Int J Mol Sci.* 2022 Feb 23;23(5):2466. doi: 10.3390/ijms23052466. PMID: 35269608; PMCID: PMC8910571.






Polymeric micelles are good candidates to improve the pharmacokinetics and the tumor uptake of aptamer-based probes




Castelli R, Ibarra M, Faccio R, Miraballes I, Fernández M, Moglioni A, Cabral P, Cerecetto H, Glisoni RJ, Calzada V. T908 Polymeric Micelles Improved the Uptake of Sgc8-c Aptamer Probe in Tumor-Bearing Mice: A Co-Association Study between the Probe and Preformed Nanostructures. *Pharmaceuticals* (Basel). 2021 Dec 23;15(1):15. doi: 10.3390/ph15010015. PMID: 35056072; PMCID: PMC8780797.





Aptamers as drug-delivery system



Sicco E.; Almeida L.; Moreno M.; **Calzada V***; Cerecetto H. Chemical conjugations of Sgc8-c with the lymphoma drug dasatinib to generate selective biotherapeutics. *Aptamers*. 11:19942 2021.

Sicco E, Cerecetto H, Calzada V, Moreno M. Targeted-Lymphoma Drug Delivery System Based on the Sgc8-c Aptamer. *Cancers* (Basel). 2023 Feb 1;15(3):922. doi: 10.3390/cancers15030922. PMID: 36765879; PMCID: PMC9913644.





Aptamer_uy group

- SELEX → more information

- Sequencing and data analyses

- Screening

- Characterization and optimization





The 9th International Electronic Conference on Medicinal Chemistry

01-30 November 2023 | Online



Versatility



The 9th International Electronic Conference on Medicinal Chemistry

01-30 November 2023 | Online



THANKS

FOR WOMEN IN SCIENCE



AGENCIA NACIONAL
DE INVESTIGACIÓN
E INNOVACIÓN



United Nations
Educational, Scientific and
Cultural Organization

L'ORÉAL



UNIVERSIDAD
DE LA REPÚBLICA
URUGUAY



PEDECIBA
MEC-UDELAR



Ministerio
de Educación
y Cultura