

Proceeding Paper

Enhancing Corn Yield Prediction in Iowa: A Concatenate-Based 2D-CNN-BILSTM Model with Integration of Sentinel-1/2 and SoilGRIDs Data [†]

Mahdiyeh Fathi ¹, Reza Shah-Hosseini ^{1,*} and Armin Moghimi ²

¹ School of Surveying and Geospatial Engineering, College of Engineering, University of Tehran, Tehran, Iran; mahdiyeh.fathi@ut.ac.ir

² Ludwig-Franzius-Institute for Hydraulic, Estuarine and Coastal Engineering, Leibniz University Hannover, Hanover, Germany; moghimi@lufi.uni-hannover.de

* Correspondence: rshahosseini@ut.ac.ir

[†] Presented at the 5th International Electronic Conference on Remote Sensing, 7–21 November 2023; Available online: <https://ecrs2023.sciforum.net/>.

Abstract: Ensuring food security in precision agriculture demands early prediction of corn yield in the USA at international, regional, and local levels. Accurate corn yield estimation can play a crucial role in averting famine by offering insights into food availability during the growing season. To address this, we propose a Concatenate-based 2D-CNN-BILSTM model that integrates Sentinel-1, Sentinel-2, and Soil-GRIDs (global gridded soil information) data for corn yield estimation in Iowa State from 2018 to 2021. This approach utilizes Sentinel-2 features, including spectral bands (Blue, Green, Red, Red Edge 1/2/3, NIR, n-NIR, and SWIR 1/2), and vegetation indices (NDVI, LSWI, DVI, RVI, WDRVI, SAVI, VARIGREEN, and GNDVI), alongside Sentinel 1 features (VV, VH, difference VV, and VH, and RVI), and Soil data (Silt, Clay, Sand, CEC, and pH) as initial inputs. To extract high-level features from this data each month, a dedicated 2D-CNN was designed. This 2D-CNN concatenates high-level features from the previous month with low-level features of the subsequent month, serving as input features for the model. Additionally, to incorporate single-time soil data features, another 2D-CNN was implemented. Finally, high-level features from soil, Sentinel-1, and Sentinel-2 data were concatenated and fed into a BILSTM layer for accurate corn yield prediction. Comparative analysis against random forest (RF), Concatenate-based 2D-CNN, and 2D-CNN models, using metrics like RMSE, MAE, MAPE, and the Index of Agreement, revealed the superiority of our model. It achieved an Index of Agreement of 84.67% with an RMSE of 0.698 t/ha. The Concatenate-based 2D-CNN model also performed well with an RMSE of 0.799 t/ha and an Index of Agreement of 72.71%. The 2D-CNN model followed closely with an RMSE of 0.834 t/ha and an Index of Agreement of 69.90%. In contrast, the RF model lagged with an RMSE of 1.073 t/ha and an Index of Agreement of 69.60%. Integration of Sentinel 1–2 and Soil-GRIDs data with the Concatenate-based 2D-CNN-BILSTM model significantly improved accuracy. Combining soil data with Sentinel 1–2 features reduced RMSE by 16 kg and increased the Index of Agreement by 2.59%. This study highlighted the potential of advanced Machine Learning (ML)/Deep Learning (DL) models in achieving precise and reliable predictions, which could support sustainable agricultural practices and food security initiatives.

Citation: Fathi, M.; Shah-Hosseini, R.; Moghimi, A. Enhancing Corn Yield Prediction in Iowa: A Concatenate-Based 2D-CNN-BILSTM Model with Integration of Sentinel-1/2 and SoilGRIDs Data. *2023*, *27*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: corn yield prediction; 2D-CNN-BILSTM; Sentinel-1/2; soil-Grids data

1. Introduction

Corn is a highly significant crop in the United States (U.S.) due to its abundance of protein, oil, and its high water consumption [1,2]. As the largest corn producer globally,

U.S. recorded a corn production of 15.1 billion bushels in 2021 (<https://www.nass.usda.gov/Newsroom/2022/01-12-2022.php>).

With the rapid increase in population, the use of Remote Sensing (RS) technology in agriculture has become of paramount economic importance. By using multisensor RS images, and soil and weather data, researchers can accurately predict crop yield. Optical RS data, obtained from satellites such as Sentinel-2 and Landsat-8, provide high/moderate-resolution imagery in visible and near-infrared bands. This enables precise assessment of vegetation health through indices like NDVI, while also offering detailed information on land cover for accurate crop type identification, including corn [3]. Additionally, optical RS data allows for monitoring phenological changes, facilitating the tracking of crop growth stages and overall health [4]. On the other hand, SAR (Synthetic Aperture Radar) images facilitate the structural analysis of vegetation by using different polarizations and microwave frequencies. Additionally SAR signals penetrates cloud cover, providing valuable insights into the physical structure of crops [5]. The applications of RS in agriculture are diverse and encompass product and irrigation management, predicting crop performance, disease and fertilizer management, as well as crop classification, among others [3]. However, the effectiveness of these applications hinges on various factors including temperature, rainfall, growth indicators, soil type, genotype structure, management practices, and nutrient elements [6]. Additionally, radiometric distortions have the potential to adversely affect the spectral bands of optical RS images [7]. To mitigate these challenges and enhance the accuracy of yield predictions, a multi-faceted approach is recommended. This involves integrating RS data with advanced machine learning (ML) models and employing data fusion techniques.

For example, Ma et al. suggested the Bayesian neural network to estimate corn yield using MODIS images, GLDAS dataset, PRISM dataset, and SSURGO at the county level in the United States between 2005 and 2019 [8]. Desloires et al. introduced a stack of machine learning techniques, namely RF, SVR, XG-Boost, and MLP, to predict corn yield based on Sentinel-2 images captured at field-scale in Iowa and Nebraska from 2017 to 2021 [9]. Khaki et al. proposed the Deep-Corn network for enhancing crop yield at the field-scale by counting corn kernels, which used a shortened VGG-16 for feature extraction at different scales [10]. Shah-Hosseini et al. developed the Stacked LASSO method for predicting corn yield in Illinois, Indiana, and Iowa between 2000 and 2018 using observed corn yield, management data, plant population, planting date, and environmental features (weather and soil) [11]. Shah-Hosseini et al. also proposed a new CNN-DNN method for estimating corn using historical management, environmental, and yield data in the United States from 1980 to 2019 [12]. San et al. suggested the CNN-RNN method for predicting corn yield using MODIS images, weather data, and soil features to extract multi-level spatiotemporal features at the county level from 2013 to 2016 [13]. Dhaliwal et al. proposed the Random Forest model for predicting corn yield using crop management data, weather data, and field-level data in the United States between 1992 and 2018 [14]. Shah-Hosseini et al. suggested combining the APSIM model with machine learning methods using plant population, planting date, and weather data for estimating corn yield in the United States between 1984 and 2018 [15].

Recent studies have demonstrated satisfactory outcomes in estimating crop yield [11,16–19]. However, they have less regarded the combination of radar and optical images, along with soil data for corn yield prediction. Moreover, most of the studies used CNN-LSTM for feature extraction, and they have not fully explored the benefits of combining high-level features from the previous month with low-level features from the subsequent month to improve corn yield [20]. Also, they mostly have employed Long Short-Term Memory (LSTM) networks for yield prediction, overlooking the potential of Bidirectional-LSTM (Bi-LSTM) networks which can integrate both past and future information to enhance corn yield forecasting [21].

In response to these limitations, we introduce a novel Concatenate-Based 2D-CNN-BiLSTM model for corn yield estimation at the county level in Iowa. Leveraging Sentinel-

1 and Sentinel-2 images along with Soil-GRIDs, which provide global gridded soil information, our model aims to enhance performance during the growing season. This model offers an innovative approach to feature integration, effectively capturing short-term fluctuations and long-term trends in corn growth patterns. Additionally, the incorporation of Soil-GRIDs data provides crucial insights into soil characteristics, augmenting the model's capacity to account for diverse soil conditions.

2. Materials and Methods

2.1. Study Area

The study area was located in IOWA State of U.S., which (see Figure 1). The research was done during the years 2018 to 2021 on corn. Corn is planted in Iowa when the soil is warm enough for the seeds to grow but not too early to avoid frost damage. The timing varies depending on the location, with southern counties planting as early as April and northern counties waiting several weeks later. Farmers in Iowa typically begin harvesting corn in mid-September, with the majority of the harvest taking place in October. However, in cooler years, the harvest may not take place until November (<https://www.iowacorn.org/education/faqs>).




Figure 1. Study Area.

2.2. Dataset

A variety of dataset was used to forecast corn yield from 2018 to 2021, including Sentinel-1 SAR (COPERNICUS/S1_GRD), Sentinel-2 SR (S2_SR_HARMONIZED), Soil-Grids (<https://www.isric.org/explore/soilgrids>), USDA Yield (<https://quickstats.nass.usda.gov/>), Crop Land Data Layer (CDL), and County Boundaries data [22,23]. The Sentinel 1 and 2 was downloaded from the Google Earth Engine (GEE) cloud computing platform [24]. Table 1 displays the statistical characteristics of yield observations for both the training and test datasets.

Table 1. Sample plot yield statistics for year in study area.

Type	Year	Number of Samples	Min (Ton Ha ⁻¹)	Max (Ton Ha ⁻¹)	Mean (Ton Ha ⁻¹)	Std (Ton Ha ⁻¹)
train	2018	93	9.38	14.18	12.13	1.25
train	2019	88	9.50	14.73	12.19	1.06
train	2020	95	5.54	12.99	10.92	1.17
test	2021	84	9.57	14.49	12.57	1.05

2.3. Methodology

The aim of the proposed method is to improve corn yields prediction accuracy at the county level in IOWA State in-during growing season prior to harvest on the month of August. As displayed in Figure 2, our proposed method includes two main steps: 1 – extracting features derived from Sentinel-1, Sentinel-2, and Soil-Grids in the GEE system, and 2—using the proposed Concatenate-Based 2D-CNN-BiLSTM model to predict corn yield. The details of each steps briefly have been explained in the following subsections.




Figure 2. Flowchart of the proposed method.

2.3.1. Feature Selection

As discussed before, various features derived from Sentinel1 \2 and Soil-Grids were used to predict corn yield in our study area. Informative spectral bands of Sentinel-2 images (i.e., Blue, Green, Red, Red Edge 1/2/3, NIR, n-NIR, and SWIR 1/2 bands), along with its Vis (i.e., NDVI, LSWI, DVI, RVI, WDRVI, SAVI, VARI-GREEN, and GNDVI) were used as input optical features [25–32]. Additionally, VV, VH, difference of VV, and VH, and Radar Vegetation Index (RVI) [33] were extracted from Sentinel-1 SAR images [34]. Soil data including silt, clay, sand, cec, and pH was also collected at various depths ranging from 0 cm to 200 cm to construct our feature set (<https://www.isric.org/explore/soilgrids>).

2.3.2. Corn Yields Prediction Using the Concatenate-Based 2D-CNN-BiLSTM Model

As prediction of corn yield is so challenging, improvement of the advance and novel deep learning model for accurately predicting corn yield is important. In this way we proposed the Concatenate-Based 2D-CNN-BiLSTM model which have two main parts (see Figure 3) including feature extraction using a 2D-CNN network, and corn yield prediction using a Bi-LSTM network. The 2D-CNN network extracts high-level spatial features from input data and concatenates them with low-level features from subsequent months [35].

Additionally, a separate 2D-CNN network was created to incorporate single-time soil data features. Finally, the high-level features from soil, Sentinel-1, and Sentinel-2 data were concatenated and fed into a Bi-LSTM layer to accurately predict corn yield. The Bi-LSTM layer is able to overcome significant time lags between inputs across any time period, and enhance its ability to represent temporal patterns at different frequencies using backward and forward information [21]. This makes it particularly advantageous for analyzing crop growth cycles of varying durations. Monthly-Block consists of Conv2D-1 > Linear activation function > Concatenate layer > Conv2D > Linear activation function. Monthly-composites ($X_{Monthly}$) pass through the Monthly-Block, and monthly-features ($F_{Monthly}$) are extracted. In addition, Soil-Block consists of Conv2D-3 > Linear activation function. Soil features pass through the Soil-Block, and soil-features (S) are then extracted. $F_{Monthly}$ and S are then concatenated together, and fed into Bi-LSTM layer with ReLU activation function to predict corn yield. Finally, the output of the Bi-LSTM layer passes through a dense layer with linear activation function to obtain yield values.

Overall, our Concatenate-Based 2D-CNN-BiLSTM model is a promising approach for accurately predicting corn yield by incorporating various data sources and effectively capturing temporal patterns.




Figure 3. Flowchart of the proposed Concatenate-Based 2D-CNN-BiLSTM model for corn yield prediction.

3. Results and Discussion

For this study, a total of 250, 27, and 83 samples were selected for training, validation, and testing the Concatenate-Based 2D-CNN-BiLSTM model, respectively. The Conv2D-1, and Conv2D-2 layers were set to have 16, and 22 filters, respectively, with a kernel size of 1×1 . The Bi-LSTM layer had 16 filters. The model was trained using the *Adam* optimizer for 30 epochs with a batch size of 10. The best weight was obtained based on the minimum Validation-Loss. The performance of the proposed model was compared with Concatenate-Based 2D-CNN, 2D-CNN, and RF in two scenarios: (1) using Sentinel-1 and -2 data, and (2) using both Sentinel-1 and -2 data along with Soil-Grids. Table 2 displays the performance of the proposed models and the compared models, measured in terms of RMSE, MAPE, MAE, RRMSE, and Index of Agreement (D).

Table 2. Comparison of Performance of Proposed Concatenate-Based 2D-CNN-BiLSTM Model versus Other Considered Methods for Corn Yield Prediction.

Model	Sentinel 1&2					Sentinel 1&2 and Soil-Grids				
	RMSE	RRMSE	MAE	MAPE	D	RMSE	RRMSE	MAE	MAPE	D
Proposed Model	0.714	5.68	0.561	4.55	82.08	0.687	5.46	0.566	4.47	84.67
Concatenate-Based 2D-CNN	0.849	6.75	0.686	5.60	67.58	0.799	6.35	0.620	5.02	72.71
2D-CNN	0.848	6.74	0.694	5.64	64.80	0.834	6.63	0.677	5.51	69.90
RF	1.089	8.66	0.935	7.95	69.04	1.073	8.54	0.918	7.78	69.60

Table 2 reveals that the Concatenate-Based 2D-CNN-BiLSTM model outperforms the Concatenate-Based 2D-CNN, 2D-CNN, and RF methods significantly. The best performance of the Concatenate-Based 2D-CNN-BiLSTM model is achieved when combining Sentinel-1&2 and Soil-Grids, with an RMSE of 0.698 (t/ha), MAPE of 4.47%, MAE of 0.566 (t/ha), RRMSE of 5.46%, and D of 84.67%. Our proposed model improves D by 14.77% compared to the 2D-CNN.

Figure 4 depicts the scatter plots of predicted yield versus observed yield between our proposed method and compared methods in 2021. The scatter plots demonstrate that the fit line is close to the diagonal line in the Concatenate-Based 2D-CNN model and far in the RF model.




Figure 4. The scatter plots of predicted yield versus observed yield between our proposed method and compared methods in 2021.

Additionally, Figure 5 illustrates that the proposed model outperforms the compared models, resulting in a reduction in Error maps and generate the brighter error map. This confirms the efficacy of utilizing soil-Grids data for yield estimation.




Figure 5. Error maps generated by the proposed and compared models in 2021.

A visual representation of the distribution of corn yield value was presented in Figure 6, which compared the USDA yield with the predicted yield obtained from our proposed method. The results displayed in Figure 6 indicate a significant level of agreement

between the observed and predicted corn yield, thereby reinforcing the reliability and accuracy of our proposed method's predictions.




Figure 6. Map of USDA corn yield, and predicted corn yield in 2021.

4. Conclusions

Forecasting corn yield is a crucial aspect of agriculture management in IOWA. Recent studies have demonstrated that remote sensing, soil data, and Deep Learning methods are effective in estimating corn yield. In order to accurately predict corn yield, it is important to consider both temporal and spatial features. To achieve this, we propose a novel Concatenate-Based 2D-CNN-BiLSTM model that extracts both spatial and temporal features. The CNNs extract spatial features, while Bi-LSTM extracts temporal features. The inputs for our model include remote sensing data (Sentinel-1&2) and Soil-Grids data. We conducted experiments with the proposed model on IOWA Corn from 2018 to 2021 at the county-level. Our results demonstrate the effectiveness and advantages of our approach over other methods. By considering both spatial and temporal features, our model is able to accurately forecast corn yield, which can aid in making informed decisions for agriculture management in IOWA.

Author Contributions: Conceptualization, M.F., R.S.-H. and A.M.; Methodology, M.F., R.S.-H. and A.M.; Project administration, R.S.-H.; Resources, M.F., R.S.-H. and A.M.; Validation, M.F., R.S.-H. and A.M.; Supervision, R.S.-H.; Writing—original draft, M.F., R.S.-H. and A.M.; Writing—review and editing, M.F., R.S.-H. and A.M. All authors have read and agreed to the published version of the manuscript.

Funding: This article was not received funding.

Institutional Review Board Statement:

Informed Consent Statement:

Data Availability Statement:

Acknowledgments: The authors would like to thank the U.S. Department of Agriculture (USDA) for providing the yield values. We extend our heartfelt thanks to Google for providing us with Google Earth Engine and Google Colab. These platforms have been instrumental in developing our deep learning models, enabling efficient geospatial analysis and model training.

Conflicts of Interest:

References

1. Fathi, M.; Shah-Hosseini, R.; Moghimi, A. Comparison of Some Deep Neural Networks for Corn and Soybean Mapping in Iowa State using Landsat imagery. *Earth Obs. Geomat. Eng.* **2022**, *6*.
2. Fathi, M.; Shah-Hosseini, R. Automatic Corn and Soybean Mapping Based on Deep Learning Methods (case Study: Hamilton, Hardin, Boone, Story, Dallas, Polk, and Jasper Counties in Iowa State). *ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci.* **2023**, *10*, 187–193.

3. Ali, A.M.; Abouelghar, M.; Belal, A.; Saleh, N.; Yones, M.; Selim, A.I.; Amin, M.E.; Elwesemy, A.; Kucher, D.E.; Maginan, S. Crop yield prediction using multi sensors remote sensing. *Egypt. J. Remote Sens. Space Sci.* **2022**, *25*, 711–716.
4. Amani, M.; Kakooei, M.; Moghimi, A.; Ghorbanian, A.; Ranjgar, B.; Mahdavi, S.; Davidson, A.; Fiset, T.; Rollin, P.; Brisco, B. Application of google earth engine cloud computing platform, sentinel imagery, and neural networks for crop mapping in Canada. *Remote Sens.* **2020**, *12*, 3561.
5. Moghimi, A.; Mohammadzadeh, A.; Khazai, S. Integrating thresholding with level set method for unsupervised change detection in multitemporal SAR images. *Can. J. Remote Sens.* **2017**, *43*, 412–431.
6. Dharani, M.; Thamilselvan, R.; Natesan, P.; Kalaivaani, P.; Santhoshkumar, S. Review on crop prediction using deep learning techniques. In Proceedings of the Journal of Physics: Conference Series, International E-Conference on Data Analytics, Intelligent Systems and Information Security & ICDIIS 2020, Pollachi, India, 11–12 December 2020; p. 012026.
7. Moghimi, A.; Celik, T.; Mohammadzadeh, A. Tensor-based keypoint detection and switching regression model for relative radiometric normalization of bitemporal multispectral images. *Int. J. Remote Sens.* **2022**, *43*, 3927–3956.
8. Ma, Y.; Zhang, Z.; Kang, Y.; Özdoğan, M. Corn yield prediction and uncertainty analysis based on remotely sensed variables using a Bayesian neural network approach. *Remote Sens. Environ.* **2021**, *259*, 112408.
9. Desloires, J.; Ienco, D.; Botrel, A. Out-of-year corn yield prediction at field-scale using Sentinel-2 satellite imagery and machine learning methods. *Comput. Electron. Agric.* **2023**, *209*, 107807.
10. Khaki, S.; Pham, H.; Han, Y.; Kuhl, A.; Kent, W.; Wang, L. Deepcorn: A semi-supervised deep learning method for high-throughput image-based corn kernel counting and yield estimation. *Knowl.-Based Syst.* **2021**, *218*, 106874.
11. Shahhosseini, M.; Hu, G.; Archontoulis, S.V. Forecasting corn yield with machine learning ensembles. *Front. Plant Sci.* **2020**, *11*, 1120.
12. Shahhosseini, M.; Hu, G.; Khaki, S.; Archontoulis, S.V. Corn yield prediction with ensemble CNN-DNN. *Front. Plant Sci.* **2021**, *12*, 709008.
13. Sun, J.; Lai, Z.; Di, L.; Sun, Z.; Tao, J.; Shen, Y. Multilevel deep learning network for county-level corn yield estimation in the us corn belt. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **2020**, *13*, 5048–5060.
14. Dhaliwal, D.S.; Williams, M.M. Sweet corn yield prediction using machine learning models and field-level data. *Precis. Agric.* **2023**, 1–14.
15. Shahhosseini, M.; Hu, G.; Huber, I.; Archontoulis, S.V. Coupling machine learning and crop modeling improves crop yield prediction in the US Corn Belt. *Sci. Rep.* **2021**, *11*, 1606.
16. Rao, C.; Liu, Y. Three-dimensional convolutional neural network (3D-CNN) for heterogeneous material homogenization. *Comput. Mater. Sci.* **2020**, *184*, 109850.
17. Mohite, J.; Sawant, S.; Pandit, A.; Agrawal, R.; Pappula, S. Soybean Crop Yield Prediction by Integration of Remote Sensing and Weather Observations. *Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.* **2023**, *48*, 197–202.
18. Schwalbert, R.A.; Amado, T.; Corassa, G.; Pott, L.P.; Prasad, P.V.; Ciampitti, I.A. Satellite-based soybean yield forecast: Integrating machine learning and weather data for improving crop yield prediction in southern Brazil. *Agric. For. Meteorol.* **2020**, *284*, 107886.
19. You, J.; Li, X.; Low, M.; Lobell, D.; Ermon, S. Deep gaussian process for crop yield prediction based on remote sensing data. In Proceedings of the AAAI Conference on Artificial Intelligence, San Francisco, CA, USA, 4–9 February 2017.
20. Sun, J.; Di, L.; Sun, Z.; Shen, Y.; Lai, Z. County-level soybean yield prediction using deep CNN-LSTM model. *Sensors* **2019**, *19*, 4363.
21. Siami-Namini, S.; Tavakoli, N.; Namin, A.S. The performance of LSTM and BiLSTM in forecasting time series. In Proceedings of the 2019 IEEE International Conference on Big Data (Big Data), Los Angeles, CA, USA, 9–12 December 2019; pp. 3285–3292.
22. Malenovsky, Z.; Rott, H.; Cihlar, J.; Schaepman, M.E.; García-Santos, G.; Fernandes, R.; Berger, M. Sentinels for science: Potential of Sentinel-1, -2, and -3 missions for scientific observations of ocean, cryosphere, and land. *Remote Sens. Environ.* **2012**, *120*, 91–101.
23. Boryan, C.; Yang, Z.; Mueller, R.; Craig, M. Monitoring US agriculture: The US department of agriculture, national agricultural statistics service, cropland data layer program. *Geocarto Int.* **2011**, *26*, 341–358.
24. Amani, M.; Ghorbanian, A.; Ahmadi, S.A.; Kakooei, M.; Moghimi, A.; Mirmazloumi, S.M.; Moghaddam, S.H.A.; Mahdavi, S.; Ghahremanloo, M.; Parsian, S. Google earth engine cloud computing platform for remote sensing big data applications: A comprehensive review. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **2020**, *13*, 5326–5350.
25. Sonobe, R.; Yamaya, Y.; Tani, H.; Wang, X.; Kobayashi, N.; Mochizuki, K.-i. Crop classification from Sentinel-2-derived vegetation indices using ensemble learning. *J. Appl. Remote Sens.* **2018**, *12*, 026019.
26. Gitelson, A.A.; Kaufman, Y.J.; Merzlyak, M.N. Use of a green channel in remote sensing of global vegetation from EOS-MODIS. *Remote Sens. Environ.* **1996**, *58*, 289–298.
27. Wang, C.; Wu, Y.; Hu, Q.; Hu, J.; Chen, Y.; Lin, S.; Xie, Q. Comparison of Vegetation Phenology Derived from Solar-Induced Chlorophyll Fluorescence and Enhanced Vegetation Index, and Their Relationship with Climatic Limitations. *Remote Sens.* **2022**, *14*, 3018.
28. Richardson, A.J.; Everitt, J.H. Using spectral vegetation indices to estimate rangeland productivity. *Geocarto Int.* **1992**, *7*, 63–69.
29. Christian, J.I.; Basara, J.B.; Lowman, L.E.; Xiao, X.; Mesheske, D.; Zhou, Y. Flash drought identification from satellite-based land surface water index. *Remote Sens. Appl. Soc. Environ.* **2022**, *26*, 100770.

30. Broge, N.H.; Leblanc, E. Comparing prediction power and stability of broadband and hyperspectral vegetation indices for estimation of green leaf area index and canopy chlorophyll density. *Remote Sens. Environ.* **2001**, *76*, 156–172.
31. Eng, L.S.; Ismail, R.; Hashim, W.; Baharum, A. The use of VARI, GLI, and VIgreen formulas in detecting vegetation in aerial images. *Int. J. Technol.* **2019**, *10*, 1385–1394.
32. Huete, A.R. A soil-adjusted vegetation index (SAVI). *Remote Sens. Environ.* **1988**, *25*, 295–309.
33. Arii, M.; van Zyl, J.J.; Kim, Y. A general characterization for polarimetric scattering from vegetation canopies. *IEEE Trans. Geosci. Remote Sens.* **2010**, *48*, 3349–3357.
34. Holtgrave, A.-K.; Röder, N.; Ackermann, A.; Erasmi, S.; Kleinschmit, B. Comparing Sentinel-1 and-2 data and indices for agricultural land use monitoring. *Remote Sens.* **2020**, *12*, 2919.
35. Yu, C.; Han, R.; Song, M.; Liu, C.; Chang, C.-I. A simplified 2D-3D CNN architecture for hyperspectral image classification based on spatial–spectral fusion. *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.* **2020**, *13*, 2485–2501.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.