

Mineralogical Characterization of PM₁₀ over the Central Himalayan Region †

Sakshi Gupta^{1,2*}, Priyanka Srivastava³, Manish Naja³, Nikki Choudhary^{1,2} and Sudhir Kumar Sharma^{1,2}

¹ CSIR-National Physical Laboratory, Dr. K S Krishnan Road, New Delhi-110012, India; sakshigupta21096@gmail.com (S.G.); nikichoudhary1607@gmail.com (N.C.); sudhircsir@gmail.com (S.K.S)

² Academy of Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India

³ Aryabhata Research Institute of Observational Sciences (ARIES), Nainital 263002, Uttarakhand, India; srivastava.priyanka@nies.go.jp (P.S); manish@aries.res.in (M.N)

* Correspondence: sakshigupta21096@gmail.com

† Presented at the 6th International Electronic Conference on Atmospheric Sciences, 15-30 October 2023. Available online: <https://ecas2023.sciforum.net>.

Abstract: The air quality of Himalayan region of India is deteriorating due to the increasing load of particulate matter that is emitted from various local and regional sources as well as transit of dust-related pollutants from the Indo-Gangetic Plain (IGP) and surrounding areas. In this study, the mineralogical characteristics of coarse mode particulate matter (PM₁₀) has been analysed using X-Ray Diffraction (XRD) technique from January–December, 2019 over Nainital (29.39°N, 79.45°E; altitude: 1958 m above mean sea level), a central Himalayan region of India. XRD analysis of PM₁₀ samples showed the presence of clay minerals, crystalline silicate minerals, carbonate minerals, and asbestiform minerals. It is shown that the quartz minerals with significant levels of crystallinity were present in all the samples. Other minerals that are contributing to the soil dust are also observed in the analysis (CaFe₂O₄, CaCO₃, CaMg(CO₃)₂, calcium ammonium silicate hydrate (C-A-S-H), gypsum, kaolinite, illite, augite, and montmorillonite). The minerals, ammonium sulphate, hematite, and magnetite were also found in the samples and are suggested to be from the biogenic and anthropogenic activities including biomass burning, fuel combustion, vehicle exhaust, construction activities etc. This study indicated that the majority of the minerals in PM₁₀ that were present in this Himalayan region are from soil/crustal dust.

Keywords: PM₁₀; Himalayas; XRD; Dust; Clay Minerals; Sources

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Last-name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The Indian Himalayan Region (IHR) is renowned for its pristine nature, ecological fragility, abundant biodiversity, and remarkable vulnerability, making it one of the most crucial regions on the Earth [1-3]. The central Himalayan region of India is widely recognized for its unique geological and environmental attributes [4-6]. Urbanization in and around the Himalayas has led to increased energy consumption, resulting in disturbances to the temperature of the Himalayas and the degradation of its air quality [7,8]. One of the main reasons of deteriorating air quality is increase in the load of particulate matter (PM) emissions over the region. These PM emissions are known to generally arise from local and regional sources as well as migration of dust-related pollutants from Indo-Gangetic Plain (IGP) and surrounding areas [4, 5, 9]. Hence, to achieve better understanding of the composition, sources, and potential impact of PM in the central Himalayan region, mineralogical characterization of PM₁₀ has been conducted. This study aims to analyze and identify the mineral components present in the PM, particularly focusing on the coarse mode particulate matter (PM₁₀) fraction.

2. Materials and Methods

2.1. Study area and Sampling

The study was conducted in ARIES, Nainital, a central Himalayan region of India (29.39°N, 79.45°E, 1959 m amsl) (Figure 1). 24h PM₁₀ sampling was performed from January- December, 2019 using a high-volume sampler with average flow rate of 1.2 m³ min⁻¹ with a flow accuracy of ±2 % of full scale. Detailed information regarding sampling procedures and instrumentation can be found in earlier publications [5].




Figure 1. Study Area: ARIES, Nainital (Source: Google Earth).

2.2. X-Ray Diffraction Analysis

X-ray Diffraction (XRD) analysis was performed on the collected PM₁₀ samples to determine the mineralogical characteristics. The XRD technique made this possible by identifying and quantifying the mineral contents in the sample based on their unique diffraction patterns. The XRD measurements were conducted using a Rigaku Ultima IV instrument. The samples were scanned during the XRD investigation at a Bragg angle (2 θ) varying from 10 to 60 degrees to gather the X-ray diffraction data. 3° per minute was the scanning speed employed. As the X-ray source, a copper (Cu) K α -line with a wavelength of 1.54 Å was used. Mineral identification was carried out by comparing the peak positions (2 θ) from the XRD data with the reported literature [10-12] and RRUFF database of a reference standard.

3. Results and Discussion

3.1. Mineralogical composition

The XRD analysis of PM₁₀ revealed the presence of various minerals. Illite, kaolinite, montmorillonite, quartz, dolomite, calcite, magnetite, hematite, gypsum, halite, mascagnite, augite, albite, wollastonite, and calcium aluminium silicate hydrate (C-A-S-H) are the common minerals that were detected in all the samples (Table 1). Figure 2 shows the XRD pattern of the mineral content that were present in PM₁₀ samples. It is important to note that these minerals may be found in various environmental contexts, and their presence can have different implications depending on the concentrations. Additionally, some of the minerals mentioned can have multiple sources and pathways of formation. Understanding their sources and impacts is crucial for environmental and health considerations.




Figure 2. XRD pattern for the PM₁₀ sample. Quartz (Q), dolomite (D), albite (Al), augite (Au), illite (I), kaolinite (K), montmorillonite (M), Hematite (H), magnetite (Mg), gypsum (G), wollastonite (W), mascalinite (AS), and halite (H).

3.2. Soil-dust composition

We used quartz fibre filter for sampling of PM₁₀ samples that has silicate composition and also it is the mineral that is present in soil/ crustal dust samples. Quartz minerals with significant crystallinity were consistently detected in all the samples, indicating their ubiquitous presence. Previous studies illustrated that the presence of quartz mineral in the samples were due to the geographical characteristics of the respective locations i.e., soil, land cover, land use pattern, road length, etc. [10-15]. Dolomite, albite and augite (small amount) are other minerals detected in PM₁₀ samples which have natural/geological origin i.e., originated from soil and road dust, weathering of rocks, etc. [10, 11, 16].

Table 1. Minerals, corresponding XRD peak positions (2θ).

Mineral	Chemical Composition	2θ
Quartz	SiO ₂	20.64, 26.50, 40.46
Dolomite	(Ca, Mg (CO ₃) ₂)	30.70, 37.56, 50.52, 50.94
Augite	(Ca, Mg, Fe) ₂ Si ₂ O ₆	19.80, 30.70, 34.64, 40.46, 40.84, 41.92, 49.56
Albite	Na(AlSi ₃ O ₈)	13.74, 14.74, 15.96, 23.28, 24.60, 27.96, 53.26
Calcite	CaCO ₃	23.28, 29.36, 36.06, 47.88, 48.30
Kaolinite	Al ₂ O ₃ .2SiO ₂ .2H ₂ O	19.80, 20.64, 21.34, 23.28, 24.60, 36.06, 37.56, 38.96, 40.46, 40.85, 47.88, 52.72, 55.22, 56.74
Illite	(K, H ₃ O ⁺) (Al, Mg, Fe) ₂ (Si, Al) ₄ O ₁₀ [(OH) ₂ , (H ₂ O)]	17.36, 26.82
Montmorillonit e	(Na,Ca) _{0.3} (Al,Mg) ₂ SiO ₄ (OH) ₂ .n H ₂ O	19.80, 21.34, 34.64
Magnetite	Fe ₃ O ₄	56.74
Hematite	Fe ₂ O ₃	40.84, 49.56
Gypsum	CaSO ₄ .2H ₂ O	20.64, 23.28
Halite	NaCl	31.88, 56.74

Mascagnite	$(\text{NH}_4)_2\text{SO}_4$	20.64, 29.36, 34.64, 38.96
Wollastonite	CaSiO_3	23.28, 51.96, 53.26
C-A-S-H	$\text{Ca}_{12}\text{Al}_2\text{Si}_{18}\text{O}_{51}(\text{OH})_2 \cdot 18\text{H}_2\text{O}$	29.36

3.3. Anthropogenic contribution 104

The minerals calcite, kaolinite, illite, montmorillonite, magnetite, hematite, gypsum, and wollastonite were found in PM₁₀ samples that are originated from both geogenic as well as biogenic and anthropogenic sources. The minerals illite [10, 11, 17], kaolinite [10-12, 16], and montmorillonite [11, 14] were primarily originated from geological processes and various industrial processes, agricultural activities, combustion activities (fossil fuel, biomass burning, etc.) and were also responsible for the generation of these minerals in the air [5, 10, 14, 16]. Hematite [11, 14, 15] and magnetite [12] both are Fe containing minerals that primarily originated from soil erosion and weathering of rocks, dust storm and also from various human activities such as iron and steel production, combustion (vehicular emission, power plants), iron and steel wear from brake pads and tires [13-16]. Candeias et al., (2020) illustrated that the mineral containing Fe, Cu, Zn, S, Al, Ti and Sb composition were originated through anthropogenic activities like brake pads, brake disc abrasion, and road wears etc. [15]. Gypsum [12], C-A-S-H [10, 12] and wollastonite [12] are the minerals in which Ca majorly contributed in their composition. These minerals are originated through mining, construction and demolition activities [10, 12]. Mascagnite i.e., ammonium sulphate is majorly contributing through secondary reaction that is occurring in the atmosphere [12]. The main sources of mascagnite in PM₁₀ are fossil fuel burning (coal and oil), industrial emissions, biomass burning, waste incineration, etc., [6]. Halite is the salt mineral that is used as de-icing agent during winters [12]. Various studies illustrated the transport of dust aerosols from Indo-Gangetic Plain (IGP), the Thar desert, Bay of Bengal (BoB), and other regional countries towards the IHR [3, 5].

4. Conclusion 126

Through the mineralogical characterization of PM₁₀ collected over central Himalayan region of India from January- December, 2019, valuable insights have been gained regarding types of mineral present in the airborne particles. Present study sheds light on the sources and origin of these minerals, such as natural dust, anthropogenic emissions, or a combination thereof. Certain minerals such as quartz, dolomite, albite, and augite have been found in the study have natural origin. Additionally, minerals like illite, kaolinite, montmorillonite, hematite, magnetite, gypsum, calcium aluminium silicate, ammonium sulphate, halite, etc., associated with the biogenic and anthropogenic activities like combustion, mining, construction, demolition, etc., have also been detected. Hence, by understanding the mineralogical characteristics of PM, policymakers, scientists, and environmentalists can gain crucial insights into the sources, composition, and potential health impacts of PM.

Author Contributions: Chemical analysis, writing—original draft preparation, writing—review and editing, S.G.; sample collection, chemical analysis, data curation, writing—review and editing, P.S.; N.C.; S.G.; conceptualization, investigation, supervision, funding acquisition, S.K.S.; M.N.; All authors have read and agreed to the published version of the manuscript. 139-142

Funding: The authors also acknowledge the Department of Science and Technology (DST), New Delhi, India for financial support for this study (DST/CCP/Aerosol/88/2017). 143-144

Institutional Review Board Statement: Not applicable. 145

Informed Consent Statement: Not applicable. 146

Data Availability Statement: The datasets are available with corresponding author and will be provided on reasonable request. 147-148

Acknowledgments: The authors are thankful to the Director, CSIR-NPL, New Delhi and Head, Environmental Sciences & Biomedical Metrology Division, CSIR-NPL, New Delhi for their encouragement for this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Yuan, Q., Wan, X., Cong, Z., Li, M., Liu, L., Shu, S., Liu, R., Xu, L., Zhang, J., Ding, X., Li, W. In situ observations of light-absorbing carbonaceous aerosols at Himalaya: Analysis of the south Asian sources and trans-Himalayan valleys transport pathways. *J. Geophys. Res.* 2020, 125, e2020JD032615. <https://doi.org/10.1029/2020JD032615>
2. Yang, J., Ji, Z., Kang, S., Tripathee, L. Contribution of south Asian biomass burning to black carbon over the Tibetan Plateau and its climatic impact. *Environ. Pollut.* 2021, 270, 116195. <https://doi.org/10.1016/j.envpol.2020.116195>
3. Rai, A., Mukherjee, S., Choudhary, N., Ghosh, A., Chatterjee, A., Mandal, T., Sharma, S.K., Kotnala, R. Seasonal transport pathway and sources of carbonaceous aerosols at an urban site of eastern Himalaya. *Aerosol Sci. Eng.* 2021, 5, 318–343. <https://doi.org/10.1007/s41810-021-00106-5>
4. Sharma, S.K., Choudhary, N., Srivastava, P., Naja, M., Vijayan, N., Kotnala, G., Mandal, T.K. Variation of carbonaceous species and trace elements in PM₁₀ at a mountain site in the central Himalayan region of India. *J. Atmos. Chem.* 2020, 77, 1–14. <https://doi.org/10.1007/s10874-020-09402-9>
5. Choudhary N., Srivastava P., Dutta M., Mukherjee S., Rai A., Kuniyal J.C., Lata R., Chatterjee A., Naja, M., Vijayan, N., Mandal, T.K., Sharma, S.K. Seasonal Characteristics, Sources and Pollution Pathways of PM₁₀ at High Altitudes Himalayas of India. *Aerosol Air Qual. Res.* 2022, 22 (7). <https://doi.org/10.4209/aaqr.220092>.
6. Choudhary, N., Rai, A., Kuniyal, J.C., Srivastava, P., Lata, R., Dutta, M., Ghosh, A., Dey, S., Sarkar, S., Gupta, S., Chaudhary, S., Thakur, I., Bawari, A., Naja, M., Vijayan, N., Chatterjee, A., Mandal, T.K., Sharma, S.K., Kotnala, R.K. Chemical Characterization and Source Apportionment of PM₁₀ Using Receptor Models over the Himalayan Region of India. *Atmosphere.* 2023, 14, 880. <https://doi.org/10.3390/atmos14050880>.
7. Ram, K., Sarin, M.M. Spatio-temporal variability in atmospheric abundances of EC, OC and WSOC over Northern India. *J. Aerosol Sci.* 2010, 41, 88–98. <https://doi.org/10.1016/j.jaerosci.2009.11.004>
8. Sharma, S.K., Mukherjee, S., Choudhary, N., Rai, A., Ghosh, A., Chatterjee, A., Vijayan, N., Mandal, T. Seasonal variation and sources of carbonaceous species and elements in PM_{2.5} and PM₁₀ over the eastern Himalaya. *Environ. Sci. Pollut. Res.* 2021, 28, 51642–51656. <https://doi.org/10.1007/s11356-021-14361-z>
9. Jain, S., Sharma, S.K., Srivastava, M.K., Chatterjee, A., Singh, R.K., Saxena, M., Mandal, T.K. Source Apportionment of PM₁₀ Over Three Tropical Urban Atmospheres at Indo-Gangetic Plain of India: An Approach Using Different Receptor Models. *Arch. Environ. Contam. Toxicol.* 2019, 76(1), 114–128. <https://doi.org/10.1007/s00244-018-0572-4>.
10. Bora, J., Deka, P., Bhuyan, P. *et al.* Morphology and mineralogy of ambient particulate matter over mid-Brahmaputra Valley: application of SEM–EDX, XRD, and FTIR techniques. *SN Appl. Sci.* 3, 137. <https://doi.org/10.1007/s42452-020-04117-8>
11. Neupane B.B., Sharma A., Giri B., Joshi M.K., 2020. Characterization of airborne dust samples collected from core areas of Kathmandu Valley. *Heliyon.* 2021, 6 (4). <https://doi.org/10.1016/j.heliyon.2020.e03791>.
12. Kimothi, S., Chilkoti, S., Rawat, V., Thapiyal, A., Gautam, A.S., Gautam, S. Micro- to macro-scaling analysis of PM_{2.5} in sensitive environment of Himalaya, India. *Geological Journal.* 2023, 1–19. <https://doi.org/10.1002/gj.4765>.
13. Kumar, S., Jain, M.K. Characterization and morphometric study of household settled dust: A case study in Dhanbad, the coal capital of India. *Applied Geochemistry.* 2022, 144, 105398. <https://doi.org/10.1016/j.apgeochem.2022.105398>.
14. Senthil Kumar, R., Rajkumar, P. Characterization of minerals in air dust particles in the state of Tamilnadu, India through FTIR, XRD and SEM analyses. *Infrared Physics & Technology.* 2014, 67, 30–41. <https://doi.org/10.1016/j.infrared.2014.06.002>.
15. Candeias, C.; Vicente, E.; Tomé, M.; Rocha, F.; Ávila, P.; Célia, A. Geochemical, Mineralogical and Morphological Characterisation of Road Dust and Associated Health Risks. *Int. J. Environ. Res. Public Health.* 2020, 17, 1563. <https://doi.org/10.3390/ijerph17051563>
16. Gunawardana C, Goonetilleke A, Egodawatta P, Dawes L, Kokot S. Source characterisation of road dust based on chemical and mineralogical composition. *Chemosphere.* 2012, 87(2):163–70. <https://doi.org/10.1016/j.chemosphere.2011.12.012>.
17. Nowak, S., Lafon, S., Caquiereau, S., Journet, E., Laurent, B. Quantitative study of the mineralogical composition of mineral dust aerosols by X-ray diffraction. *Talanta.* 2018, 15, 186: 133–139. <https://doi.org/10.1016/j.talanta.2018.03.059>.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.