

Proceeding Paper

# Uranyl Acetate, a Lewis Acid Catalyst for Acetoxylation of Monoterpenic and Steroidal Alcohols <sup>†</sup>

Juan Enrique Tacoronte Morales <sup>1,\*</sup>, Carla Bernal Villavicencio <sup>1</sup>, Xavier Leopoldo Gracia Cervantes <sup>1</sup>,  
Maria Elizabeth Canchingre <sup>1</sup> and Maria Teresa Cabrera Pedroso <sup>2</sup>

<sup>1</sup> Multidisciplinary Research Group, GIM-FACI, Faculty of Science & Technology, Chemical Engineering Coordination, Campus New Horizons, Technical University of Esmeraldas, Esmeraldas, Ecuador; email1@email.com (C.B.V.); email1@email.com (X.L.G.C.); email1@email.com (M.E.C.)

<sup>2</sup> Facultad de Ciencias de la Salud, Universidad de Las Americas, UDLA, UDLA-Park, Quito 170124, Ecuador; email1@email.com

\* Correspondence: jetacoronte@yahoo.com; Tel.: +593-991702915

<sup>†</sup> Presented at the 27th International Electronic Conference on Synthetic Organic Chemistry (ECSOC-27), 15–30 November 2023; Available online: <https://ecsoc-27.sciforum.net/>.

**Abstract:** The use of heterogeneous catalytic systems in the production of esters of secondary alcohols (monoterpenic and steroidal) allows obtaining satisfactory yields of these derivatives widely used in the industry. The use of the heterogeneous system uranyl acetate/chloroform-acetonitrile, under laboratory conditions, allows to reach yields higher than 90%, optimizing the purification steps and without the use of extreme conditions (inert atmosphere, high temperatures or corrosive agents). All compounds were characterized by the use of spectroscopic techniques.

**Keywords:** uranyl acetate; esterification; monoterpenic and steroidal alcohols; esters

## 1. Introduction

Esters have high potential in industry and pharmaceuticals because of their wide use in fragrances, flavors, surfactants, plasticizers, drugs and as solvents [1]. Generally, they are synthesized by catalytic acetylation of alcohols with, pyridine, 4-(dimethylamino) pyridine (DMAP) or 4-pyrrolidinopyridine (PPy) in presence of acyl chlorides or acid anhydrides [2,3]. Acetylation is one of the primary reactions in organic synthesis, since acetyl groups are useful for protecting various kinds of functional groups, such as alcohols, amines, phenols and thiols, among others [4–7]. It has been developed, at laboratory or industrial scale, under heterogeneous and homogeneous catalytic conditions as well as in the presence of acidic or basic catalysts, including biogenic catalytic extracts [8–12].

These methods suffer from inherent disadvantages such as high cost of the catalysts employed at laboratory and industrial scales, low stability of catalytic systems in the presence of moisture, poor regioselectivity in case of allylic alcohols, elimination byproducts in tertiary alcohols, racemization of optically active alcohols, epimerization of steroidal and tedious workup during and purification procedures. In this context, one of the most versatile ways for protecting hydroxyl groups in alcohols is the Lewis-mediated esterification using a variety of acylation reagents such as acid anhydrides and proper catalysts.

In [13], was reported that terminal olefins were converted selectively towards the Markovnikov's products, 1-methylalkylacetate without isomerization using vanadium oxide in presence of trifluoroacetic acid. This prompted us to develop a new Lewis acid catalyst with acetate ligands on uranium oxide and its acetoxy salts for acetylation reactions in a different solvent system.

We report herein a heterogeneous system, uranyl (VI) acetate /acetonitrile/ chloroform, for acetylation of alcohols (monoterpenic and steroidal) with acetic anhydride at

**Citation:** Morales, J.E.T.; Villavicencio, C.B.; Cervantes, X.L.G.; Canchingre, M.E.; Pedroso, M.T.C. Uranyl Acetate, a Lewis Acid Catalyst for Acetoxylation of Monoterpenic and Steroidal Alcohols. *Chem. Proc.* **2023**, *14*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s): Name

Published: 15 November 2023



**Copyright:** © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

laboratory scale. Incidentally, this is one of the few reports on the heterogenization of a homogeneous system, guided by the simple principle of insolubility, that evolves and functions as a genuine 'heterogeneous system' facilitating the recyclability and reusability of catalyst [14].

Uranyl acetate has been used in the photochemical oxidation of imidazolyl thione derivatives. It has also been reported for use in catalytic mixtures, alongside other uranyl salts, for highly efficient photocatalytic conversion processes from C-H to C-C bonds. Additionally, it has been utilized in photo assisted selective oxidation processes of alcohols and sulfides, as well as in the organic pollutant control [15–17].

## 2. Materials and Methods

### 2.1. General Procedures

All commercial reagents (oxides, acetoxy salts, acetic anhydride, acetic acid, and selected monoterpenic alcohols) and solvents ( $\text{CHCl}_3$ ,  $\text{CH}_3\text{CN}$ ) were purchased from Sigma-Aldrich/Merck Life Science (Darmstadt, Alemania), and they were used without any further purification or synthetic modification. All yields refer to the isolated products. Determination of the purity of substrates, related acetoxy derivatives and the monitoring of the acetoxylation reaction were accomplished by means of thin layer chromatography (TLC) on silica-gel plates 60 F-254 (Germany). All O-acetoxy derivatives were identified by their spectra and physical data. Melting points were determined by using the capillary tube method with an electrothermal 9100 apparatus.  $^1\text{H-NMR}$  spectra ( $\text{CDCl}_3$ ) were registered on Bruker AC-250 MHz spectrometer (Germany), at 25 °C, using tetramethylsilane as internal standard.

### 2.2. Preparation of the Catalyst

Uranyl (VI) acetate was prepared by refluxing uranium oxide  $\text{UO}_3$  (1.82 g) in excess of acetic anhydride (50 mL) for 2 h. The slight-yellow solid product was then filtered. Washed with  $\text{CHCl}_3$  (150 mL) and dried under vacuum at room temperature for 1 h (yield 3.5 g, 95%,  $\text{UO}_2(\text{CH}_3\text{COO})_2$ , powder insoluble in organic solvents, with slight odor of vinegar). The product obtained is kept in a closed flask at 18 °C in the absence of light until it is used in the acetylation process.

#### 2.2.1. Precautionary Measures

Uranyl acetate, a water-soluble uranium compound, is often used as a stain in electron microscopy. However, it requires the adoption of basic safety precautions, emphasizing the avoidance of any possibility of inhalation or ingestion of the material. Given the inherent toxicity of uranyl salts, when working with uranium oxide and uranyl acetate, the use of NK-45 masks, protective eyewear, and a lab coat is mandatory for amounts up to 10 g. Work periods should never exceed four hours, and gloves must be worn when handling or weighing uranyl acetate. To prevent bench surface contamination, use spill trays (metal or plastic) with disposable coverings such as bench coats, and clean the surface after use. Adopting appropriate control measures is essential to minimize risk, including:

- Reducing the amount of material handled as much as possible.
- Not exceeding a working temperature of 200 °C to avoid the thermal decomposition of uranyl acetate.
- Containing unsealed sources to prevent contamination.
- Maintaining a high level of cleanliness.
- Not disposing of uranyl acetate as ordinary waste

### 2.3. Acetylation Procedure


The alcohol (2.0 mmole), acetic anhydride (3 mmole) and the catalysts (0.2 mmole, 10 %w) were stirred in acetonitrile-chloroform (5 mL 3:2 *v/v*) at 70 °C for 3–4 h. The progress

of the reaction was controlled by TLC using ethyl-acetate-dichloromethane eluent mixture and vanilline-sulphuric acid (10%) as chromogenic agent. After completion of the reaction, the mixture was extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 30$  mL). Combined organic layers were washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . The solvent was removed and the crude products were column chromatographed on silica gel column to afford pure acetate esters which were subjected for NMR- $^1\text{H}$  spectroscopy. The catalyst was then filtered, removed, and kept for another cycle ( $5\times$ ) with previous activation at  $50^\circ\text{C}$  during 2 h.

### 3. Results and Discussion

The conversion of various alcohols, including, steroidal, and monoterpene derivatives, to their corresponding esters was easily achieved, almost quantitatively, by treating the alcohol with 1.5 equivalents of acetic anhydride at  $70^\circ\text{C}$  for 3–4 h in the presence of 10% mol of the catalyst in the described solvent mixture.

The process of acetoxylation takes place according to the proposed Scheme 1.









**Scheme 1.** Acetoxylation of alcohols with  $\text{Ac}_2\text{O}$  in the presence of Uranyl Acetate.

The results of the acetoxylation process, using the proposed catalytic system, are shown (*grosso modo*) in Table 1.

Steroidal alcohols such as progesterone, estrone and androstenol react smoothly with the catalytic system (uranyl acetate/acetonitrile-chloroform) within 3–4 h with very satisfactory yields.

**Table 1.** Acetoxylation of alcohols (steroidal and monoterpene) with  $\text{Ac}_2\text{O}/\text{UO}_2(\text{CH}_3\text{COO})_2$ .


Entry	Substrate	O-Acetoxy-Derivative <sup>a</sup>
1	 Pregnenolone	 Pregnenolone acetate (m.p.: $191.7^\circ\text{C}$ /96.8%)
2	 Estrone	 Estrone acetate (m.p.: $129.7^\circ\text{C}$ /97.3%)

3		
	Δ-4-androsten-17β-ol	Δ-4-androsten-17β-ol-acetate (m.p.: 142.6 °C/83.4%)
4	(-)-Menthol	Menthyl acetate (b.p.: 57.8 °C/82%)
5	(+/-)-endo-Norborneol	Norbornyl acetate (b.p.: 202 °C/85%)
6	α-Terpineol	(±)-α-Terpinyl acetate (b.p.: > 185 °C) <sup>b</sup>


<sup>a</sup> The reported yields of isolated products are an average from over five subsequent batches. <sup>b</sup> Was observed a sudden darkening of the reaction mixture.

In the spirostannic series, in the presence of this catalytic system, a similar behavior of the diosgenin and hecogenin substrates is observed, generating their corresponding acetates with excellent yields.

The Figure 1 depicts the results obtained for spirostannic alcohols. The reported yields of isolated products are an average from over five subsequent batches. The final product was re-chromatographed twice on silica-gel.



Diosgenin Acetate (m.p.: 195.80 °C/yield: 93.1%)



Hecogenin Acetate (m.p.: 243.60 °C/yield: 90.4%)

**Figure 1.** Spirostannic alcohol acetates obtained in the presence of  $\text{Ac}_2\text{O}/\text{UO}_2(\text{CH}_3\text{COO})_2/\text{Acetonitrile-Chloroform}/3\text{--}4\text{ h}/70\text{ }^\circ\text{C}$ .

An important features worthy to describe is that the selective acetylation of hydroxy carbonyl compounds (entries 2 and 3, Table 1) is achieved without the formation of  $\alpha,\beta$ -unsaturated carbonyl compounds due to the elimination of the resulting acetate.

The most significant achievement is the reusability of the salt of transition metal catalyst for several number of cycles (5×) with almost consistently activity. The synergistic effect of Lewis acids of insoluble solid salt, and solvent mixture, influences the acetoxylation reaction for higher activity as solid acid. The advantage of using this catalytic system, at laboratory scale, can be summarized as follows: Uranyl acetate can be easily prepared

directly from uranium oxide and acetic anhydride; the mixture of  $\text{UO}_2(\text{AcO})_2/\text{Acetonitrile-Chloroform}$  is stable and not revealed any toxic or hazardous action during the storage (1 year) and utilization at laboratory scale; the spirocetalic unit of spirostene derivatives (diosgenin and hecogenin, Figure 1) remains intact and was not observed any isomerization of the olefinic bound in steroidal alcohol; it is not necessary any inert atmosphere or dry conditions for performing the reaction; the catalyst is easy recovered by filtration and centrifugation; and could be re-used in 5 catalytic cycles after thermal activation; not observed any formation of by-products like ketene, olefine o acetoacetate derivatives. It should be noted that, under the reaction conditions, no decomposition of uranyl acetate is observed.

Some implications in kinetic resolution and mechanistic studies are underway, as well as its application to other hydroxy functionalized substrates (carbohydrates and tertiary alcohols). The mechanistic perception of the acetoxylation process in the presence of uranyl acetate as catalyst will be published soon.

In conclusion, the monoterpene and secondary steroidal alcohols are acetoxylation satisfactorily, with remarkable selectivity, and yields, using acetic anhydride in the presence of uranyl (VI) acetate as catalyst.

**Author Contributions:** J.E.T.M. Conceptualization and methodology; C.B.V. investigation and validation; X.L.G.C. formal analysis and software; M.E.C. formal analysis, investigation and project administration; M.T.C.P. writing—review and editing, resources and investigation. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no government, academic, institutional or external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** This study was sponsored, in part, by Ecuador's Prometheus Project "Development of catalytic processes under eco-sustainable conditions" 2015–2017.

**Conflicts of Interest:** The authors declare no conflict of interest. The Technical University of Esmeraldas, Ecuador, had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## References

1. Riemenscheider, W. *Ullmann's Encyclopedia of Industrial Chemistry*, 5th ed.; Verlag Chemie: Weinheim, Germany, 1997; Volume A9, p 565.
2. Hoffe, G.; Steglich, V.; Vorbruggen, H. 4-Dialkylaminopyridines as Highly Active Acylation Catalysts. *Angew. Chem. Int. Ed. Eng.* **1978**, *17*, 569–583. <https://doi.org/10.1002/anie.197805691>.
3. Scriven, E.F.V. 4-Dialkylaminopyridines: Super acylation and alkylation catalysts. *Chem. Soc. Rev.* **1983**, *12*, 129–161. <https://doi.org/10.1039/CS9831200129>.
4. Pasricha, S.; Rangarajan, T.M. Green Acetylation of Primary Aromatic Amines. *Resonance* **2023**, *28*, 325–331. <https://doi.org/10.1007/s12045-023-1551-2>.
5. Zhang, Z.; Zhao, Z.; Liu, M.; Liu, H.; Li, Q.; Xiang, J.; Wu, T.; Han, B. Acylation of phenols to phenolic esters with organic salts. *Green Chem.* **2022**, *24*, 9763–9771. <https://doi.org/10.1039/D2GC03807B>.
6. Rahmatpour, A.; Alinejad, S.; Donyapeyma, G. Noncross-linked polystyrene nanoencapsulation of ferric chloride: A novel and reusable heterogeneous macromolecular Lewis's acid catalyst toward selective acetylation of alcohols, phenols, amines, and thiols. *J. Organomet. Chem.* **2022**, *961*, 122264. <https://doi.org/10.1016/j.jorganchem.2022.122264>.
7. Sartori, G.; Ballini, R.; Bigi, F.; Bosica, G.; Maggi, R.; Righi, P. Protection (and deprotection) of functional groups in organic synthesis by heterogeneous catalysis. *Chem. Rev.* **2004**, *104*, 199–250. <https://doi.org/10.1021/cr0200769>.
8. Valentini, F.; Galloni, P.; Brancadoro, D.; Conte, V.; Sabuzi, F. A Stoichiometric Solvent-Free Protocol for Acetylation Reactions. *Front. Chem.* **2022**, *10*, 842190. <https://doi.org/10.3389/fchem.2022.842190>.
9. Chutia, R.; Chetia, B. Acetylation of alcohols, phenols and amines using waste plant extract. *SN Appl. Sci.* **2020**, *2*, 1564–1571. <https://doi.org/10.1007/s42452-020-03353-2>.
10. Mayr, S.; Zipse, H. Annelated Pyridine Bases for the Selective Acylation of 1,2-Diols. *Eur. J. Org. Chem.* **2022**, *29*, e202101521. <https://doi.org/10.1002/ejoc.202101521>.

11. Peng, P.; Linseis, M.; Winter, R.F.; Schmidt, R.R. Regioselective Acylation of Diols and Triols: The Cyanide Effect. *J. Am. Chem. Soc.* **2016**, *138*, 6002–6009. <https://doi.org/10.1021/jacs.6b02454>.
12. Estevão, M.S.; Afonso, C.A.M. Effect of a catalyst in the Acylation of Alcohols with Acetic Anhydride: Manipulation of a natural aroma. In *Comprehensive Organic Chemistry Experiments for the Laboratory Classroom*; Royal Society of Chemistry: London, UK, 2016; pp. 169–171.
13. Choudary, B.M.; Reddy, P.N. Selective Markovnikov's addition of trifluoroacetic acid to alkenes using vanadium(V) oxide. *J. Chem. Soc. Chem.* **1993**, *4*, 405–407. <https://doi.org/10.1039/c39930000405>.
14. Collis, A.E.; Horváth, I.T. Heterogenization of homogeneous catalytic systems. *Catal. Sci. Technol.* **2011**, *1*, 912–919. <https://doi.org/10.1039/c1cy00174d>.
15. Sharma, R.; Kabra, B.; Vaidya, V. Photochemical oxidation of 2-imidazolidinethione by uranyl acetate. *Asian J. Chem.* **1996**, *8*, 500–504.
16. Selishchev, D.S.; Filippov, T.N.; Lyulyukin, M.N.; Kozlov, D.V. Uranyl-modified TiO<sub>2</sub> for complete photocatalytic oxidation of volatile organic compounds under UV and visible light. *Chem. Eng. J.* **2019**, *370*, 1440–1449. <https://doi.org/10.1016/j.cej.2019.03.280>.
17. Hu, D.; Jiang, X. Stepwise benzylic oxygenation via uranyl-photocatalysis. *Green Chem.* **2022**, *24*, 124–129. <https://doi.org/10.1039/D1GC04042A>.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.