

Synthesis and Characterization of Thymol-Based Hydrophobic Deep Eutectic Solvents

Deborah Oluwatomilola Adeoye ^{a, c*}, Zaharaddeen Sani Gano^b, Omar Umar Ahmed^d, Suleiman Mohammed Shu-waa, Abdulazeez Yusuf Atta^a and Baba Yakubu Jubril^a

a – Department of Chemical Engineering, Ahmadu Bello University, Zaria, Kaduna State, Nigeria.

b – National Research Institute for Chemical Technology, Zaria, Kaduna State, Nigeria.

c – Nigerian Institute of Leather and Science Technology, Zaria, Kaduna State, Nigeria.

d – Department of Chemical and Petroleum Engineering, Bayero University, Kano, Nigeria.

* Correspondence: Author: yawehst2006@gmail.com

Abstract: The evolution of Hydrophobic Deep Eutectic Solvents (HDESs) has expanded the applications of the new generation solvents known as Deep Eutectic Solvents (DES) to include water-based operations. How stable they are in aqueous media qualifies them to be categorized as hydrophobic. This will also determine whether they are appropriate materials for water-based industrial processes or not; or whether they end up constituting a greater pollution load to those processes due to the leaching of their precursors into the aqueous media when used. This work sought to prepare HDESs from a monoterpene (thymol), and three long-chain organic acids (octanoic acid, decanoic acid, and dodecanoic acid). The physicochemical characteristics of the prepared HDESs were investigated. Thereafter, their moisture absorption capacity and stability in an aqueous environment were determined to ascertain whether they are hydrophobic as predicted.

Keywords: Hydrophobic Deep Eutectic Solvent; Long chain organic acids; Monoterpenes; Moisture content/Moisture absorption capacity

Introduction

Deep Eutectic Solvents (DESs), also known as new-generation ionic liquids are green solvents that are gradually taking over the roles of other conventional solvents. They are easy to prepare, release low volatile organic compounds, and have low cost (Francisco et al., 2013; Paiva et al., 2014). This gives them an edge over ionic liquids. DESs are defined as solvents that are produced by the interaction of two or more components: Hydrogen Bond Donor (HBD), and Hydrogen Bond Acceptor (HBA), generally associated via hydrogen bonding producing a liquid whose melting point is lower than that of the individual components. This hydrogen bond interaction produces a distinct chemical entity with a lower melting point than its precursors. So far, many DESs researched about are hydrophilic. This makes them unstable in aqueous environments, thereby limiting their applications (Passos et al., 2016; Silva et al., 2016). Because of the increase in the applications of DESs, its synthesis is currently the subject of growing interest. This has resulted in a recent investigation into the synthesis of hydrophobic Deep Eutectic Solvents (HDESs).

Synthesis of HDESs is achieved by using either water-insoluble or weakly water-soluble components. HBAs for HDES synthesis can be classified into two categories. The first is composed of ionic compounds (e.g. tetra-alkyl quaternary ammonium or phosphorous salts), while the second is composed of non-ionic compounds (e.g. monoterpenes). In the evolution of HDES synthesis, those synthesized from the combination of hydrophobic

components alone were found to be more stable in aqueous environments than those synthesized from hydrophobic and hydrophilic parts. Hydrophilic components in HDESs were soluble in aqueous phases, which reduced their stability (due to the leaching of some portion of the hydrophilic DESs) in such an environment (C. Florindo et al., 2017).

Suitable application of HDESs can only be achieved through a thorough knowledge and interpretation of their mechanism of interaction and physicochemical properties. These properties are usually attributed to different HBA and HBD combinations which create unique interactions (Chen et al., 2016). Properties such as density, viscosity, surface tension, and melting/freezing point depend on the HDES structure, while moisture absorption capacity and thermal stability determine appropriate application options. This paper sought to investigate the fundamental physicochemical properties of three (3) non-ionic HDESs prepared from thymol and long-chain saturated organic acids.

2.0. Method

2.1. Hydrophobic Deep Eutectic Solvent Synthesis

Thymol (99.5% assay) was purchased from BDH chemicals, Octanoic acid (99% assay), Decanoic acid (98% assay), and Dodecanoic acid (99% assay) from Loba Chemie. In this work, binary mixtures of thymol and carboxylic acids (octanoic acid, decanoic acid, and dodecanoic acid) were prepared by adding the components into glass vessels (at the molar ratios as contained in Table 1), using a Mettler Toledo analytical weighing balance (ME 204E). The solid mixtures were melted under stirring on a magnetic stirrer hot plate at 80°C until a homogeneous liquid mixture was obtained and cooled to room temperature.

Table 1. Composition of different hydrophobic Deep Eutectic Solvents.

HBA	HBD	Abbreviation	Mole Ratio	Physical appearance
Thymol	Octanoic acid	TC ₈	1:1	Pale yellow liquid
	Decanoic acid	TC ₁₀	1:1	Colourless liquid
	Dodecanoic acid	TC ₁₂	1:1	Golden yellow liquid

3.0. Results and Discussion

3.1. Fourier Transform Infrared of HDESs

The hydrogen bond is the primary intermolecular interaction between terpenes and organic acids in the formation of HDES (Ali et al., 2020; Zhang et al., 2012). Fourier Transform Infrared (FTIR) spectra of the different terpenes and organic acids and the resultant HDES from their combinations were determined to investigate and ascertain this interaction. As represented in Figures 1-3, all thymol-based HDESs also reflected shifts in the initial OH stretching band of their HBA (thymol) from 3175.7 cm⁻¹ to 3418 cm⁻¹ in TC₈, and TC₁₀, while in TC₁₂, the OH stretching band shifted to 3406.8 cm⁻¹. Also, there were shifts from the initial C=O bands in C₁₀ and C₁₂ from 1692.2 cm⁻¹ to 1707.1 cm⁻¹ in TC₁₀ and TC₁₂, while TC₈ maintained its original C=O band peak but had an increase in its intensity. Results obtained were compared with and validated by previous works, which showed similar shifts to higher wavelengths for the OH band of the monoterpene and the CO bands of the organic acid; thus depicting interactions between the components which resulted in the formation of DESs (Ibrahim et al., 2019; Mjalli et al., 2017; Santana et al., 2019).




Figure 1: FTIR spectra of Thymol, Octanoic acid, and TC8 HDES




Figure 2: FTIR spectra of Thymol, Decanoic acid, and TC10 HDES




Figure 3: FTIR spectra of Thymol, Dodecanoic acid, and TC12 HDES

In FTIR, an increase in the peak intensity usually means an increase in the amount (per unit volume) of the functional group associated with the molecular bond. Therefore, the increase in peak intensity and the various peak shifts observed in the spectra of the HDESs can be attributed to the intermolecular hydrogen bonding between the HBDs and HBAs.

3.2. Physicochemical properties

i. Density

Density is a crucial property of solvents that greatly influences dissolution, reaction and separation processes, determining their viability. As seen in Figure 4, the density of all thymol-based HDESs ranged between 0.8820 - 0.9357 kg/l. All HDESs revealed a linear decrease in density with a consistent increase in temperature. Hydrophobic deep eutectic solvents have been reported to possess lower densities than water (Catarina Florindo et al., 2019). The density of a deep eutectic solvent reveals a temperature-dependent behaviour, which decreases linearly with increasing temperature (Cui et al., 2017; C. Florindo et al., 2014; Ibrahim et al., 2019; Shahbaz et al., 2012). According

to (Catarina Florindo et al., 2019), the densities of thymol-based DES were ≥ 0.925 kg/l at 30°C. Moreover, density depends on the choice of the hydrogen bond donor (Andrew P. Abbott, Barron, et al., 2007; Andrew P. Abbott, Harris, et al., 2007; Cui et al., 2017; C. Florindo et al., 2014; García et al., 2015), and the molar ratio (Andrew P. Abbott et al., 2011).




Figure 4: Density-temperature graph for Thymol based HDESs

i. Viscosity

Viscosities of the prepared HDESs were observed to be temperature-dependent. It decreased linearly with corresponding temperature increases. As seen in Figure 5, the viscosity of thymol-based HDES increased from TC₈ to TC₁₂ (TC₈ < TC₁₀ < TC₁₂) and was between 7 – 17 mPa.s. The results above align with results obtained from previous work (Catarina Florindo et al., 2019). Also, from the results obtained, it was observed that an increase in the alkyl chain of the carboxylic acid (HBD) resulted in the corresponding rise in the viscosities of the HDESs. Noteworthy is the extraordinarily low viscosities (<20 mPa.s) of HDESs based on fatty acids combined with menthol and thymol. These overcame one of the significant drawbacks of hydrophilic DES, where for example, a viscosity of 859.45 mPa.s was attained for ChCl: urea (1:2) (Shekaari et al., 2017).




Figure 5: Viscosity-temperature graph for Thymol based HDESs

The viscosity of a eutectic mixture is affected by the nature of its components (Andrew P. Abbott, Barron, et al., 2007; D'Agostino et al., 2011), their molar ratio (Andrew P. Abbott et al., 2011), the temperature (A. P. Abbott et al., 2004; Andrew P. Abbott et al., 2003, 2006; Andrew P. Abbott et al., 2004; Dai et al., 2015; Kareem et al., 2010), and the water content (D'Agostino et al., 2011; Dai et al., 2015; Du et al., 2016; C. Florindo et al., 2014; Shah & Mjalli, 2014).

i. Surface Tension

Surface tension is an essential property since it is highly dependent on the intensity of the intermolecular forces taking place between the hydrogen bond donor and the hydrogen bond acceptor. It also determines the suitability of HDES in interfacial processes in which mass transfer occurs. In Figure 6, the surface tension of TC₈, TC₁₀, and TC₁₂ falls

between 26.0 – 28.5 mN/m. The surface tension of the HDESs was also temperature-dependent as it decreased with increasing temperature. Studies have shown that surface tension decreases linearly with increasing temperature (García et al., 2015; Lapeña et al., 2019; Nunes et al., 2019). The surface tension of all HDES was observed to be <30mN/m, and as exemplified in previous works, decreased with increasing temperature.




Figure 6: Surface tension-temperature graph for Thymol based HDESs

3.3. Hydrophobicity test

Figure 7 shows the image of the HDESs immediately after agitation. An oil-in-water emulsion-like mixture was observed. The samples were left for 24 hours to see if the emulsion formed any phase separation. After 24 hours, clear and distinctive phase separation was noticed with water (more dense phase) seen at the bottom of the bottle and the HDES (less dense phase) being on top, as seen in Figure 8.




Figure 7: HDESs in water immediately after agitation




Figure 8: HDESs in water 24 hours after agitation

Figure 9 is a chart indicating the moisture content determined in each HDES after its interaction with water for 24 hours. The HDESs TC₈, TC₁₀ and TC₁₂ had the following moisture content 7.80%, 3.73% and 3.10% respectively. It could be seen that thymol-based HDESs had a lower moisture absorption capacity than their menthol-based counterparts. It was also reported by (Catarina Florindo et al., 2019), that an increase in the alkyl chain of the HBDs resulted in increased hydrophobicity of the HDESs.

It is highly recommended to determine the chemical stability of deep eutectic solvents that find applications in aqueous environments to ascertain that there is no contamination of the water phase with the DES and no loss of the DES structure due to water absorption. The ability of an HDES to maintain its structure and not lose its integrity in the presence of water is of enormous importance.




Figure 9: Percentage Moisture content graph for HDESs (test for hydrophobicity)

4.0. Conclusion

Three (3) hydrophobic deep eutectic solvents from thymol and long-chain organic acids were successfully prepared, all in ratio 1:1. All were liquids at room temperature, with TC₈ having a pale yellow colour, TC₁₀ being colourless and TC₁₂ having a golden yellow colour. The prepared HDESs were characterized using FTIR, and the spectra revealed remarkable shifts in the O-H stretching bands and C=O stretching bands of the HDESs compared with their precursors. The changes in the OH stretching bands and C=O stretching bands resulted from the intermolecular hydrogen bond formed between the starting materials to give HDES.

Also, physicochemical analysis was carried out on the synthesized HDESs, which revealed their excellent characteristics. This better informs researchers on their best areas of application, potentially in the removal of recalcitrant aromatic contaminants in wastewater. The density of thymol-based HDESs was within 0.925 – 0.940 kg/l; all densities were found to decrease with an increase in temperature. The viscosity of the HDESs was found to be <20 mPa.s and decreased with an increase in temperature. Similar to viscosity, the surface tension of HDESs also reduced with an increase in temperature and was observed to be <30 mN/m. Finally, the extent of moisture absorption into the matrix of the HDESs was determined to be between 3.10 – 7.80 %. It was observed that the degree of hydrophobicity increased with an increase in the alkyl chain of the organic acids (HBDs).

References

- Abbott, A. P., Boothby, D., Capper, G., Davies, D. L., & Rasheed, R. (2004). Deep Eutectic Solvents Formed Between Choline Chloride and Carboxylic Acids. *Journal of American Chemical Society*.
- Abbott, Andrew P., Barron, J. C., Ryder, K. S., & Wilson, D. (2007). Eutectic-based ionic liquids with metal-containing anions and cations. *Chemistry - A European Journal*, 13(22), 6495–6501. <https://doi.org/10.1002/chem.200601738>
- Abbott, Andrew P., Capper, G., Davies, D. L., Rasheed, R. K., & Tambyrajah, V. (2003). Novel solvent properties of choline chloride/urea mixtures. *Chemical Communications*, 1, 70–71. <https://doi.org/10.1039/b210714g>
- Abbott, Andrew P., Capper, G., & Gray, S. (2006). Design of improved deep eutectic solvents using hole theory. *ChemPhysChem*, 7(4), 803–806. <https://doi.org/10.1002/cphc.200500489>
- Abbott, Andrew P., Harris, R. C., & Ryder, K. S. (2007). Application of hole theory to define ionic liquids by their transport properties. *Journal of Physical Chemistry B*, 111(18), 4910–4913. <https://doi.org/10.1021/jp0671998>
- Abbott, Andrew P., Capper, G., Davies, D. L., & Rasheed, R. K. (2004). *Ionic Liquid Analogues Formed from Hydrated Metal Salts*. 3769–3774. <https://doi.org/10.1002/chem.200400127>
- Abbott, Andrew P., Harris, R. C., Ryder, K. S., Agostino, C. D., Gladden, F., & Mantle, M. D. (2011). *Green Chemistry Glycerol eutectics as sustainable solvent systems* †. 82–90. <https://doi.org/10.1039/c0gc00395f>
- Ali, H. H., Ghareeb, M. M., Al-Remawi, M., & T. Al-Akayleh, F. (2020). New insight into sin7gle phase formation of capric acid/menthol eutectic mixtures by Fourier-transform infrared spectroscopy and differential scanning calorimetry. *Tropical Journal of Pharmaceutical Research*, 19(2), 361–369. <https://doi.org/10.4314/tjpr.v19i2.19>
- Chen, J., Liu, M., Wang, Q., Du, H., & Zhang, L. (2016). Deep eutectic Solvent-Based Microwave-Assisted method for extraction of hydrophilic and hydrophobic components from radix salviae miltiorrhizae. *Molecules*, 21(10). <https://doi.org/10.3390/molecules21101383>
- Cui, Y., Li, C., Yin, J., Li, S., Jia, Y., & Bao, M. (2017). Design, synthesis and properties of acidic deep eutectic solvents based on choline chloride. In *Journal of Molecular Liquids* (Vol. 236). Elsevier B.V. <https://doi.org/10.1016/j.molliq.2017.04.052>
- D'Agostino, C., Harris, R. C., Abbott, A. P., Gladden, L. F., & Mantle, M. D. (2011). Molecular motion and ion diffusion in choline chloride based deep eutectic solvents studied by 1H pulsed field gradient NMR spectroscopy. *Physical Chemistry Chemical Physics*, 13(48), 21383–21391. <https://doi.org/10.1039/c1cp22554e>
- Dai, Y., Witkamp, G. J., Verpoorte, R., & Choi, Y. H. (2015). Tailoring properties of natural deep eutectic solvents with water to facilitate their applications. *Food Chemistry*, 187, 14–19. <https://doi.org/10.1016/j.foodchem.2015.03.123>
- Du, C., Zhao, B., Chen, X. B., Birbilis, N., & Yang, H. (2016). Effect of water presence on choline chloride-2urea ionic liquid and coating platings from the hydrated ionic liquid. *Scientific Reports*, 6(July), 1–14. <https://doi.org/10.1038/srep29225>

14. Florindo, C., Oliveira, F. S., Rebelo, L. P. N., Fernandes, A. M., & Marrucho, I. M. (2014). Insights into the synthesis and properties of deep eutectic solvents based on cholinium chloride and carboxylic acids. *ACS Sustainable Chemistry and Engineering*. <https://doi.org/10.1021/sc500439w>
15. Florindo, C., Branco, L. C., & Marrucho, I. M. (2017). Development of hydrophobic deep eutectic solvents for extraction of pesticides from aqueous environments. *Fluid Phase Equilibria*, *448*, 135–142. <https://doi.org/10.1016/j.fluid.2017.04.002>
16. Florindo, Catarina, Branco, L. C., & Marrucho, I. M. (2019). Quest for Green-Solvent Design: From Hydrophilic to Hydrophobic (Deep) Eutectic Solvents. *ChemSusChem*, *12*(8), 1549–1559. <https://doi.org/10.1002/cssc.201900147>
17. Francisco, M., Van Den Bruinhorst, A., & Kroon, M. C. (2013). Low-transition-temperature mixtures (LTTMs): A new generation of designer solvents. *Angewandte Chemie - International Edition*, *52*(11), 3074–3085. <https://doi.org/10.1002/anie.201207548>
18. García, G., Aparicio, S., Ullah, R., & Atilhan, M. (2015). Deep eutectic solvents: Physicochemical properties and gas separation applications. *Energy and Fuels*, *29*(4), 2616–2644. <https://doi.org/10.1021/ef5028873>
19. Ibrahim, R. K., Hayyan, M., AlSaadi, M. A., Ibrahim, S., Hayyan, A., & Hashim, M. A. (2019). Physical properties of ethylene glycol-based deep eutectic solvents. *Journal of Molecular Liquids*, *276*, 794–800. <https://doi.org/10.1016/j.molliq.2018.12.032>
20. Kareem, M. A., Mjalli, F. S., Hashim, M. A., & Alnashef, I. M. (2010). Phosphonium-based ionic liquids analogues and their physical properties. *Journal of Chemical and Engineering Data*, *55*(11), 4632–4637. <https://doi.org/10.1021/je100104v>
21. Lapeña, D., Lomba, L., Artal, M., Lafuente, C., & Giner, B. (2019). The NADES glyceline as a potential Green Solvent: A comprehensive study of its thermophysical properties and effect of water inclusion. *Journal of Chemical Thermodynamics*, *128*, 164–172. <https://doi.org/10.1016/j.jct.2018.07.031>
22. Mjalli, F. S., Murshid, G., Al-Zakwani, S., & Hayyan, A. (2017). Monoethanolamine-based deep eutectic solvents, their synthesis and characterization. *Fluid Phase Equilibria*, *448*, 30–40. <https://doi.org/10.1016/j.fluid.2017.03.008>
23. Nunes, R. J., Saramago, B., & Marrucho, I. M. (2019). Surface Tension of dl -Menthol:Octanoic Acid Eutectic Mixtures. *Journal of Chemical and Engineering Data*, *64*(11), 4915–4923. <https://doi.org/10.1021/acs.jced.9b00424>
24. Paiva, A., Craveiro, R., Aroso, I., Martins, M., Reis, R. L., & Duarte, A. R. C. (2014). Natural deep eutectic solvents - Solvents for the 21st century. *ACS Sustainable Chemistry and Engineering*, *2*(5), 1063–1071. <https://doi.org/10.1021/sc500096j>
25. Passos, H., Tavares, D. J. P., Ferreira, A. M., Freire, M. G., & Coutinho, J. A. P. (2016). Are Aqueous Biphasic Systems Composed of Deep Eutectic Solvents Ternary or Quaternary Systems? *ACS Sustainable Chemistry and Engineering*, *4*(5), 2881–2886. <https://doi.org/10.1021/acssuschemeng.6b00485>
26. Santana, A. P. R., Mora-Vargas, J. A., Guimarães, T. G. S., Amaral, C. D. B., Oliveira, A., & Gonzalez, M. H. (2019). Sustainable synthesis of natural deep eutectic solvents (NADES) by different methods. *Journal of Molecular Liquids*, *293*, 111452. <https://doi.org/10.1016/j.molliq.2019.111452>
27. Shah, D., & Mjalli, F. S. (2014). Effect of water on the thermo-physical properties of Reline: An experimental and molecular simulation based approach. *Physical Chemistry Chemical Physics*, *16*(43), 23900–23907. <https://doi.org/10.1039/c4cp02600d>
28. Shahbaz, K., Baroutian, S., Mjalli, F. S., Hashim, M. A., & Alnashef, I. M. (2012). Densities of ammonium and phosphonium based deep eutectic solvents: Prediction using artificial intelligence and group contribution techniques. *Thermochimica Acta*, *527*, 59–66. <https://doi.org/10.1016/j.tca.2011.10.010>
29. Shekaari, H., Zafarani-Moattar, M. T., & Mohammadi, B. (2017). Thermophysical characterization of aqueous deep eutectic solvent (choline chloride/urea) solutions in full ranges of concentration at T = (293.15–323.15) K. *Journal of Molecular Liquids*, *243*, 451–461. <https://doi.org/10.1016/j.molliq.2017.08.051>
30. Silva, N. H. C. S., Pinto, R. J. B., Freire, C. S. R., & Marrucho, I. M. (2016). Production of lysozyme nanofibers using deep eutectic solvent aqueous solutions. *Colloids and Surfaces B: Biointerfaces*, *147*, 36–44. <https://doi.org/10.1016/j.colsurfb.2016.07.005>