

Proceeding Paper

# Hydrothermal Alteration Features Enhancement and Mapping Using High-Resolution Hyperspectral Data <sup>†</sup>

Soufiane Hajaj <sup>1,\*</sup>, Abderrazak El Harti <sup>1</sup>, Amine Jellouli <sup>1</sup>, Saloua Mnissar Himyari <sup>2</sup> and Abderrazak Hamzaoui <sup>2</sup>

<sup>1</sup> Geomatics, Georesources and Environment Laboratory, Faculty of Sciences and Techniques, Sultan Moulay Slimane University, Beni Mellal 23000, Morocco; a.elharti@usms.ma (A.E.H.)

<sup>2</sup> National Office of Hydrocarbons and Mines—Office National des Hydrocarbures et des Mines (ON-HYM), Moulay Hassan Boulevard, Rabat, Morocco

\* Correspondence: soufianehajaj13@gmail.com, soufiane.hajaj@usms.ma (S.HAJAJ)

<sup>†</sup> Presented at the 5th International Electronic Conference on Remote Sensing, 7–21 November 2023; Available online: <https://ecrs2023.sciforum.net/>

**Abstract:** Hydrothermal alteration mapping is considered as widely adopted step in mineral exploration of numerous ore deposits. In this work the wavelength mapping and relative absorption band depth (RBD) method were applied to map hydrothermal alterations in a site from the abandoned mine of Idikel, western Anti-Atlas, Morocco. Fe<sup>2+</sup>/Fe<sup>3+</sup>, Al-OH, Mg-Fe-OH/ CO<sub>3</sub> hydrothermal alteration minerals were targeted based on HyMap airborne imaging spectroscopy data. Using wavelength mapping approach, the 900 to 1205 nm, 2094 to 2217 nm, and 2264 to 2318 nm ranges were selected to map Fe<sup>2+</sup>/Fe<sup>3+</sup>, Al-OH, Mg-Fe-OH/ CO<sub>3</sub> absorption features, respectively. By carefully selecting these spectral ranges, the study aimed to achieve accurate and reliable mapping of hydrothermal alteration features. The highest interpolated depth of Al-OH features was matched with a major cluster of pixels at 2200 nm. Mg-Fe-OH/ CO<sub>3</sub> highest interpolated depth was depicted at 2300 nm. Fe<sup>2+</sup>/Fe<sup>3+</sup> highest interpolated depth was depicted between 900 and 1000. The relative absorption band depth method was also applied to enhance the detectability of hydrothermal alteration minerals. This method involves assessing the depth of absorption bands associated with the target minerals, allowing for a detailed characterization of the alteration features. The combination of both wavelength mapping and enhancement methods contributed to a comprehensive and robust hydrothermal alteration mapping process. The identification of Fe<sup>2+</sup>/Fe<sup>3+</sup>, Al-OH, and Mg-Fe-OH/CO<sub>3</sub> manifestations provided valuable insights into potential mineralization zones within the study area. Overall, this research contributes to the advancement of hydrothermal alteration mapping using hyperspectral data by selecting the required HyMap bands for mapping targeted alterations. The combination of wavelength mapping and enhancement methods proves to be a powerful approach for accurately identifying and characterizing hydrothermal alteration features using specific hyperspectral channels. The findings from this study can aid future mineral exploration endeavors in similar geological settings, providing valuable guidance for locating potential mineral resources in mountainous and challenging terrains.

**Keywords:** hyperspectral; hydrothermal alteration; RBD; wavelength mapping

## 1. Introduction

Hydrothermal alteration mapping is a crucial step in mineral exploration for a wide range of ore deposits. It involves the identification and characterization of mineral alterations in geological formations, providing valuable insights into potential mineralization zones. Indeed, within the realm of hydrothermal alteration minerals, a significant subset exhibits spectral features within the visible and infrared spectrum, spanning the range of 300 to 3000 nanometers [1]. This spectral activity signifies the minerals' capacity to selec-

tively absorb incident electromagnetic radiation at particular wavelengths, while concurrently reflecting the remaining wavelengths. This absorption of energy results in the emergence of distinct "spectral absorption features," which possess diagnostic qualities that can be harnessed for the identification of minerals or mineral assemblages [2]. Recently, imaging spectroscopy data from various airborne and spaceborne platforms demonstrated a high ability in mineral mapping investigations [3–5].

In this study, we applied advanced techniques for hydrothermal alteration mapping using HyMap airborne imaging spectroscopy data in the context of the abandoned Idikel mine site in the western Anti-Atlas region of Morocco. This research focuses on the utilization of wavelength mapping and the relative absorption band depth method to pinpoint hydrothermal alteration minerals, specifically targeting Fe<sup>2+</sup>/Fe<sup>3+</sup>, Al-OH, and Mg-Fe-OH/CO<sub>3</sub> manifestations. By strategically selecting spectral ranges, we aim to achieve precision and reliability in mapping these alterations. Furthermore, we employed the RBD method to enhance the detection of hydrothermal alteration minerals, enabling a more detailed characterization of these features. The combined use of wavelength mapping and enhancement methods results in a comprehensive and robust hydrothermal alteration mapping process.

By identifying and characterizing Fe<sup>2+</sup>/Fe<sup>3+</sup>, Al-OH, and Mg-Fe-OH/CO<sub>3</sub> manifestations, this research provides valuable insights into potential mineralization zones within the study area. It contributes significantly to the advancement of hydrothermal alteration mapping using hyperspectral data by selecting specific HyMap bands for targeted alteration mapping. The integration of wavelength mapping and enhancement techniques is a powerful approach for accurately identifying and characterizing hydrothermal alteration features. This study holds the potential to assist future mineral exploration endeavors in similar geological settings, offering valuable guidance for locating potential mineral resources in challenging terrains, such as mountainous regions.

## 2. Material and Methods

### 2.1. Geological Setting and Alterations

The geology of the Kerdous inlier located in the western Anti Atlas belt, Morocco was principally structured by the Eburnean and Pan-African orogeny [6,7]. The geological map used for the study area located at east Kerdous inlier was digitalized from the geological map of Tafraout, The Paleoproterozoic formations are materialized by metamorphic series, including gneisses, schist, and micaschist. The Neoproterozoic is unconformity deposited, and it is intersected by rhyolitic vulcanites, granites, quartzites, ignimbrites, and Ultimate conglomerates of the Adoudounien base (see Figure 1) [7].

In the eastern region of Kerdous, historical mining operations are evident through the presence of the disused Manganese Idikel mine. Within the Ultimate conglomerates and quartzites lithological units, three distinct sites exhibit favorable conditions for conducting mineral exploration campaigns which is verified by utilizing hyperspectral survey techniques [8]. The close spatial association between the NE-SW and NW-SE lineaments and alteration zones highlighted an important tectonic control of mineralization in the study area [8]. The alteration minerals in the study area include muscovite, dolomite, pyrophyllite, illite, haematite, kaolinite, and montmorillonite. Hydrothermal alteration potentially related to paleo-relief occurrence includes silicification, sericitization, and dolomitization of the host rocks (for additional details, see [8]).




Figure 1. Geological map of the Ameln Valley area [9].

### 2.2. HyMap Data and Remote Sensing Techniques

The HyMap hyperspectral imaging system is originally developed by Integrated Spectronic in Sydney, Australia, and subsequently operated by HyVista [10], captures spectral information across 128 distinct spectral bands spanning the VNIR (Visible and Near-Infrared) and SWIR (Short-Wave Infrared) regions (from 0.45  $\mu\text{m}$  to 2.5  $\mu\text{m}$ ) [11]. The HyMap dataset offers a high spatial resolution of approximately 5 meters and an average spectral resolution of 15 nanometers.

RBD method uses a three-point formulation, wherein the sum of the bands displaying spectral shoulders is divided by the band representing the minimum of the absorption feature [12]. This technique possesses the capability to discern targets even under varying illumination conditions [13]. The RBD approach has been effectively employed to emphasize spectral attributes associated with alteration mineral zones or mineral assemblages, specifically those of the argillic, phyllic, and propylitic [14]. The development of three distinct RBDs within the spectral wavelength range of HyMap has enabled the mapping of manifestations related to  $\text{Fe}^{3+}$ - $\text{Fe}^{2+}$  by dividing the sum of the Band 21 (755 nm) and 51 (1205 nm) by B31 (900) and multiplied by the band 21 (755 nm) by the Band 4 (456 nm).

Additionally, Al-OH is mapped by dividing the sum of the band 101 (2094 nm) and 111 (2268 nm) by the band 108 (2217 nm). The enhancement of Mg-Fe-OH/CO<sub>3</sub> alteration minerals can be performed by dividing the sum of the Band 111 (2268 nm) and 120 (2414 nm) by the band 114 (2318 nm) [8,15]. This capability is grounded in the distinctive absorption features exhibited by these minerals [1]. In the current study, the RBD-derived images/maps were used to target Fe<sup>3+</sup>/Fe<sup>2+</sup> alteration minerals including hematite and goethite showing absorption features from 450 to 1200 nm [16], the Al-OH minerals including kaolinite, Muscovite/illite, pyrophyllite, montmorillonite showing absorption features near 2200 nm [16], and dolomite showing absorption features near 2318 nm [16]. According to alteration minerals absorption features, the employability of these RBDs can be beneficial to map other minerals with common criteria, including calcite, chlorite, and epidote for Mg-Fe-OH/CO<sub>3</sub> alteration minerals, as well as limonite and jarosite for Fe<sup>3+</sup>/Fe<sup>2+</sup> alteration minerals [16].


The minimum wavelength mapper approach relies on the analysis of spectral absorption characteristics, including parameters like position and depth. It facilitates the generation of a visual representation, presented as a color-coded map, which serves as a proxy for mineral composition, mineral abundance, and spatial distribution [17]. The Minimum Wavelength Mapper method excels in computing a proxy for mineral composition and demonstrates its proficiency in detecting compositional alterations through the identification of subtle shifts in wavelength positions. These attributes underscore the method's significance in mineralogical analysis. In contrast, techniques based on mineral endmembers or library spectra, such as the spectral angle mapper (SAM) and linear spectral unmixing (LSU), lack the capacity to discern minor alterations in the wavelength positions of absorption features or represent them as distinct endmembers [18].

### 3. Results and Discussion


Figure 2 shows the results of hydrothermal alteration mapping around the abandoned Idikel mine. Minimum wavelength mapping results exhibit the high abundance of Al-OH alteration in Ultimate conglomerate lithological unit as highlighted in purple color, while this alteration is absent in carbonates units (Figure 2). The high alteration minerals abundance is related essentially to argillic and phyllic alterations [8]. Besides Fe<sup>2+</sup>/Fe<sup>3+</sup> alterations were mapped in blue tones in gneisses and some parts of Ultimate conglomerates units. Based on previous mapping results in the Ameln valley shear zone, the Fe<sup>2+</sup>/Fe<sup>3+</sup> features could be related to Hematite. Mg-Fe-OH/CO<sub>3</sub> alterations were mapped yellow in carbonates units represented by base limestone and dolomite and lower limestone and dolomite which can be due to dolomite mineral. While a little abundance of Mg-Fe-OH/CO<sub>3</sub> minerals can be detected in Paleoproterozoic schist due to the presence of recent sediments (Figure 2). Figure 3 displays the plot of the absorption features depth as a function of the wavelength position for each result by the wavelength mapping. Figure 3A demonstrated that the interpolated minimum wavelength was centered around 2200 nm. Figure 3B shows several features, the main interpolated depth of the wavelength was identified near 900 nm. However, the presence of several interpolated depths can explain the relative weak abundance of Fe<sup>2+</sup>/Fe<sup>3+</sup> within the study area. Figure 3C shows that the main interpolated depth between 2285 nm and 2300 nm.

For RBD, Al-OH alteration minerals, which are mostly found in the Ultimate conglomerates unit and gneiss unit are depicted as bright pixels on the RBD1 image-map (Figure 2). For Fe<sup>2+</sup>/Fe<sup>3+</sup> alteration minerals, these are often linked to lower limestone and dolomite and the Ultimate conglomerates lithological units. This alteration is enhanced as bright pixels on the RBD2 image-map (Figure 2). In the Adoudounien formations, the RBD3 exhibits a high spatial distribution of Mg-Fe-OH/CO<sub>3</sub> particularly (carbonates) in the lithological units of base limestone and dolomite and lower limestone and dolomite (Figure 2).

The current work method involves assessing the depth of absorption bands associated with the target minerals, allowing a detailed characterization of the alteration features. The combination of both wavelength mapping and enhancement methods contributed to a comprehensive and robust hydrothermal alteration mapping process. The mapping results following minimum wavelength and RBD technique have revealed a global agreement. In some cases, limited information was revealed by using RBD technique for visualizing targeted minerals in the study area, which can be observed by an overspread of bright pixels, surpassing the specified alteration features. Besides, it was demonstrated that establishing the minimum wavelength technique within the three specified intervals proved to be useful in verifying the RBD-derived alteration maps. Accordingly, the detailed insights including the absorption depth percentage and the electromagnetic spectrum positions of the enhanced alteration features have been extracted successfully (See figure 2 and 3). The minimum wavelength mapping method is highly sensitive to small changes in mineral composition or percentage. This sensitivity allows it to create a single color-coded map that serves as a proxy for mineral composition, abundance, and spatial patterns [17]. The identification of Fe<sub>2+</sub>/Fe<sub>3+</sub>, Al-OH, and Mg-Fe-OH/CO<sub>3</sub> manifestations provided valuable insights into potential mineralization zones within the study area.



**Figure 2.** The results of the wavelength mapping and relative absorption band depth for the three targeted hydrothermal alteration features (i.e., Al-OH, Fe<sub>2+</sub>/Fe<sub>3+</sub>, and Mg-Fe-OH/CO<sub>3</sub>, from the left to the right) in the study area.



**Figure 3.** The plot of the absorption features depth as a function of the wavelength position for each result of the wavelength mapping for Al-OH (A), Fe<sup>2+</sup>/Fe<sup>3+</sup> (B), and Mg-Fe-OH/CO<sub>3</sub> (C) hydrothermal alteration minerals.

#### 4. Conclusion

In conclusion, this study has demonstrated the effectiveness of utilizing advanced techniques, including wavelength mapping and the RBD method, for the precise mapping of hydrothermal alterations in the Idikel mine site within the western Anti-Atlas of Morocco. By specifically targeting Fe<sup>2+</sup>/Fe<sup>3+</sup>, Al-OH, and Mg-Fe-OH/CO<sub>3</sub> hydrothermal alteration minerals through HyMap airborne imaging spectroscopy data, we have achieved accurate and reliable mapping of these features. The integration of wavelength mapping and enhancement techniques has proven to be a powerful approach for accurately characterizing hydrothermal alteration features using specific hyperspectral channels. It is highlighted that applying minimum wavelength technique within the three specified intervals provides a useful insights and a further verification of the RBD-derived alteration maps. The current study findings holds the potential to guide future mineral exploration efforts in similar geological settings, particularly in challenging mountainous terrains, facilitating the discovery of potential mineral resources.

**Acknowledgments:** We show appreciation to the ONHYM for providing the HyMap data. Also, the authors would like to thank The 5th International Electronic Conference on Remote Sensing organization committee for providing the opportunity to discuss the present work's findings. The authors extend their gratitude to the reviewers for their valuable feedback, instrumental in the refinement of the paper's findings.

**Conflicts of Interest:** The authors declare that they have no conflict of interest. It is affirmed that the manuscript has been comprehensively reviewed and endorsed by all the named authors.

#### References

1. Clark, R.N.; Gallagher, A.J.; Swayze, G.A. Material absorption band depth mapping of imaging spectrometer data using a complete band shape least-squares fit with library reference spectra. In Proceedings of the Proceedings of the second airborne visible/infrared imaging spectrometer (AVIRIS) workshop, 1990; pp. 176-186.
2. Hecker, C.; van Ruitenbeek, F.J.; van der Werff, H.M.; Bakker, W.H.; Hewson, R.D.; van der Meer, F.D. Spectral absorption feature analysis for finding ore: A tutorial on using the method in geological remote sensing. *IEEE geoscience and remote sensing magazine* 2019, 7, 51-71.
3. Cardoso-Fernandes, J.; Santos, D.; Lima, A.; Teodoro, A.C. Assessing the PRISMA Potential for Mineral Exploration to Vector Low-Grade Lithium Deposits. In Proceedings of the IGARSS 2022-2022 IEEE International Geoscience and Remote Sensing Symposium, 2022; pp. 5985-5988.
4. Pour, A.B.; Guha, A.; Crispini, L.; Chatterjee, S. Editorial for the Special Issue Entitled Hyperspectral Remote Sensing from Spaceborne and Low-Altitude Aerial/Drone-Based Platforms—Differences in Approaches, Data Processing Methods, and Applications. 2023, 15, 5119.

5. Yao, F.; Liu, S.; Wang, D.; Geng, X.; Wang, C.; Wang, Y. Review on the development of multi-and hyperspectral Remote sensing technology for exploration of copper-gold deposits. *Ore Geology Reviews* 2023, 105732.
6. Choubert, G. Histoire géologique du Précambrien de l'Anti-Atlas; Éditions du Service géologique du Maroc: 1964.
7. Hassenforder, B. Evolution tectono-métamorphique du socle du Kerdous (Anti-Atlas occidental, Maroc) dans le cadre des orogènes éburnéenne et pan-africaine. *Sciences Géologiques, bulletins et mémoires* 1978, 31, 21-31.
8. Hajaj, S.; El Harti, A.; Jellouli, A.; Pour, A.B.; Himyari, S.M.; Hamzaoui, A.; Bensalah, M.K.; Benaouiss, N.; Hashim, M. HyMap imagery for copper and manganese prospecting in the east of Ameln valley shear zone (Kerdous inlier, western Anti-Atlas, Morocco). *Journal of Spatial Science* 2023, 1-22.
9. Hajaj, S.; El Harti, A.; Jellouli, A.; Pour, A.B.; Mnissar Himyari, S.; Hamzaoui, A.; Hashim, M. Evaluating the Performance of Machine Learning and Deep Learning Techniques to HyMap Imagery for Lithological Mapping in a Semi-Arid Region: Case Study from Western Anti-Atlas, Morocco. *Minerals* 2023, 13, 766.
10. Cocks, T.; Jenssen, R.; Stewart, A.; Wilson, I.; Shields, T. The HyMap™ airborne hyperspectral sensor: The system, calibration and performance. In *Proceedings of the Proceedings of the 1st EARSeL workshop on Imaging Spectroscopy, 1998*; pp. 37-42.
11. Hausknecht, P. Hymap Data L2/3 Processing, J Report for the Geological Survey of Namibia 2005, 1, 48.
12. Crowley, J.K.; Brickey, D.W.; Rowan, L.C. Airborne imaging spectrometer data of the Ruby Mountains, Montana: mineral discrimination using relative absorption band-depth images. *Remote Sensing of Environment* 1989, 29, 121-134.
13. Lillesand, T.; Kiefer, R.W.; Chipman, J. *Remote sensing and image interpretation*; John Wiley & Sons: 2015.
14. Beiranvand Pour, A.; S. Park, T.-Y.; Park, Y.; Hong, J.K.; M Muslim, A.; Läufer, A.; Crispini, L.; Pradhan, B.; Zoheir, B.; Rahmani, O. Landsat-8, advanced spaceborne thermal emission and reflection radiometer, and WorldView-3 multispectral satellite imagery for prospecting copper-gold mineralization in the northeastern Inglefield Mobile Belt (IMB), northwest Greenland. *Remote Sensing* 2019, 11, 2430.
15. Hajaj, S.; El Harti, A.; Jellouli, A.; Pour, A.B.; Himyari, S.M.; Hamzaoui, A.; Bensalah, M.K.; Benaouis, N.; Hashim, M. HyMap airborne imaging spectroscopy for mineral potential mapping of cupriferous mineralization in a semi-arid region based on pixel/sub-pixel hydrothermal alteration minerals mapping—A case study; *Copernicus Meetings*: 2023.
16. Hunt, G.R. Spectral signatures of particulate minerals in the visible and near infrared. *Geophysics* 1977, 42, 501-513.
17. Bakker, W.; van Ruitenbeek, F.; van der Werff, H. Hyperspectral image mapping by automatic color coding of absorption features. In *Proceedings of the 7th EARSEL workshop of the Special Interest Group in imaging spectroscopy: final programme, 11-13 April 2011, Edinburgh, UK, 2011*.
18. Portela, B.; Sepp, M.D.; van Ruitenbeek, F.J.; Hecker, C.; Dilles, J.H. Using hyperspectral imagery for identification of pyrophyllite-muscovite intergrowths and alunite in the shallow epithermal environment of the Yerington porphyry copper district. *Ore geology reviews* 2021, 131, 104012.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.