

Management and quality of irrigation water in the sustainable Selenium-enriched rice (*Oryza sativa* L.) production †

Ana Coelho Marques ^{1,2,*}, Diana Daccak ^{1,2}, Inês Carmo Luís ^{1,2}, Ana Rita F. Coelho ^{1,2}, Cláudia Campos Pessoa ^{1,2}, Manuela Simões ^{1,2}, Paula Scotti-Campos ^{2,3}, Ana Sofia Almeida ^{2,4}, Maria Graça Brito ^{1,2}, José Carlos Kullberg ^{1,2}, Carlos Galhano ^{1,2}, Fernando Reboredo ^{1,2}, Paulo Legoinha ^{1,2}, José C. Ramalho ^{2,5}, José Manuel N. Semedo ^{2,3}, Isabel P. Pais ^{2,3}, Maria Manuela Silva ^{1,2}, Maria Fernanda Pessoa ^{1,2}, Lourenço Palha ⁶, Cátia Silva ⁶ and Fernando C. Lidon ^{1,2}

¹ Earth Sciences Department, NOVA School of Science and Technology (FCT NOVA), Campus de Caparica, 2829-516 Caparica, Portugal; d.daccak@campus.fct.unl.pt (D.D.); idc.rodrigues@campus.fct.unl.pt (I.C.L.); arf.coelho@campus.fct.unl.pt (A.R.F.C.); c.pessoa@campus.fct.unl.pt (C.C.P.); mmsr@fct.unl.pt (M.S.); mgb@fct.unl.pt (M.G.B.); jck@fct.unl.pt (J.C.K.); acag@fct.unl.pt (C.G.); fhr@fct.unl.pt (F.R.); pal@fct.unl.pt (P.L.); mma.silva@fct.unl.pt (M.M.S.); mfgp@fct.unl.pt (M.F.P.); fjl@fct.unl.pt (F.C.L.)

² GeoBioTec Research Center, NOVA School of Science and Technology (FCT NOVA), Caparica, Portugal; paula.scotti@iniav.pt (P.S.C.); sofia.almeida@iniav.pt (A.S.A.); cochichor@mail.telepac.pt (J.C.R.); jose.semedo@iniav.pt (J.M.N.S.); isabel.pais@iniav.pt (I.P.P.)

³ Instituto Nacional de Investigação Agrária e Veterinária, I.P. (INIAV), Avenida da República, Quinta do Marquês, 2780-157 Oeiras, Portugal;

⁴ Instituto Nacional de Investigação Agrária e Veterinária, I.P. (INIAV), Estrada de Gil Vaz 6, 7351-901 Elvas, Portugal;

⁵ PlantStress & Biodiversity Lab., Centro de Estudos Florestais (CEF), Associate Laboratory TERRA, Instituto Superior Agronomia (ISA), Universidade de Lisboa (ULisboa), Quinta do Marquês, Av. República, 2784-505 Oeiras and Tapada da Ajuda, 1349-017 Lisboa, Portugal;

⁶ Centro de Competências do Arroz (COTArroz), 2120-014 Salvaterra de Magos, Portugal; l.palha@cotarroz.pt (L.P.); catia.leonardo.silva@gmail.com (C.S.)

* Correspondence: amc.marques@campus.fct.unl.pt; Tel.: +351 212 948 573

† Presented at the 2nd International Online Conference on Agriculture, 01–15 November 2023; Available online: <https://sciforum.net/conference/IOCAG2023>

Abstract: There is a growing need for strategic actions involving efficient water use, sustainable agricultural production, and food security. Agricultural productivity can be improved through good agricultural practices based on water quality management, new genetically modified resources, and using precision agriculture. This study aimed to monitor the crop water (supply, irrigation, and flooding) of an advanced rice (*Oryza sativa* L.) line of the breeding program (OP 1509) subjected to Selenium (Se) enrichment. Water lines in paddy rice field were monitored by Unmanned Aerial Vehicles (UAVs). The parameters of pH, pHS, electrical conductivity, temperature, HCO₃⁻, Cl⁻, SO₄²⁻, PO₄³⁻, Na⁺, K⁺, Ca²⁺, and Mg²⁺ were analyzed. According to the Piper diagram, the samples were classified as sodium chloride bicarbonate (supply) and sodium bicarbonate chloride (irrigation and flooding). The Langelier Saturation Index (LSI) was calculated and indicated that waters are good to use in agricultural practices. According to the Wilcox classification, regarding agriculture use, the samples were classified as C2S1 (supply and irrigation) and C3S1 (flooding). The Selenium contents were analyzed by atomic absorption and significant differences were observed in rice grains, with a maximum content of 10 mg.kg⁻¹. In conclusion, water quality is in accordance with the parameters for use in this crop and the workflow used improved the grain quality.

Keywords: Langelier Saturation Index; *Oryza sativa* L.; Piper and Wilcox diagram; Precision agriculture; Water Management and Quality

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Last-name

Published: date

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

There is a growing need for strategic actions involving the efficient use of water, sustainable agricultural production, and food security, considering climate change [1]. Efficient use and sustainable supplies of water in the agriculture, aquaculture, live-stock, and agroforestry sectors are the future way [1]. Rice (*Oryza sativa* L.) is one of the most important crops in the world [2] and requires a high amount of water to grow [3]. For this reason, it is important to know the quality of the water. The water can be classified according to the chemical composition by Piper or /and Stiff diagrams in functions of the relative predomination of specific ions, namely of the main anions and cations. The suitability of the water for agricultural use is based on the Wilcox diagram where the conductivity ranges between classes C1–C4 and the Sodium Adsorption Ratio index (SAR) between classes S1–S4. The SAR index is very important to classifying irrigation waters, considering that if it is high in Na^+ and low in Ca^{2+} , the cation exchange complex can become saturated with Na^+ [4] in water and affect crop growth. Thus, the use of good agricultural practices based on pest control, soil, and water management using precision agriculture, and new genetic materials can contribute to increased productivity. Plant breeding allows the development of plants with desirable agronomic characteristics, such as tolerance to climatic conditions, disease resistance, and higher grain yield. In this way, it is possible to improve the quality of the final product [5]. Biofortification is an effective and sustainable method for increasing micronutrient levels in rice to mitigate deficiencies in people who primarily eat rice [6].

In this sense, this study aims management of irrigation water based on its composition and pick up culture images with drones to support the sustainable production of rice with added value.

2. Materials and Methods

2.1. Experimental Fields and Biofortification Workflow

In a paddy rice field located at the center of Portugal (Salvaterra de Magos), an advanced rice line (OP 1509) of the breeding program carried out by Instituto Nacional de Investigação Agrária e Veterinária (INIAV) was tested. The experimental design was a factorial arrangement: 2 concentrations (0 and $300\text{g Se}\cdot\text{ha}^{-1}$), 2 Se forms (sodium selenate and sodium selenite), and 4 replicates in a total of 16 plots. Selenium applications included 3 foliar applications, which occurred at the end of booting, anthesis, and at the milky grain stage. The 1st application was $500\text{g Se}\cdot\text{ha}^{-1}$, however, the plants showed symptoms of toxicity, which is why the remaining applications were with $300\text{g Se}\cdot\text{ha}^{-1}$. The trial was carried out from 4th June to 23rd October 2019.

2.2. Precision Agriculture–Characterization of Water Lines

The experimental rice field was flown with Unmanned Aerial Vehicles (UAVs) synchronized by Global Position System (GPS), according to Coelho et al. [7]. The flight was performed (18th July) after crop implementation to classify surface water drainage (water lines) in the experimental paddy rice field according to Direção Geral de Agricultura e Desenvolvimento Rural [8].

2.3. Water Analysis

Physical parameters (pH, electrical conductivity, and temperature) were measured with a multiparameter analyzer (Consort C6030) coupled with SP21 and SK20 T electrodes. The chemical parameters such as bicarbonate (HCO_3^-) were analyzed by titration using 100 mL of sample, 0.1 % methyl orange, and 0.1 N hydrochloric acid as titrant [9]. Photometry was carried out using a spectroquant (NOVA 60, Merck, Darmstadt–Germany) to quantify the anions such as chloride (Cl^-), sulphate (SO_4^{2-}), and phosphate (PO_4^{3-}) using the specific kits 1.14897, 1.14779 and 1.14842, respectively. The ionic chromatography was used to measure cations calcium (Ca^{2+}) magnesium (Mg^{2+}), sodium (Na^+), and potassium (K^+) using a chromatograph (Metrohm, 761 Compact IC) and pre-column

(Metrosep cation 1-2, 6.1010.000). The 10 μL of sample previously prepared in eluent were injected at a flow rate of 1.00 mL/min. [10]. Water composition were projected on Piper diagram in order to classify water facies hydrochemistry [11], on Wilcox diagram (considering Sodium Adsorption Ratio, SAR) to assess water irrigation purposes [12] and on Stiff Diagram which represents an irregular polygonal pattern of anion and cation abundance and allows water samples to be compared [4]. The Piper, Wilcox and Stiff diagrams were obtained using the Grapher software (version 16.3.410). Langelier Saturation Index (LSI) and equilibrium pHs were calculated to assess the corrosive or incrusting action [13].

2.4. Atomic Absorption–Quantification of Selenium in Paddy Rice Grains

Selenium content was analyzed by atomic absorption spectrophotometry (Perkin Elmer AAnalyst 200), followed the methods described by Carrondo et al. and Reboledo et al. [14,15]. In short, an acid digestion procedure was carried out with a mix of HNO_3 – HCl with a ratio of 4:1.

2.4. Statistical Analysis

Data were statistically analyzed using a One-way ANOVA ($p \leq 0.05$) to assess differences among treatments. A Tukey's test was performed for mean comparison using IBM SPSS Statistics 20 program.

3. Results

The geomorphology of the experimental rice field is characterized by different elevations, the eastern part being higher (Figure 1). Thus, the elevation was measured to determine water drainage trends. Accordingly, the water lines created in the experimental field follow a westerly direction (Figure 1).



Figure 1. Orthophotomap of elevation and water lines in paddy rice field of OP 1509 genotype at 18th July.

The pH of the water samples ranged from 6.3–6.9, while the pHs ranged between 8.4–8.9 (Table 1). Electrical conductivity of the water varied from 376 $\mu\text{S}/\text{cm}$ to 420 $\mu\text{S}/\text{cm}$, in supply (A), and irrigation (B), respectively. Moreover, the high EC is observed in flooding (C) sample such as 800 $\mu\text{S}/\text{cm}$ (Table 1). In sample A, the concentration of bicarbonate (HCO_3^-), chloride (Cl^-), and sulphate (SO_4^{2-}) was 1.1 meq/L, 1.4 meq/L, and 0.9 meq/L, respectively (Figure 2). Regarding B and C samples, the concentration of HCO_3^- , Cl^- , and SO_4^{2-} ranged from 0.9–1.4 meq/L, 2.1–3.7 meq/L, and 0.8–1.5 meq/L, respectively. In both water samples, the concentration of phosphate (PO_4^{3-}) is low (< 0.04 meq/L). Additionally, the Langelier Saturation Index (LSI) was the same for samples A and C (-1.8) while that of sample B was -2.6 (Table 1).

Table 1. Values of temperature (°C), pH, Electrical Conductivity (EC, $\mu\text{S}/\text{cm}$), anions (meq/L), cations (meq/L), pHs, and Langelier Saturation Index (LSI) in the water supply (A), irrigation water (B), and flooding water (C) samples from the paddy rice field of OP 1509 genotype.

Parameter	A	B	C
Temperature	20	20	20
pH	6.9	6.3	6.6
EC	376	420	800
HCO ₃ ⁻	1.1	0.9	1.4
Cl ⁻	1.4	2.1	3.7
SO ₄ ²⁻	0.9	0.8	1.5
PO ₄ ³⁻	< 0.04	< 0.04	< 0.04
Ca ²⁺	0.5	0.7	1.4
Mg ²⁺	0.4	0.4	0.6
Na ⁺	1.3	1.7	2.7
K ⁺	0.2	0.06	0.4
pH's	8.4	8.9	8.4
LSI	-1.8	-2.6	-1.8

According to the Piper diagram, the chemical composition of the water samples is sodium chloride bicarbonate (A) and sodium bicarbonate chloride (B and C) (Figure 2). By Wilcox classification, water samples A and B are C2S1 while sample C is C3S1. Thus, the Sodium Adsorption Rate Index (SAR) is 1.93, 2.29, and 2.70 for samples A, B, and C respectively (Figure 2).




Figure 2. Classification of hydrochemical facies of waters (A, B, and C samples) and its adequation of irrigation purposes by Piper and Wilcox diagrams.

The Stiff diagrams show that Cl⁻ is the dominant anion and the same for cation Na⁺ followed by Mg²⁺ and Ca²⁺, in both samples (Figure 3). The highest concentrations (cations and anions) are observed in sample C.




Figure 3. Representation of water samples (A, B, and C) in Stiff diagrams.

In the control, Selenium (Se) content was not detected, however, in biofortified grains significant differences were reported (Table 2). Selenite application showed a 10.10 mg.kg⁻¹ of increased while selenate showed a 6.00 mg.kg⁻¹.

Table 2. Mean of values (n = 4) ± standard error of Se content in paddy rice flour of OP 1509 genotype. Letters *a* and *b* revealed significant differences between treatments ($p \leq 0.05$).

Treatment	Concentration (mg.kg ⁻¹)
Control	-
Na ₂ SeO ₄ (300g Se.ha ⁻¹)	6.00 ± 1.09b
Na ₂ SeO ₃ (300g Se.ha ⁻¹)	10.10± 0.75a

4. Discussion

The quality of water used in agriculture affects crops. Thus, to increase macro or microelement contents in a crop, such as rice (*Oryza sativa* L.), it is necessary to preserve the quality of the water used. Therefore, it is extremely important to evaluate the suitability of irrigation water. All crops take water from where it is most readily available (within the rooting depth), however, each crop has its absorption pattern [16]. The geomorphology of the paddy rice field was characterized by orthophotomap where the elevation and the water lines created in the experimental field were observed. The lower elevation zone (west zone) corresponds to the flatter surface and is more favorable to water accumulation and consequently greater infiltration (Figure 1). The higher areas of the experimental field (east zone) represent areas that promote surface water runoff and consequently reduced water infiltration (Figure 1). If surface drainage occurs, it follows the direction of the estimated water lines. The samples showed a pH ranging from 6.3–6.9 (Table 1), which is in line with their use for agricultural purposes [16]. A value outside the normal range may cause a nutritional imbalance or may contain a toxic ion [13]. The salinity hazard is measured by the increase in Electrical Conductivity (EC) because most are the most influential in water quality in crops. In your study, the increase of EC in samples A (376 µS/cm), B (420 µS/cm), and C (800 µS/cm) were probably caused by the incorporation of salts from fertilizers applied in the paddy rice field, which were moved at the time of drainage. According to the literature, EC below 1000 µS/cm is adequate for agriculture [17]. Rice is a moderately sensitive crop to salinity and from 3000 µS/cm there is a reduction in plant growth [18]. Similarly, to EC, the concentration of anions and cations was higher in sample C (Table 1). The high concentration of bicarbonate (HCO₃⁻) in water can increase the sodium (Na⁺) content in the soil due to the precipitation of calcium (Ca²⁺) and magnesium (Mg²⁺). In this study, the HCO₃⁻ content does not present a degree of restriction for the use of water in irrigation because it is below 1.5 meq/L [13]. Also, through the Langelier Saturation Index (LSI), which relates pH to equilibrium pH (pH_s), the values are < 0. Thus, the results suggest that the water is undersaturated in calcium carbonate (CO₃Ca) with a high tendency for corrosive action when circulating in pipes with this composition [19]. According to the Piper diagram, based on the contribution of the cations and anions, the samples were classified as sodium chloride bicarbonate (A) and sodium bicarbonate chloride (B and C) (Figure 2). To classify the water as suitable for irrigation, the Wilcox diagram was plotted [20]. The EC was plotted against the Sodium Adsorption Rate (SAR) index (relative proportion of Na⁺ to Ca²⁺ and Mg²⁺ concentration), which represents the suitability of the water for agricultural use [12]. The values ranged from 1.93–2.70 for samples A and C, respectively (Figure 2). According to the Wilcox classification, all samples have low sodium hazard (S1). Regarding the salinity hazard, samples A (C2S1) and B (C2S1) present medium while sample C (C3S1) presents high. Thus, water with a C2 rating (250–750 µS/cm) can promote plant stress due to the accumulation of salts in the soil and should be used whenever there is a moderate degree of leaching. Also, water in the C3 class (750–2250 µS/cm) affects most plants, requiring good drainage, leaching, and careful irrigation. According to the Stiff diagram the evolution of anions and cations from sample A to B is observed (Figure 3). Rice plants are the most sensitive to humidity, temperature, and water conditions [21]. Based on this, breeding techniques ensure the availability of

varieties adapted to different environmental conditions [5]. The increase of Se depends on the characteristics of each variety, the form of Se used, and concentration [22]. Independently of the foliar pulverization, the results showed a Se increment of 10.10 mg.kg⁻¹ (sodium selenite) and 6.00 mg.kg⁻¹ (sodium selenate) in paddy rice grains (Table 2). Our findings are following the literature that points to selenite as more effective than selenate [23].

Considering that part of the plant development cycle takes place in water, the conditions of irrigation water, supply, and waterlogging are very important for obtaining quality grain. Thus, water monitoring for parameters such as pH, EC, pHs, ISL, and the quantification of anions and cations is important, allowing corrective action to be taken in a timely way.

5. Conclusions

This study focused on monitoring the water management and quality of an experimental field of an advanced rice (*Oryza sativa* L.) line of the breeding program (OP 1509) subjected to Selenium (Se) enrichment. The water lines were monitored using Unmanned Aerial Vehicles (UAVs). Water samples were collected from water supply (A), irrigation water (B), and flooding water (C), and physicochemical composition (pH, EC, anions, and cations concentration) were monitored. The pH at the equilibrium point (pHs) and Langelier Saturation Index (LSI) were also considered. According to the Piper diagram, the samples were classified as sodium chloride bicarbonate (sample A) and sodium bicarbonate chloride (samples B and C), and their suitability for agricultural use was classified as C2S1 (samples A and B) and C3S1 (sample C) according to the Wilcox diagram. Thus, the use of quality water contributed to the production of rice grain enriched with Se (up to 10 mg.kg⁻¹).

Author Contributions: Conceptualization, A.C.M. and F.C.L.; methodology, M.S., M.G.B., J.C.K., C.G. and F.C.L.; investigation, A.C.M., D.D., I.C.L., A.R.F.C. and C.C.P.; formal analysis, A.C.M., D.D., I.C.L., A.R.F.C., C.C.P. and M.S.; software, M.S., J.C.K. and M.G.B.; resources, M.S., M.G.B., J.C.K., C.G., F.R., P.L., J.C.R., J.M.N.S., I.P.P., M.M.S., M.F.P., L.P., C.S. and F.C.L.; writing-original draft preparation, A.C.M.; writing-review and editing, A.C.M. and F.C.L.; supervision, M.S., P.S.-C. and A.S.A.; project administration, F.C.L.; funding acquisition, F.C.L.

Funding: The research was funded by PDR2020, grant number 101-030671 and Fundação para a Ciência e a Tecnologia, I.P. (FCT) 2022.10859.BD.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: The authors give thanks to Lourenço Palha, Cátia Silva (COTArroz) and Orivárzea (Orizicultores do Ribatejo, S.A.) for technical assistance. We also thank the research centers (GeoBioTec) UIDB/04035/2020, (CEF) UIDB/00239/2020 and Associate Laboratory TERRA (LA/P/0092/2020) for support facilities.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Food and Agriculture Organization of the United Nations (FAO). Coping with water scarcity in agriculture. *FAO*, Rome, Italy. Available online: <https://www.fao.org/3/i6459e/i6459e.pdf> (accessed on 26 July 2023).
2. Fukagawa, N.K. & Ziska, L.H. Rice: Importance for global nutrition. *J. Nutr. Sci. Vitaminol.* **2019**, *65*. <https://doi.org/10.3177/jnsv.65.s2>.
3. Mallareddy, M.; Thirumalaikumar, R.; Balasubramanian, P. et al. Maximizing water use efficiency in rice farming: A comprehensive review of innovative irrigation management technologies. *Water* **2023**, *15*, 1802. <https://doi.org/10.3390/w15101802>.
4. Singh, A.K. & Kumar, S.R. Quality assessment of groundwater for drinking and irrigation use in semi-urban area of Tripura, India. *Ecol. Environ. Conserv.* **2015**, *21*, 97–108. ISSN: 0971-765X.
5. Dantas, S.J. From conventional to genomic: A review on rice breeding methods. *SouzaEAD* **2023**, *61*, 1–17. ISSN: 2595-5934.
6. Das, K.; Roy, P. & Tiwari, R.K.S. Biofortification of rice, an impactful strategy for nutritional security: Current perspectives and future prospect. *Plant-Based Diet* **2023**. <https://doi.org/10.5772/intechopen.110460>.

7. Coelho, A.R.F.; Lidon, F.C.; Pessoa, C.C. et al. Can foliar pulverization with CaCl_2 and $\text{Ca}(\text{NO}_3)_2$ trigger Ca enrichment in *Solanum Tuberosum* L. tubers? *Plants* **2021**, *10*, 245. <https://doi.org/10.3390/plants10020245>.
8. Direção Geral de Agricultura e Desenvolvimento Rural. Carta de Capacidade de Uso do Solo de Portugal—Bases e Normas Adotadas na Sua Elaboração, 6th ed.; Ministério da Economia, Secretaria de Estado da Agricultura, Serviço de Reconhecimento e de Ordenamento Agrário: Lisboa, Portugal, **1972**, 25–26.
9. Rodier, J.; Legube, B.; Merlet, N. L'Analyse de l'Eau, 9th ed.; Dunod: Paris, France, **2009**, 1579. ISBN 9782100072460.
10. Simões, M. Métodos cromatográficos, volumétricos e potenciométricos para análise química quantitativa de água subterrânea e sua aplicação no aquífero cenozóico da Bacia do Baixo Tejo, Portugal. *Geociências Rev. Unesp. São Paulo* **2008**, *27*, 161–169.
11. Piper, A.M. A Graphic Procedure in the Geochemical Interpretation of Water Analyses, Ground Water Note 12; US Geological Survey: Washington, DC, USA, **1953**.
12. Wilcox, L. Classification and Use of Irrigation Waters, Circular, N. 969; USDA: Washington, DC, USA, **1955**.
13. Oliveira, I. & Varela, M. A qualidade da água dos pequenos regadios individuais do Alentejo. In *Proceedings of the I National Congress of Irrigation and Drainage*. Beja, Portugal, **2005**, 5-8.
14. Carrondo, M.; Reboredo, F.; Ganho, R.; Santos Oliveira, J.F. Heavy metal analysis of sediments in Tejo estuary, Portugal, using a rapid flameless atomic absorption procedure. *Talanta* **1984**, *31*, 561–564.
15. Reboredo, F.H.S.; Ribeiro, C.A.G. Vertical distribution of Al, Cu, Fe and Zn in soil salt marshes of the Sado estuary, Portugal. *Int. J. Environ. Stud.* **1984**, *23*, 249–253.
16. Ayers, R. & Westcot, D. Water Quality for Agriculture. Food and Agriculture Organization of the United Nations, Rome, Italy **1985**, *29*, 174.
17. Decreto-Lei nº236/98. Diário da República. I Série A-176, 01/08/1998, 3677-3722.
18. Tanji, K.K. & Kielen, N.C. Agricultural Drainage Water Management in Arid and Semi-Arid Areas. *FAO Irrigation and Drainage* **2002**, *61*, 188.
19. Granjeiro, M.; Santiago, M.; Silva, C. et al. Razões iónicas e índices de Langelier e Larson no estudo hidrogeoquímico das águas subterrâneas no município de Quixeré—Ceará. In *Proceedings of the XVIII Simpósio Brasileiro de Recursos Hídricos*, Campo Grande: Brazil **2009**, 1- 16.
20. Singh, K.K.; Tewari, G.; Kumar, S. Evaluation of groundwater quality for suitability of irrigation purposes: A case study in the Udham Singh Nagar, Uttarakhand. *J Chem* **2020**, 1-15. <https://doi.org/10.1155/2020/6924026>.
21. Ray, M. Influence of Different Weather Parameters on Rice Production-A Review. *ALS* **2016**, *5*, 5776–5782. ISSN: 2278-3849.
22. Wang, Y.D.; Wang, X.; Wong, Y.S. Generation of selenium-enriched rice with enhanced grain yield, selenium content and bioavailability through fertilisation with selenite. *Food Chem.* **2013**, *141*, 2385–2393. <https://doi.org/10.1016/j.foodchem.2013.05.095>.
23. Li, H.F.; McGrath, S.P.; Zhao, F.J. Selenium uptake, translocation and speciation in wheat supplied with selenate or selenite. *New Phytol.* **2008**, *178*, 92–102. <https://doi.org/10.1111/j.1469-8137.2007.02343.x>.