

Potential of *Achyrocline satureioides* ethanolic extract in the control of *Spodoptera littoralis*

Rafaela Regina Fantatto ^{1*} and Rosemeire Cristina Linhari Rodrigues Pietro ^{2*}¹ Department of Drugs and Medicines, São Paulo State University UNESP; rrfbio@hotmail.com² Department of Drugs and Medicines, São Paulo State University UNESP; rosemeire.pietro@unesp.com

* Correspondence: rrfbio@hotmail.com; Tel.: 911962926

Abstract: The species *Spodoptera littoralis* is one of the most destructive agricultural pests in tropical and subtropical area causing significant losses to farmers and the alternative forms of control is necessary. This study aims to evaluate ethanolic extracts from leaves and inflorescences of *Achyrocline satureioides* as a potential control agent for *Spodoptera littoralis*. The ethanolic extract was prepared via maceration and the Bioassay was carried out in 5 replications with sixth instar *S. littoralis* larvae and used the Food Deterrence methodology that remained feeding until the larvae ate 75% of the control or sample. The results demonstrated that the concentration of 10 mg/mL of the extracts from *A. satureioides* leaves and inflorescences inhibited 37.23% and 22.72% of *S. littoralis* feeding, respectively, indicating that the extracts have an inhibitory effect.

Keywords: pests; *A. satureioides*; *S. littoralis*

1. Introduction

The global population on the rise, there is an increasing need to bolster food production. According to a 2009 study conducted by the FAO [1], it is projected that a 70% increase in production will be necessary by the year 2050. This push for expanded production brings with it concerns about conserving natural resources, which in turn drives the exploration of more ecologically sound production methods and the reduction of agrochemical usage. In this context, organic agriculture gains prominence as a promising alternative [2]. In the realm of organic agriculture, alternative methods of pest control have been used. This approach involves utilizing organisms like entomopathogenic fungi, bacteria, nematodes, and plant extracts to combat pests. The use of plant extracts for pest control carries several advantages, including low financial cost, biodegradability, and the absence of the side effects associated with traditional pesticides [3]. An example of a high-impact agricultural pest is the species *Spodoptera littoralis*, commonly known as the cotton leafworm. This pest is particularly destructive in tropical and subtropical regions, feeding on at least 87 types of plants, including vital crops like tomatoes, bell peppers, cotton, and corn [4,5] However, controlling this pest has often been achieved through the use of synthetic chemical insecticides with broad-spectrum activity, resulting in low selectivity. The continuous and exclusive application of these compounds has led to the emergence of resistant populations, compromising control efficacy [6,7]. Given this scenario, the present study focused on evaluating the potential of the extract from *Achyrocline satureioides*, a species belonging to the Asteraceae family found in various regions of South America. This plant is recognized for its acaricidal and nematicidal potential [8]. The study aimed to investigate the effectiveness of this extract against *Spodoptera littoralis* larvae through a choice assay. The evaluation of this extract holds significant relevance in the pursuit of developing more effective and sustainable approaches for agricultural pest control.

2. Methodology

Citation: To be added by editorial staff during production.

Academic Editor: Firstname Last-name

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

2.1. Plant material

Inflorescences and leaves of *Achyrocline satureioides* were obtained through a collaborative agreement between the Faculty of Pharmaceutical Sciences of Araraquara (FCFAR-UNESP) and the Multidisciplinary Center for Chemical, Biological, and Agromomic Research (CPQBA UNICAMP). The extracts were prepared for maceration using methodology development for Fantatto et al., 2022 [9] by allowing them to remain in contact with absolute ethanol at a ratio of 1:10 for 7 days. After this period, the extract was filtered, concentrated through rotary evaporation, and stored in amber vials until use.

2.2. *S. littoralis* larvae obtention

S. littoralis larvae were reared on an artificial diet containing bell pepper plants (*Capsicum annuum*) and maintained at a temperature of 22 ± 1 °C, relative humidity > 70%, and a photoperiod of 16:8 h (light:dark) in a growth chamber.

2.3. Choice Feeding Assay

For the choice feeding assay, the extract was utilized at a concentration of 10 mg/mL, and 5 replicates were conducted for both the extract and control. Petri dishes were coated with agar culture medium at a concentration of 10g/1000 mL of water. After solidification, four holes were created in the agar, and parts of *Capsicum annuum* with approximately 1 cm in diameter were positioned to be affixed but not submerged. Ten microliters of the 10 mg/mL extract were applied to two leaves on each plate, while two leaves remained untreated as controls. Additionally, two *Spodoptera littoralis* larvae were placed on each plate in the same positions. The assay continued until the larvae consumed 75% of either the control or the treatments. Percent feeding reduction (% FR) was determined for the equation $\% FR = [1 - (\text{treatment consumption} / \text{control consumption})] \times 100$ [10,11].

3. Results and discussion

After the larvae had eaten 75% of the control group, they were removed from the plates and leftover leaves were removed and pasted onto a sheet of white paper, separating them into treatment and control (Figure 1). The sheet was scanned and measurements were taken following the methodology of Reina et. al, 2001 [11].




Figure 1. (A) Choice Feeding Assay (B) Leaves considered for the inhibition calculation.

The results revealed that a concentration of 10 mg/mL of extracts from the leaves and inflorescences of *A. satureioides* inhibited 37.23% and 22.72% of *S. littoralis* feeding, respectively. A similar outcome was observed in a study conducted by Hussein et al. in 2023 [12], where the methanolic extract of *S. terebinthifolius* exhibited a feed deterrent activity

of 21.9% at a concentration of 5 mg/mL. *S. terebinthifolius* contains a substantial amount of tannins that deactivate the digestive enzymes in its leaves, forming a tannin-protein complex that is challenging to digest and adversely affects the growth and survival of insects. In the results obtained in the current study using *A. saturoioides* extracts, the observed effect can be attributed to the presence of phenolic compounds, specifically quercetin, isoquercetin, and caffeic acid, which are the predominant molecules in this species [9,13]. Caffeic acid aids plants in defending themselves against predators, pests, and diseases by inhibiting the growth of insects, fungi, and bacteria [14]. Quercetin, on the other hand, is a flavonoid known for its significance in the production of bioinsecticides. Its role in plant metabolism has been associated with a defense mechanism that alters the palatability and nutritional value of the plant, reducing its attractiveness to insects [15,16]. This suggests that the extracts possess an inhibitory effect on the feeding behavior of this caterpillar.

4. Conclusion

The results presented are promising and new trials will be carried out to promote better use of the species *A. saturoioides* in the control of agricultural pests.

References

1. FAO. O estado da comida e agricultura. Organização das Nações Unidas para Alimentação e Agricultura, 2009. Disponível em: . Acessado em 21 de junho de 2021.
2. Babin, D. Protected areas: Combining biodiversity conservation and sustainable development: Foundations and recommendations for a development cooperation strategy on protected area management. Institut Français de la Biodiversité, 2013.
3. Rodrigues, J.S., De Castro, R.M., Da Silva, M.G.G., & Da Silva, D.A. (2017). Atividade inseticida de extratos vegetais e seletividade a insetos benéficos. *Revista Semiárido De Visu*, 5(3), 138-148.
4. Carlos, L.V. Espécies contabilizam 100 das espécies exóticas mais invasivas da Europa. Espécies contabilizam 100 das espécies mais invasivas da Europa. 3: Springer Science; 2008. pág. 269-374.
5. Salama, H.S., Dimetry, N.Z., & Salem, S.A. (1971). On the host preference and biology of the cotton leaf worm *Spodoptera littoralis* Bois. *Zeitschrift für Angewandte Entomologie*, 67(1-4), 261-266.
6. Hawkins, N.J., Bass, C., Dixon, A., & Neve, P. (2019). The evolutionary origins of pesticide resistance. *Biological Reviews*, 94(1), 135-155. <https://doi.org/10.1111/brv.12440>.
7. Kranthi, K.R., Jadhav, D.R., Kranthi, S., Wanjari, R.R., Ali, S.S., & Russell, D.A. (2002). Insecticide resistance in five major insect pests of cotton in India. *Crop protection*, 21(6), 449-460. [https://doi.org/10.1016/S0261-2194\(01\)00131-4](https://doi.org/10.1016/S0261-2194(01)00131-4).
8. Fantatto, R.R., Gainza, Y.A., Figueiredo, A., Sorrechia, R., Chagas, A.C.D.S., & Pietro, R.C.L.R. (2022). The association of extracts of *Achyrocline saturoioides* and the fungus *Beauveria bassiana* against the tick *Rhipicephalus microplus*. *Experimental and Applied Acarology*, 87(4), 351-363. <https://doi.org/10.1007/s10493-022-00736-5>.
9. Fantatto, R.R., de Souza Chagas, A.C., Gainza, Y.A., Politi, F.A.S., de Souza Mesquita, L.M., Vilegas, W., ... & Pietro, R.C.L.R. (2022). Acaricidal and anthelmintic action of ethanolic extract and essential oil of *Achyrocline saturoioides*. *Experimental Parasitology*, 236, 108252. <https://doi.org/10.1016/j.exppara.2022.108252>.
10. Bentley, M.D., Leonard, D.E., Stoddard, W.F., & Zalkow, L.H. (1984). Pyrrolizidine alkaloids as larval feeding deterrents for spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Annals of the Entomological Society of America*, 77(4), 393-397. <https://doi.org/10.1093/aesa/77.4.393>.
11. Reina, M., González-Coloma, A., Gutiérrez, C., Cabrera, R., Rodríguez, M.L., Fajardo, V., & Villarroel, L. (2001). Defensive chemistry of *Senecio miser*. *Journal of Natural Products*, 64(1), 6-11. <https://doi.org/10.1021/np000114o>.
12. Hussein, H.S., Salem, M.Z., Soliman, A.M., & Eldesouky, S.E. (2023). Comparative study of three plant-derived extracts as new management strategies against *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). *Scientific Reports*, 13(1), 3542. <https://doi.org/10.1038/s41598-023-30588-x>.
13. Salgueiro, A.C., Folmer, V., da Rosa, H.S., Costa, M.T., Boligon, A.A., Paula, F.R., ... & Puntel, G.O. (2016). In vitro and in silico antioxidant and toxicological activities of *Achyrocline saturoioides*. *Journal of ethnopharmacology*, 194, 6-14. <https://doi.org/10.1016/j.jep.2016.08.048>.
14. Tošović, J. (2017). Spectroscopic features of caffeic acid: Theoretical study. *Kragujevac Journal of Science*, (39), 99-108.
15. Padial, I.M.P.M., Souza, S.A.D., Malaquias, J.B., Cardoso, C.A.L., Pachú, J.K.D.S., Fioratti, C.A.G., & Mussury, R.M. (2023). Leaf Extracts of *Miconia albicans* (Sw.) Triana (Melastomataceae) Prevent the Feeding and Oviposition of *Plutella xylostella* (Linnaeus, 1758) (Lepidoptera: Plutellidae). *Agronomy*, 13(3), 890. <https://doi.org/10.3390/agronomy13030890>.
16. Sayed, R.A., Mahfouz, H.M., Ali, M.M., & ElBassiony, M.N. (2017). Insecticidal properties of some plant extracts against cotton leafworm *Spodoptera littoralis* (BOISD). *Sinai Journal of Applied Sciences*, 6(2), 111-120. <https://doi.org/10.21608/sinjas.2017.78777>.

Disclaimer/Publisher's Note The authors declare no conflict of interest.