

The Urgency of Implementing Field Research for Fir Forest Conservation and Management: Case Studies in Central Greece

¹Panagiotis P. Koulelis, ¹Panos V. Petrakis

¹Institute of Mediterranean Forest Ecosystems, ELGO DIMITRA, www.fria.gr, Terma Alkmanos 11528 Athens
pkoulelis@elgo.gr, pvpetrakis@elgo.gr

INTRODUCTION & AIM

During the last decades, severe drought events, characterized by high temperatures and low precipitation, caused intense forest dieback episodes across Europe [1, 2]. In addition, Mediterranean regions might become most vulnerable to tree species loss, mainly due to increased frequency and intensity of drought events [3, 4]. The forest ecosystems of Southern Europe, during the last century have been subject to direct abiotic disturbances, e.g., droughts and other climatic factors [e.g. 5, 6]. In Greece, water scarcity is probably the main climatic constraints limiting forest growth. Studies have presented a robust correlation between available water and tree ring width and tree growth in different Mediterranean forests and for different tree species [e.g. 7, 8, 9, 10, 11, 12, 13, 14]. Forest ecosystems are unique in having a detailed description of historical events and disturbances imprinted in the tree-ring record. Fire is the most important disturbance, and it is found that fire scarred trees have this event depicted in their tree-rings. The climate is another factor that leaves an imprint in the tree-ring record. Defoliation events are often found in the studies of time series of tree-ring widths, and they are separated from drought induced events by simultaneously studying non-host tree-ring widths. of major defoliators

The urgency of implementing field research for fir forest conservation and management is evident, especially as Mediterranean forest ecosystems face new challenges due to rising temperatures during the growing season. Ecosystems that fail to adapt or have no time to adapt will likely disappear.

METHOD

In the previously mentioned studies, significant emphasis was placed on methodologies involving the use of drought indices alongside measurements of average tree ring width (Fig. 1), and the development of a common nonlinear function for determining a suitable tree growth model over time, along with correlations between these factors. However, the reduction in radial growth is obscured within the obtained time series, and to uncover the most critical factors, advanced bio-mathematical and/or physico-chemical techniques are also required. These include time series analysis using multivariate singular spectral analysis and examining the amount of carbon-13 (¹³C) in the α -cellulose of tree rings, which can reveal hidden factors causing oscillations and variations in tree ring width patterns. Implementing this approach alongside the previously mentioned methods may lead to more validated results.

RESULTS & DISCUSSION

Species-specific sensitivity to warmer or drier climate could affect tree growth behavior directly. The predicted increase in the severity and frequency of drought events [e.g. 15, 16] globally, might have major impacts on tree growth [e.g. 2]. We revealed a reduced growth index for two different altitude stands regarding fir plots on Mt. Giona after 1999 indicating unfavorable growth conditions in the two last decades. The reduced Average Ring width Index (ARWI) according to the Standard Precipitation Index (SPI), is collateral to those two decades of mild or moderate precipitation conditions, wet or dry. One extreme drought event and one extreme wet event were investigated during a period of 55 years. Those extreme events were translated to negative or positive changes in growth. Other milder events were slightly imprinted on the measurable tree ring growth. The average maximum temperature during the growing season and in specific months (April, July, and August) was linked to tree growth for Mt. Giona stands. Evapotranspiration during this period inversely affected fir growth. One of the stands showed a pronounced decline in tree ring growth (ARWI < 0.6), especially in recent years [12].

A decline in growth was observed by our team, after 1998 for two elevations in Mt. Giona, not connected with SPI but seems to be associated with observed defoliations of the insect *Choristoneura murinana* (Lepidoptera, Tortricidae) or European Fir Budworm (EFB). Finally, our 5 years research revealed and confirmed that EFB causes sporadically severe defoliation to Greek fir at Mt. Giona and Mt. Parnassus (Photo 1) and the relation between climate and tree ring width is disturbed because of the EFB attack at Mt. Giona. Our findings so far, encompassing biometric data, tree ring analysis, observations on plant communities, climatic response, and insect infestations—primarily EFB—across various stands on the mountains, have revealed significant local infestations of varying degrees. In many instances, these infestations were detected in adult trees, particularly in sunny areas or near country roads. Furthermore, our research has revealed the varied ability of the fir trees to adapt to both minor and significant climatic variations.

Monitoring of Greek fir forests in central Greece shows the need for immediate research in the Giona and Parnassus mountains, focusing on radial growth, insect infestations, and climate impacts. Urgent action is needed to address both abiotic and biotic threats. The proposed research aims to monitor, preserve, and protect the fir trees, utilizing new knowledge for informed decision-making in their management. The project's scope includes studying the growth characteristics of the fir forest, mitigating threats from biological factors (primarily EFB) by involving pheromones and other biological methods, promoting natural regeneration, preserving biodiversity, and evaluating the water status of the fir trees in response to climate change. Investigating their interactions and understanding the ecosystem's status concerning the previously mentioned aspects seems to be a significant research priority.




Figure 1. Master Average Tree Ring Width indices for three stands at different elevations [988m (S1), 1.274m (S2), and 1.257m (S3)] in Mt. Giona over time[12]



Photo 1. EFB infestations closely to beekeeper facilities (Photo credit: Panagiotis P. Koulelis)

CONCLUSION

A comprehensive understanding of tree growth across various sites, species, climates, and management practices is vital for foresters and stakeholders. Past evaluations of Greece's National Biodiversity Strategy have shown limited progress in addressing new challenges and threats. The proposed project, specifically focused on the Greek fir, aligns with the National Biodiversity Strategy and supports its goals within the framework of conservation, restoration, and strengthening nationwide enhancement efforts.

REFERENCES

- Peñuelas, J., Lloret, F., Montoya, R.: Severe drought effects on Mediterranean woody flora in Spain. *Forest Science*, 47: 214–218 (2001).
- Dobbertin, M.: Tree growth as indicator of tree vitality and of tree re-action to environmental stress: a review. *European Journal of Forest Research*, 124: 319–333 (2005).
- IPCC, Climate Change 2007: impacts, adaptation and vulnerability. In Parry, M.L., Canzi-ani, O.F., Palutikof, J.P., Van Der Linden, P.J., Hanson, C.E. (eds). Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press. 22 p (2007).
- IPCC: Managing the risks of extreme events and disasters to advance climate change adaptation. In Field, C.B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastran-drea, M.D., Mach, K.J., Plattner, G.K., Allen, S.K., Tignor, M., Midgley, P.M. (eds). A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change. Cambridge, UK, and New York, NY, USA: Cambridge University Press. 582 p (2012).
- McDowell, N., Pockman, W.T., Allen, C.D., Breshears, D.D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D., Yepez, E.A.: Mechanisms of plant survival and mortality during drought: Why do some plants survive while others succumb to drought? *New Phytologist*, 178(4): 719–739. Author, F., Author, S., Author, T.: Book title. 2nd edn. (2008).
- Allen, C.D., Macalady, A.K et al.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 259, 660–684 (2010).
- Sarris, D., Christodoulakis, D., Körner, C.: Recent decline in precipitation and tree growth in the eastern Mediterranean. *Global Change Biology*, 13: 1–14 (2007).
- Papadopoulos, A.: Tree-ring patterns and climate response of Mediterranean fir populations in Central Greece. *Dendrochronologia*, 40: 17–25 (2016).
- Koulelis, P. P., Daskalakou, E. N., & Ioannidis, K. E.: Impact of regional climatic conditions on tree growth on mainland Greece. *Folia Oecologica*, 46(2), 127–136 (2019).
- Koulelis, P.P., Fassouli, V.P., Petrakis, P.V., Ioannidis, K.: The impact of selected climatic factors on the growth of Greek fir on Mount Giona in mainland Greece based on tree ring analysis. *Austrian Journal of Forest Science*, 1, 1–30 (2022).
- Koulelis, P. P., Proutsos, N., Solomou, A. D., Avramidou, E. V., Malliarou, E., Athanasiou, M., ... & Petrakis, P. V. (2023). Effects of climate change on Greek forests: A review. *Atmosphere*, 14(7), 1155. <https://doi.org/10.3390/atmos14071155>
- Koulelis, P. P., & Petrakis, P. V. (2023). Brief Overview of Greek Fir Radial Growth in Response to Climate and European Fir Budworm: Three Case Studies from Giona Mountain, Central Greece. *Climate*, 11(4), 78. <https://doi.org/10.3390/cli11040078>
- Koulelis, P. P., Fassouli, V. P., Solomou, A. D. Petrakis, P. V. (2022). Monitoring Greek fir forests offers useful information regarding radial growth, insect infestations, plant communities and climatic response: Aggregate results. SSRN Electronic Journal (preprint) Available at SSRN: <https://ssrn.com/abstract=4256604> or <http://dx.doi.org/10.2139/ssrn.4256604>
- Petrakis, P. V, Koulelis, P. P., Fassouli, V. P., & Solomou, A. D.: Preliminary results of European budworm *Choristoneura murinana* (Hubner) impact on Greek fir radial growth at Mts Parnassus and Giona. *Folia Oecologica*, 49(2), 102–109 (2022).
- Giorgi, F., Im E-S., Coppola, et al.: Higher Hydroclimatic Intensity with Global Warming. *Journal of Climate*, 24(20):5309–24 (2011).
- Field, C.B., Barros, V., Stocker, T.F., Dahe, Q.: Managing the risks of extreme events and disasters to advance climate change adaptation: Special report of the Intergovernmental Panel on Climate Change. 1st ed. Cambridge: Cambridge Univ. Press (2012).