


# Crystal structure of H<sub>4</sub>L (N-N'-Bis(o- hydroxybenzoyl) 1,4 - diaminobutane

G. González-Riopedre, M.I. Fernández-García, M. Maneiro, A. M  
Gonzalez-Noya, Y. Perez , B. Fernández, M.A. Vázquez-  
Fernández

*Dpto. de Química Inorgánica. Facultad de Ciencias. Univ. de  
Santiago de Compostela. Lugo 27002 (Spain).  
misabel.fernandez.garcia@usc.es*

# ABSTRACT

- A bis-amide bis-phenoxy  $N_2O_2$  ligand was obtained from the 2:1 molar reaction of phenyl salicylate and the diamine, 1,4-Diaminobutane, to yield  $H_4L$ . The ligand has been characterised by elemental analysis, IR, and  $^1H$  and  $^{13}C$  NMR spectroscopies, mass spectrometry (ES) and X ray diffraction spectroscopy




# INTRODUCTION

---

- Our interest in this type of ligands derives from the known ability of ligands containing amide groups to stabilise high oxidation states of metal ions when coordinated with the deprotonated nitrogen atom.
- The oxygen amide atom of these ligands can act as a bridge between neighbouring metallic atoms and afford multi-dimensional and supramolecular structures.
- For this purpose, we selected the tetraanionic  $H_4L$  bisamido-bisphenoxy ligand. We have thought that the utilization of tetraanionic bis-amide bis-phenoxy ligands should favour the aggregation of neighbouring complexes through these donors atoms (amide and phenoxy oxygen).
- In previous papers<sup>1</sup> we reported on the synthesis, structures and properties of Mn(III) complexes with asymmetrical trianionic amido-imino-phenoxy ligands.
- Herein, we present the tetraanionic ligand,  $H_4L$  that contain six potential donor atoms: two amide nitrogen, two phenoxy and two amide oxygen atoms

# SYNTHESIS

- H<sub>4</sub>L were prepared according to the literature<sup>2</sup>, in this case by reaction in a 2:1 molar reaction of phenyl salicylate and the diamine at 180-190 °C for 1 h.
- The product was treated with diethyl ether to obtain a white powder at almost quantitative yield.
- White crystals of H<sub>4</sub>L, suitable for single crystal X-ray diffraction studies, were obtained by slow evaporation of methanol solution of the ligand.




1,2-bis(2-hydroxybenzen-2-carboxamide)-butylendiamine  
**H<sub>4</sub>L**



- The compound was also characterised by elemental analysis, IR, and <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy.

# RESULTS

- Ligand H4L:
- M.p.  $\longrightarrow$  110 °C. R  $\longrightarrow$  55%
- Anal Calc. for C<sub>18</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub>: C, 65.8; H, 6.1; N, 8.5. Found: C, 66.2; H, 5.6; N, 8.6%. MS ES (m/z): 329;
- IR (KBr, cm<sup>-1</sup>):  $\nu$ (N-H) 3407,  $\nu$ (O-H) 3057,  $\nu$ (amide I) 1643,  $\nu$ (amide II) 1586,  $\nu$ (C-O) 1259.



- <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, ppm):  $\delta$  9.17 (s: singlet, 2H), 7.20-7.80 (m: multiplet, 4H), 3.34-3.28 (m, 4H), 1.55 (m, 2H).
- <sup>13</sup>C NMR (DMSO-d<sub>6</sub>, ppm):  $\delta$  133.9 (C-OH), 161.6 (C=O), 27.5 (CH<sub>2</sub>), 39.0 (CH<sub>2</sub>).



# RESULTS

- Crystal data and structure refinement for H<sub>4</sub>L are listed in table 1. Crystal structure, with the numbering scheme, is shown in figure 1. Selected bond lengths and angles, as well as potential hydrogen bonds and bonding scheme, are listed in table 2 and table 3.




Figure 1

# DISCUSSION

- In the crystal structure of the title ligand  $C_{18}H_{20}N_2O_4$ , exists two not-identical parts divided by the C14-C14B.
- The slight asymmetry given is showed by the slight different distances collected in Table 1.


**Table 1:** Selected bond lengths (Å) and angles (°) for  $H_4L$

O(20)-C(2)	1.357(2)	O(20B)-C(2B)	1.354(2)
O(20) - H(20)	0.98(24)	O(20B) - H(20B)	0.96 (2)
O(15)-C(11))	1.2593(19)	O(15B)-C(11B)	1.257(2)
C(1) - C(2)	1.414 (2)	C(1B) - C(2B)	1.406 (2)
C(1) - C(11)	1.482 (2)	C(1B) - C(11B)	1.491 (2)
C(11) - N(12)	1.330(2)	C(11B) - N(12B)	1.333(2)
N(12)- C(13)	1.465(2)	N(12B)- C(13B)	1.459 (2)
N(12)- H(12)	0.851(2)	N(12B)- H(12B)	0.873(14)
C(13) - C(14)	1.519(2)	C(13B) - C(14B)	1.517 (3)
C(2) – O(20)-H(20)	102.5(14)	C(2B) – O(20B)- H(20B)	100.7(16)
O(15)-C(11))C(1)	120.13(15)	O(15B)-C(11B)-C(1B)	120.47(15)
N(12)-C(13)- H(13A)	109.2	N(12B)-C(13B)- H(13C)	109.1
N(12)-C(11)-C(1)	118.75(15)	N(12B)-C(11B)-C(1B)	118.60(16)
C(11)-N(12)-C(13)	124.18(16)	C(11B)-N(12B)- C(13B)	123.57(15)
N(12)-C(13)-C(14-)	112.18(15)	N(12)-C(13)-C(14-)	112.70(15)

# DISCUSSION

- The C(11)-O(15) and C(11B)-O(15B) distances of 1.259 and 1.257 Å respectively are consistent with C=O double bonding of amide groups. The oxygen O(20) and O(20B) atoms are forming phenolic groups, and they present distances O(20)-C(2) and O(20B)-C(2B) of 1.357 and 1.354 Å respectively, corresponding to the expected single bonds. O(20)-H(20) distances 0.98 (4) Å are longer than the O(20B)-H(20B) 0.96 Å.
- This molecule has intramolecular hydrogen bonds between the phenolic oxygens (O(20) and O(20B)) and amide oxygens (O(15), O(15B)). Another hydrogen bond is the interaction between the amide nitrogen and amide oxygen N(12)--H(12)..O(15), and N(12B)--H(12B)..O(15B)\* of a neighboring molecule. Moreover  $\pi$ -stacking interactions are further appreciated between the benzene rings (Table 3, Figure 2). N(12) --H(12) ..O(15)\*\*

**Figure 2.** Stick diagram for H<sub>4</sub>L showing the hydrogen bonding





# REFERENCES

---

- 1.-a) Gustavo González- Riopedre, Manuel R. Bermejo, M. Isabel. Fernández-García, Ana.M. González-Noya, Rosa Pedrido, M Jesús Rodríguez-Doutón, M. Maneiro, *Inorg. Chem*, 2015, **54(6)**, 2512; b) Manuel R. Bermejo, Ana M. González Noya, Victor Abad, M. Isabel Fernández, Marcelino Maneiro, Rosa Pedrido and Miguel Vázquez. *Eur. J. Inorg. Chem*, 2004, **18**, 3696.
- 2.- H. Ojima and K. Aichi, *Nippon Kagaku Kaishi* 1967, **88 (3)** 329-333.