

The role of excipients in neglected tropical diseases

Dolores R. Serrano^{1,±}, Aikaterini Lalatsa^{2,±}, M. Auxiliadora Dea-Ayuela^{3,*}


¹Departamento de Farmacia y Tecnología Farmacéutica, Facultad de Farmacia, Universidad Complutense de Madrid, Plaza Ramon yCajal s/n, Madrid, 28040, Spain.

²School of Pharmacy and Biomedical Sciences, University of Portsmouth, St. Michael's Building, White Swan Road, Portsmouth PO12DT, U.K.

³Departamento de Farmacia, Facultad de Ciencias de la Salud, Universidad Cardenal Herrera-CEU, Moncada, Valencia, 46113, Spain. * mdea@uch.ceu.es

± First co-authorship shared

Keywords: Excipients, leishmaniasis, amphiphile, surfactant, lipids, Eudragit, Labrasol, medium chain fatty acids, lauric acid, sodium deoxycholate

Graphical Abstract	Abstract.
	<p>Abstract. Leishmaniasis is a neglected tropical disease responsible for the ninth largest disease burden in the world. Excipients are necessary for ensuring the stability and bioavailability of currently available antileishmaniasis drugs. In a recent work, we have evaluated the <i>in vitro</i> activity of 30 commercially available excipients against different <i>Leishmania</i> spp., their cytotoxicity and potential use for inclusion in novel formulations.</p>

References

[Curr Top Med Chem.](https://www.ncbi.nlm.nih.gov/pubmed/28730958) 2017 Jul 19. <https://www.ncbi.nlm.nih.gov/pubmed/28730958>