



## Prediction of RIFIN proteins with gene orientation network indices

Viviana F. Quevedo-Tumailli <sup>a, b</sup>, Bernabé Ortega-Tenezaca <sup>a, b, c</sup>, Julio César Vargas-Burgos <sup>b</sup>, Alejandro Pazos-Sierra <sup>a</sup> and Humbert González-Díaz <sup>d, e, \*</sup>


<sup>a</sup> *RNASA-IMEDIR, Computer Science Faculty, University of A Coruña, 15071, A Coruña, Spain.*

<sup>b</sup> *Universidad Estatal Amazónica UEA, Puyo, Pastaza, Ecuador*

<sup>c</sup> *Universidad Regional Autónoma de los Andes UNIANDES-Puyo, Ecuador*

<sup>d</sup> *Dept. of Organic Chemistry II, University of the Basque Country UPV/EHU, 48940, Leioa, Biscay, Spain*

<sup>e</sup> *IKERBASQUE, Basque Foundation for Science, 48011, Bilbao, Biscay, Spain*



**Keywords:** Malaria; *Plasmodium* sp. proteome; Chromosome microstructure; Gene orientation; Complex Networks; Machine Learning

\* Corresponding author: H.G.D. ([humberto.gonzalezdiaz@ehu.es](mailto:humberto.gonzalezdiaz@ehu.es))

### References

1. Junker, B. H.; Koschuetzki, D.; Schreiber, F., Exploration of biological network centralities with CentiBiN. *BMC Bioinformatics* 2006, 7, (1), 219.
2. Haye, A.; Albert, J.; Rooman, M., Modeling the *Drosophila* gene cluster regulation network for muscle development. *PLoS One* 2014, 9, (3), e90285.
3. Inoue, L. Y.; Neira, M.; Nelson, C.; Gleave, M.; Etzioni, R., Cluster-based network model for time-course gene expression data. *Biostatistics* 2007, 8, (3), 507-25.
4. Lavazec, C.; Sanyal, S.; Templeton, T. J., Hypervariability within the Rifin, Stevor and Pfmc-TM superfamilies in *Plasmodium falciparum*. *Nucleic Acids Res* 2006, 34, (22), 6696-707.
5. Hurst, L. D.; Williams, E. J.; Pal, C., Natural selection promotes the conservation of linkage of co-expressed genes. *Trends Genet* 2002, 18, (12), 604-6.
6. Kustatscher, G.; Grabowski, P.; Rappsilber, J., Pervasive coexpression of spatially proximal genes is buffered at the protein level. *Mol Syst Biol* 2017, 13, (8), 937.