

# Green, microwave-assisted synthesis of O-perbutyrylated-alkyl-glycosides<sup>†</sup>

Emmanuel Pérez-Escalante<sup>1</sup>, Luis Guillermo González-Olivares<sup>1</sup>, Araceli Castañeda-Ovando<sup>1</sup>, Verónica Salazar-Pereda<sup>1</sup>, John F. Trant<sup>2</sup>, Mirandeli Bautista-Ávila<sup>3</sup> and Sergio Alatorre-Santamaría<sup>4\*</sup>

<sup>1</sup> Universidad Autónoma del Estado de Hidalgo. Área Académica de Química. Carretera Pachuca-Tulancingo km 4.5, 42184. Mineral de la Reforma, Hidalgo, México.

<sup>2</sup> University of Windsor. Department of Chemistry and Biochemistry. 401 Sunset Ave., N9B 3P4. Windsor, Ontario, Canada

<sup>3</sup> Universidad Autónoma del Estado de Hidalgo. Área Académica de Farmacia. Carretera Pachuca-Actopan s/n, 42160. San Agustín Tlaxiaca, Hidalgo, México.

<sup>4</sup> Universidad Autónoma Metropolitana, Unidad Iztapalapa. División de Ciencias Biológicas y de la Salud. Departamento de Biotecnología. Av. San Rafael Atlixco 186, 09340. Ciudad de México, México.

\* Correspondence: salatorre@xanum.uam.mx

† Presented at the 23rd International Electronic Conference on Synthetic Organic Chemistry, November 15 – December 15, 2019.

E-mail addresses: emmanuel\_perez@uaeh.edu.mx (E. Pérez-Escalante), lgonzales@uaeh.edu.mx (L.G. González-Olivares), ovandoa@uaeh.edu.mx (A. Castañeda-Ovando), salazar@uaeh.edu.mx (V. Salazar-Pereda), j.trant@uwindsor.ca (J.F. Trant), mirandeli@hotmail.com (M. Bautista-Ávila).

Received: date; Accepted: date; Published: date

**Abstract:** Chemical synthesis of carbohydrates is a challenging task. Several protection and deprotection steps of hydroxyl groups are required to ensure regioselective formation of the glycosidic bond. Usually, it is achieved through acylation where conventional heating is combined with addition of Lewis acids as catalysts. This traditional approach has two drawbacks; it is time consuming and often catalysts are hazardous to the environment. An alternative route relies on application of microwaves and/or other Lewis acids with less or no toxicity. Such combination would reduce reaction times and offer a benign synthetic strategy to obtain peracylated compounds.

The current work describes an efficient and environmentally mild synthesis of peracylated glycosides with potential application in enzymatic preparation of carbohydrates. Model compound O-perbutyrylated-phenyl-galactose was synthesized using imidazole as catalyst in the microwave-assisted process. The acylation protocol was optimized, and the target sugar was obtained at 50% yield after 1 h. In conclusion, the combination of imidazole and microwaves provides an excellent alternative to swiftly synthesize peracylated glycosides in a benign way.

**Keywords:** perbutyrylated sugars; imidazole; microwaves; green chemistry.

## 1. Introduction

Traditional carbohydrates synthesis involves multiple protection/deprotection steps, where ester derivatives are the most commonly used protecting groups [1,2]. Carbohydrate esters are obtained via acylation using conventional heating and Lewis acids as catalysts [3]. These protocols often require long reaction times, and the catalysts used may pose health and environmental threats, like in the case of pyridine [4].

Considering the negative environmental impact of the traditional approach, more sustainable protocols for carbohydrate esterification are needed. The benign synthetic route encompasses substitution of hazardous catalyst such as pyridine with imidazole, [5] as well as the use of microwaves under solvent-free conditions [6]. Such strategy follows the basic principles of green chemistry defined as avoidance of the toxic wastes production, and/or preference for chemicals with low toxicity and environmental impact [6].

Although perbutyrylated glycosides have been shown to possess different biopharmaceutical and antitumor properties, they have been hardly investigated as potential peracylated carbohydrate precursors [7,8]. Thus, the present work aimed at developing an efficient and environmentally mild synthesis of peracylated glycosides which can be further used as building blocks in enzymatic preparation of wide variety of sugars.

## 2. Methods

*Chemicals and equipment:* All reagents and solvents were at least ACS grade and were used without further purification. Thin layer chromatography (TLC) was performed with SiO<sub>2</sub> plates Kieselgel, UV<sub>254</sub> and the mobile phase consisted of a mixture of petroleum ether and ethyl acetate (1:0.1), using aqueous sulfuric acid solution (5%) as stain mixture. Nuclear Magnetic Resonance Spectroscopy (NMR) was carried out in CDCl<sub>3</sub> using a Bruker Advance 400 Spectrometer.

*Synthesis of O-perbutyrylated-phenyl-galactose at room temperature:* 100 mg of 1-phenyl-β-D-galactopyranoside (0.39 mmol) and 16 mg of imidazole (0.23 mmol) were dissolved in 4 mL of dried acetonitrile under N<sub>2</sub> atmosphere. Then, 640 μL (3.91 mmol) of butyric anhydride were added and the reaction was maintained for 60 h at room temperature. Product was purified by flash chromatography using a mixture of petroleum ether and ethyl acetate as eluent. Pure compound was characterized by <sup>1</sup>H and <sup>13</sup>C-NMR.


*Synthesis of O-perbutyrylated-phenyl-galactose by microwave heating:* 100 mg of 1-phenyl-β-D-galactopyranoside (0.39 mmol) and 83.7 mg of imidazole (1.2 mmol) were transferred to a G30 vessel (Anton Paar). Afterwards, 2 mL of acetonitrile (HPLC grade) and 2 mL of butyric anhydride (12.22 mmol) were added to the reaction tube, and the system was heated in microwaves at 140 °C during 60 min. Product was recovered with CH<sub>2</sub>Cl<sub>2</sub> and the organic phase was washed with deionized water, 0.1 M acetates buffer (pH=5.5) and 1 M KHCO<sub>3</sub>. Pure product was characterized by <sup>1</sup>H and <sup>13</sup>C-NMR.

*Characterization of O-perbutyrylated-phenyl-galactose:* Product was obtained as pale yellow oil. R<sub>f</sub> (petroleum ether:EtOAc; 1:0.1): 0.55; <sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>): δ 7.28 (m, 2H, H9 and H11), 7.06 (t, J = 7.4 Hz, 1H, H10), 6.99 (dd, J = 8.6 Hz, 0.9 Hz, 2H, H8 and H12), 5.55 – 5.46 (m, 2H, H6), 5.14 (dd, J = 10.5 Hz, 3.4 Hz, 1H, H3), 5.05 (d, J = 8.0 Hz, 1H, H1), 4.23 (dd, J = 11.1 Hz, 7.0 Hz, 1H, H4), 4.14 (dd, J = 11.2 Hz, 6.1 Hz, 1H, H2), 4.10 – 4.05 (m, 1H, H5), 2.46 – 2.36 (m, 2H, H16), 2.27 (dt, J = 7.3 Hz, 4.4 Hz, 4H, H20 and H28), 2.21 (t, J = 7.3 Hz, 2H, H24), 1.69 (dt, J = 9.6 Hz, 4.8 Hz, 2, H27), 1.66 – 1.55 (m, 6H, H15, H19 and H23), 0.99 (t, J = 7.4 Hz, 3H, H26), 0.91 (dt, J = 7.5 Hz, 4.5 Hz, 9H, H14, H18 and H22); <sup>13</sup>C-NMR (101 MHz, CDCl<sub>3</sub>): δ 173.01-172.11 (C13, C17, C21 and C25), 157.13 (C7), 129.65 (C9 and C12), 123.33 (C10), 117.00 (C8 and C12), 99.81 (C1), 71.31 (C3), 70.85 (C5), 68.56 (C2), 66.88 (C4), 61.44 (C6), 36.03 (C16, C20, C24 and C28), 18.78 – 18.17 (C15, C19, C23 and C27), 13.69 (C14, C18, C22 and C26).

### 3. Results and discussion

In order to synthesize *O*-perbutyrylated-phenyl-galactose (Figure 1) in an environmentally benign approach, a methodology described by Tiwari et al. [5] was followed and further optimized to meet green chemistry principles. When the original protocol was followed, in which room temperature was applied, esterification was completed after 60 h. <sup>1</sup>H-NMR spectroscopy showed characteristic signals of the product between 0.91 and 2.46 ppm corresponding to the multiplicity of the methyl and methylene groups of the *O*-butyryl chain. Likewise, <sup>13</sup>C-NMR spectroscopy showed the characteristic shifts for carbons in the *O*-butyryl chain: carbonyl groups were found around 172 ppm, while methyl and methylene group chemical shifts were identified between 13 and 36 ppm, respectively. In these reaction conditions the product was afforded at 33% yield.

With the aim to offer a microwave-assisted method to *O*-perbutyrylated-phenyl-galactose and peracylated sugars in general, the same conditions used previously were tested using microwaves instead of conventional heating. Nonetheless, TLC analysis showed an incomplete consumption of starter material after 1 h of reaction. In fact, the addition of more butyric anhydride and imidazole was needed to complete reaction, extending the process until 2 or 3 h. In order to improve reaction time, the protocol was modified by increasing the concentration of both imidazole (0.3 mmol per hydroxyl group) and butyric anhydride (2 mL; 12.22 mmol). Under these new conditions, a total esterification of *O*-phenyl-galactose was reached in only 1 h and the purified product was obtained at 50% yield.



**Figure 1.** Chemical structure of the target sugar.

### 4. Conclusions

An efficient and environmentally benign methodology was applied to synthesize *O*-perbutyrylated glycosides using imidazole as a catalyst. Optimization of the reaction conditions involved the combination of imidazole and microwaves which afforded the best overall yield in considerably reduced time. Finally, this green synthetic strategy has a great potential to be applied in the enzymatic preparations of complex carbohydrates.

**Acknowledgments:** E. Pérez-Escalante would like to acknowledge Consejo Nacional de Ciencia y Tecnología (CONACYT) for granting the Ph.D. scholarship number 473399.

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

1. Ferrier, R.J. An Historical Overview. In *The organic chemistry of sugars*, 1st ed.; Levy, D.E., Fügedi, P., Eds.; CRC Press: Boca Raton, USA, 2006; pp. 3-24.
2. Bissaro, B.; Monsan, P.; Faure, R.; O'Donohue, M.J. Glycosynthesis in a waterworld: new insight into the molecular basis of transglycosylation in retaining glycoside hydrolases. *Biochem. J.* **2015**, *467*, 17-35.

3. Filice, M.; Guisan, J.M.; Terreni, M.; Palomo, J.M. Regioselective monodeprotection of peracetylated carbohydrates. *Nat. Protoc.* **2012**, *7*, 1783.
4. Qiao, L.; Wang, J.L. Microbial degradation of pyridine by *Paracoccus sp.* isolated from contaminated soil. *J. Hazard. Mater.* **2010**, *176*, 220-225.
5. Tiwari, P.; Kumar, R.; Maulik, P.R.; Misra, A.K. Efficient acetylation of carbohydrates promoted by imidazole. *Eur. J. Org. Chem.* **2005**, *2005*, 4265-4270.
6. Farrán, A.; Cai, C.; Sandoval, M.; Xu, Y.; Liu, J.; Hernáiz, M.J.; Linhardt, R.J. Green solvents in carbohydrate chemistry: From raw materials to fine chemicals. *Chem. Rev.* **2015**, *115*, 6811-6843.
7. Carroux, C. J.; Moeker, J.; Motte, J.; Lopez, M.; Bornaghi, L.F.; Katneni, K.; Ryan, E.; Morizzi, J.; Shackelford, D.M.; Charman, S.A.; Poulsen, S.A. Synthesis of acylated glycoconjugates as templates to investigate in vitro biopharmaceutical properties. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 455-459.
8. Zi, C.T.; Yang, D.; Dong, F.W.; Li, G.T.; Li, Y.; Ding, Z.T.; Zhou, J.; Jiang, Z.H.; Hu, J. M. Synthesis and antitumor activity of novel per-butyrylated glycosides of podophyllotoxin and its derivatives. *Bioorg. Med. Chem.* **2015**, *23*, 1437-1446.



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).