

Protective Antifungal Activity of *Plantago major* Extract against the Phytopathogenic Fungi *Phytophthora cinnamomi*, *Diplodia corticola* and *Colletotrichum* Species [†]

Cláudia Ferreira ¹ and Rui Oliveira ^{1,2,3,*}

¹ Centre of Molecular and Environmental Biology (CBMA), Department of Biology, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal; email1@gmail.com

² Centre for the Research and Technology of Agro-Environmental and Biological Sciences (CITAB), Department of Biology, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

³ Centre of Biological Engineering (CEB), Department of Biology, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

* Correspondence: ruipso@bio.uminho.pt

† Presented at the 1st International Electronic Conference on Food Science and Functional Foods, 10–25 November 2020; Available online: https://foods_2020.sciforum.net/.

Submitted: date; Accepted: date; Published: date

Abstract: Synthetic fungicides for crops protection raise environmental and human concerns due to accumulation in edible vegetables, showing significant toxicity to humans, and in soil, groundwater and rivers, affecting ecological balance. In addition, they are prone to the development of resistant strains because of the single target-based mechanism of action. Plant extracts provide attractive alternatives, as they constitute a rich source of biodegradable secondary metabolites, such as phenols, flavonoids and saponins, which have multiple modes of antifungal action and lower probability of development of resistant fungi. This work has the objective of identifying plant extracts with antifungal activity aiming to contribute to food safety and sustainable agricultural practices. We selected a saponin-containing plant, *Plantago major* and extracted secondary metabolites with 50% (v/v) ethanol, dried by evaporation, and dissolved in water. For antifungal activity, the phytopathogenic fungi *Colletotrichum acutatum*, *Colletotrichum gloeosporioides*, *Colletotrichum godetiae*, *Colletotrichum nymphaeae*, *Diplodia corticola* and *Phytophthora cinnamomi* were selected because they affect fruits and vegetables, such as strawberry, almond, apple, avocado, blueberry and chestnut trees. The aqueous extract was incorporated into PDA medium at different concentrations and mycelial discs were placed in the center of each Petri dish. Growth was measured as radial mycelial growth at 3, 6, and 9 days incubation at 25 °C in the dark. Maximum growth inhibition (32.2%) was obtained against *P. cinnamomi* with 2000 µg/mL extract followed by *C. gloeosporioides* (25.7%) on the sixth day and by *C. godetiae* and *C. nymphaeae* (21.1%) on the ninth day. Results show that *P. major* presents antifungal activity in all phytopathogenic fungi tested and the extract can be used to protect important crops, by inhibiting the development of fungal infections and promoting food security and a sustainable agriculture.

Keywords: antifungal phytochemicals; natural antifungals; sustainable crop production; sustainable agriculture; *Plantago major*

1. Introduction

Crop-protecting synthetic fungicides raise environmental and human concerns due to Results accumulation in edible vegetables [1], showing significant toxicity to humans [2], and in soil [3],

groundwater and rivers [4], affecting ecological balance. *Plantago major* (Figure 1) extract is a rich source of biodegradable secondary metabolites, which have multiple modes of antifungal action and lower probability of development of resistant fungi strains, a very notorious problem with the use of synthetic fungicides [5]. The objective of this work is to evaluate the antifungal activity of *P. major* extract, as a potential replacement of synthetic fungicides, aiming to contribute to sustainable agriculture practices and food safety.



Figure 1. *Plantago major*.

2. Material and Methods

To investigate *P. major* inhibition on the mycelial growth of the phytopathogenic fungi *Colletotrichum acutatum*, *Colletotrichum gloeosporioides*, *Colletotrichum godetiae*, *Colletotrichum nymphaeae*, *Diplodia corticola* and *Phytophthora cinnamomi*, the dried plant was extracted with 50% (v/v) ethanol, the solution dried by evaporation, and the residue dissolved in water. The aqueous extract was incorporated into PDA medium at different concentrations, 100, 500, 1000 and 2000 µg/mL and mycelial discs of each fungus were placed in the center of each Petri dish. Radial mycelial growth was measured at 3, 6 and 9 days after inoculation. For each treatment, three replicates were performed. The assay ended when the negative control reached full growth. The antifungal activity of the extract was calculated in terms of inhibition percentage of mycelia growth by using the following formula:

$$\text{Inhibition (\%)} = \frac{dc - dt}{dc} \times 100,$$

where *dc* is the average increase in mycelia growth in negative control and *dt* is the average increase in mycelia growth in treated sets.

3. Results








Figure 2. Representative images of *P. major* antifungal activity at different concentrations, 100, 500, 1000 or 2000 µg/mL, against *Phytophthora cinnamomi*, on PDA solid medium, after 6 days of incubation (n = 3). Negative control (A), 100 µg/mL (B), 500 µg/mL (C), 1000 µg/mL (D) and 2000 µg/mL (E).






Figure 3. Effect of *P. major* on mycelial growth of *Phytophthora cinnamomi* (A), *Diplodia corticola* (B), *Colletotrichum gloeosporioides* (C), *Colletotrichum acutatum* (D), *Colletotrichum godetiae* (E) and *Colletotrichum nymphaeae* (F) isolates on PDA medium with incorporation of *P. major* extract. Percentage of growth inhibition determined after 3, 6 and 9 days of incubation at different concentrations of *P. major* extract, 100, 500, 1000 or 2000 µg/mL. Data are presented as mean of three independent experiments ± SD. One-way ANOVA and Kruskal Wallis test were used for multiple comparisons. Differences were considered statistically significant if $P < 0.05$. Mean values followed by the same letters are not statistically different (lowercase letters for day 3, capital letters for day 6 and Greek letters for day 9). Comparisons between different days of the same concentration are only represented if they are significant.

4. Conclusions

The extract from *P. major* has the potential to replace synthetic fungicides with convenient application programs in crops in order to control and prevent fungal growth. By inhibiting fungal growth by 20–32.2%, *P. major* extract would not be likely to promote fungal resistances and would not have impact on the environment.

P. cinnamomi

Significant difference between concentrations 100 and 2000 µg/mL on the third day.

D. corticola

Reestablishment of mycelial growth after nine days, suggesting adaptation of the fungus.

C. gloeosporioides

The extract has dose-dependent inhibition on fungal growth on the third and sixth days.

C. acutatum *C. godetiae* *C. nymphaeae*

Significant differences from day 6 to day 9 between 100 and 2000 µg / mL.

References

1. Frank, R.; Braun, H.E.; Ripley, B.D. Residues of insecticides and fungicides on Ontario-grown vegetables, 1986–1988. *Food Addit. Contam.* **1990**, *7*, 545–554, doi:10.1080/02652039009373917.
2. Wang, P.; Yang, M.; Jiang, L.; Wu, Y.J. A fungicide miconazole ameliorates tri-o-cresyl phosphate-induced demyelination through inhibition of ErbB/Akt pathway. *Neuropharmacology* **2019**, *148*, 31–39, doi:10.1016/j.neuropharm.2018.12.015.
3. Sukul, P. Enzymatic activities and microbial biomass in soil as influenced by metalaxyl residues. *Soil Biol. Biochem.* **2006**, *38*, 320–326, doi:10.1016/j.soilbio.2005.05.009.
4. Wightwick, A.; Walters, R.; Allinson, G.; Reichman, S.; Menzies, N. Environmental Risks of Fungicides Used in Horticultural Production Systems. In *Fungicides*; IntechOpen: London, UK, 2010, doi:10.5772/13032.
5. Lucas, J.A.; Hawkins, N.J.; Fraaije, B.A. The Evolution of Fungicide resistance. In *Advances in Applied Microbiology*; Elsevier: Amsterdam, The Netherlands, 2015; Volume 90, ISBN 9780128022757.



© 2020 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).