

# Genetic Analysis of Mutant Strains of *Saccharomyces cerevisiae* with Defects in Mannoprotein Synthesis <sup>†</sup>

Patricia Gil <sup>1</sup>, Alberto Martínez <sup>1</sup>, Elena Palencia <sup>1</sup>, Rocío Velázquez <sup>2</sup>, Manuel Ramírez <sup>1</sup> and Luis M. Hernández <sup>1,\*</sup>

<sup>1</sup> Departamento de Ciencias Biomédicas, Área de Microbiología, Facultad de Ciencias, Edificio Juan Remón Camacho, Avda. de Elvas s/n., 06006 Badajoz, Spain; email1@gmail.com (P.G.); email2@gmail.com (A.M.); email3@gmail.com (E.P.); email4@gmail.com (M.R.)

<sup>2</sup> Dipartimento Scienze Biomediche, Università degli Studi di Padova, 35020 Padua, Italy; email5@gmail.com

\* Correspondence: lmhernan@unex.es; Tel: +34-924-289-777

<sup>†</sup> Presented at the 1st International Electronic Conference on Food Science and Functional Foods, 10–25 November 2020; Available online: [https://foods\\_2020.sciforum.net/](https://foods_2020.sciforum.net/).

Received: 14 May 2020; Accepted: 26 August 2020; Published: date

**Abstract:** The *mn* mutants have constituted a fundamental tool in the study of the structure and biosynthesis of mannoproteins in *Saccharomyces cerevisiae*. They were isolated by the group of Dr. C.E. Ballou by random mutagenesis and a selection method using specific antibodies obtained against the wild-type strain. Initially, the mutants were characterized biochemically and in subsequent years the genes in which they were mutated were identified. All of them encode membrane proteins that catalyze the transfer of mannoses to N-oligosaccharides, sometimes isolated or as part of complexes made up of several proteins. However, the specific mutation of each of these mutants has only been identified in the case of *mn3*. In this work we have completed the characterization of the mutants by sequencing the mutated genes in each of them. As expected, they are point mutations that involve the change of one amino acid for another in the mutated protein, or for a stop signal, resulting in a truncated protein.


**Keywords:** mannoprotein; *Saccharomyces cerevisiae*; *mn* mutants; sequence

---

## 1. Introduction

The glycosylation process in eukaryotic cells has been extensively studied during the last decades and most steps have been almost completely deciphered. In these studies, the yeast *Saccharomyces cerevisiae*, a reference model organism in many laboratories, has played a central role. The process of glycosylation of N-oligosaccharides in *S. cerevisiae* is the same as that of higher eukaryotes in the early stages, which take place in the endoplasmic reticulum (ER) which lead to the synthesis of the so-called “inner core” [1]. However, in the stages that take place in the Golgi apparatus, the so-called “outer chain” is added. These stages are different from higher eukaryotes since the outer chain contains exclusively mannose and lacks other monosaccharides. All N-linked oligosaccharides in *S. cerevisiae* are “high mannose” type. In the study of the glycosylation process in *S. cerevisiae*, the isolation and characterization of defective mutants in the process has been essential. Among them, two groups are worth highlighting: the *alg* mutants (Asparagine-linked glycosylation defective) affected in the stages of the ER [1], and *mn* (mannan defective) affected in the stages that occur in the Golgi apparatus [2]. Figure 1 shows the defects of *mn* mutants in N-oligosaccharides structure. Among the latter, *mn9* presents the most drastic defect since it blocks the addition of the whole outer chain in Golgi [2,3]. The rest of *mn* mutants have also been identified by different

research groups working on glycosylation studies [2,4–9]. However, the genetic defect of *mnn* mutants has only been identified in the case of the *mnn3* mutant [10].



**Figure 1.** Chemotypes of mannoprotein mutants. Adapted from [2].

The objective of this work is to complete the genetic characterization of the *mnn* mutants, identifying the mutations in each of them, by sequencing the mutated genes in the original strains and comparing the sequences with those of the parental strain.

## 2. Materials and Methods

### 2.1. Strains

The mutants *mnn1*, *mnn2*, *mnn5*, *mnn6*, *mnn9*, *mnn10* used in this study are from the laboratory collection and were kindly supplied by Dr. C.E. Ballou some years ago. They were grown in liquid or solid YEPD medium containing 1% yeast extract, 2% peptone, 2% glucose. For solid medium, 2% agar was added.


### 2.2. Extraction of Nucleic Acids

Yeast DNA minipreps were prepared as in [11]. Yeasts cells were resuspended in 1 mL of solution I (EDTA 50 mM pH 7) centrifuged and resuspended again in 1 mL of solution II (Tris-SO<sub>4</sub> 50mM pH 9,3 + 1% mercaptoethanol). After 15 min at room temperature, they were centrifuged again and resuspended in 0.6 mL of solution III (sodium acetate 3 M). Then, 0.5 mL of equilibrate phenol is added and the mixture was incubated at room temperature for 30 min, with shaking. After centrifugation, the nucleic acids, recovered in the aqueous phase, were precipitated with 2 volumes of previously chilled isopropanol, washed with 70% ethanol, dried, and dissolved in Tris-EDTA buffer, pH 8.0.

## 3. Results and Discussion

### Identification of Mutations in MNN Genes Responsible for Synthesis of Outer Chain

Figure 2 shows the N-oligosaccharide biosynthetic pathway in the Gogi apparatus of *S. cerevisiae*. Mnn9p is part of the Mannan polymerases I and II responsible for the elongation of the backbone of mannoses linked in alpha 1–6 of the outer chain. Mnn10p is part of Mannan polymerase II, while Mnn1p, Mnn2p and Mnn5p participate in the synthesis of the side chains or branches of the main chain. Finally, Mnn6p is the transferase that catalyzes the transfer of mannose-P groups.



**Figure 2.** N-glycosylation pathway of mannoproteins in yeast Golgi [7].

**Table 1.** Changes in DNA sequence and effect on encoded protein, in *mnn* mutants as compared to wild type.

GENE	Systematic Name	Location in Chromosome	Mutation in DNA	Effect in Protein	Position in Protein
<i>MNN1</i>	YER001W	153,520..155,808	G > A	C > Y	697
<i>MNN2</i>	YBR015C	267,710..269,503	G > A	G > D	359
<i>MNN5</i>	YJL186W	80,155..81,915	G > A	W > stop signal	561
<i>MNN6</i>	YPL053C	457,118..458,458	Insertion AA	Frameshift	149
<i>MNN9</i>	YPL050C	460,779..461,966	G > A	W > stop signal	16
<i>MNN10</i>	YDR245W	952,800..953,981	G > A	W > stop signal	279

*MNN1*: located at chromosome V (YER001W), 2289 bp. It encodes an integral membrane protein with alpha-1,3-mannosyltransferase activity. The mutation in *mnn1* is a missense point mutation that changes G by A in DNA. It results in a change of cysteine by tyrosine in the protein.

*MNN2*: located at chromosome II (YBR015C), 1794 bp. It encodes an integral membrane protein with alpha-1,2-mannosyltransferase activity over mannoses linked by alpha-1,6. The mutation in *mnn2* is a missense point mutation that changes G by A in DNA. It results in a change of glycine by aspartic acid in the protein.

*MNN5*: located at chromosome V (YJL186W), 1761 bp. It encodes an integral membrane protein with alpha-1,2-mannosyltransferase activity over some mannoses linked by alpha-1,2. The mutation in *mnn5* is a nonsense point mutation that changes G by A in DNA. It results in a change of tryptophan by a stop signal the protein, thus resulting in a truncated protein.

*MNN6*: located at chromosome XVI (YPL053C), 1341 bp. It encodes an integral membrane protein with mannosylphosphate transferase activity. The mutation in *mnn6* is a frameshift mutation due to insertion of AA. The change in the reading frame most probably results in a nonfunctional protein.

*MNN9*: located at chromosome XVI (YPL050C), 1188 bp. It encodes an integral membrane protein which is part of a complex involved in outer chain elongation. The mutation in *mnn9* is a nonsense point mutation that changes (G by A) in DNA. It results in a change of tryptophan by a stop signal the protein, thus resulting in a truncated protein.

*MNN10*: located at chromosome IV (YDR245W), 1182 bp. It encodes an integral membrane protein which is part of a complex involved in outer chain elongation. The mutation in *mnn10* is a nonsense point mutation that changes G by A in DNA. It results in a change of tryptophan by a stop signal the protein, thus resulting in a truncated protein.

As expected, in most cases are point mutations with different effects: missense mutations, nonsense mutations or frameshift mutations.

In the last few years, the *mnn* mutants have been widely characterized phenotypically. With this study, the genotypic characterization is completed.

**Author Contributions:** Conceptualization, funding acquisition, writing and editing, project administration, L.M.H. Methodology and resources, P.G. Investigation: P.G., E.P. Supervision: A.M., R.V., M.R. All authors have read and agreed to the published version of the manuscript.

**Funding:** Projects GR18117 y IB16132 (Consejería de Economía, Ciencia y Agenda Digital, Junta de Extremadura), project AGL2017-87635-R (Ministerio de Economía, Industria y Competitividad, Agencia Estatal de Investigación, Gobierno de España) and Fondo Europeo de Desarrollo Regional (FEDER).

**Acknowledgments:** Patricia Gil is a recipient of a pre-doctoral fellowship from Junta de Extremadura.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Aebi, M. N-linked protein glycosylation in the ER. *Biochim. Biophys. Acta (BBA) Mol. Cell Res.* **2013**, *1833*, 2430–2437.
2. Ballou, C.E. Isolation, characterization and properties of *Saccharomyces cerevisiae mnn* mutants with nonconditional protein glycosylation defects. *Methods Enzymol.* **1990**, *185*, 440–470.
3. Hernandez, L.M.; Ballou, L.; Alvarado, E.; Tsai, P.K.; Ballou, C.E. Structure of the phosphorylated N-linked oligosaccharides from the *mnn9* and *mnn10* mutants of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **1989**, *264*, 13648–13659.
4. Jigami, Y.; Odani, T. Mannosylphosphate transfer to yeast mannan. *Biochim. Biophys. Acta* **1999**, *1426*, 335–345.
5. Rayner, J.C.; Munro, S. Identification of the *MNN2* and *MNN5* mannosyltransferases required for forming and extending the mannose branches of the outer chain mannans of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **1998**, *273*, 26836–26843.
6. Ballou, L.; Alvarado, E.; Tsai, P.K.; Dell, A.; Ballou, C.E. Protein glycosylation defects in the *Saccharomyces cerevisiae mnn7* mutant class. Support for the stop signal proposed for regulation of outer chain elongation. *J. Biol. Chem.* **1989**, *264*, 11857–11864.
7. Jungmann, J.; Rayner, J.C.; Munro, S. The *Saccharomyces cerevisiae* protein Mnn10p/Bed1p is a subunit of a Golgi mannosyltransferase complex. *J. Biol. Chem.* **1999**, *274*, 6579–6585.
8. Lussier, M.; Sdicu, A.-M.; Bussey, H. The *KTR* and *MNN1* mannosyltransferase families of *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* **1999**, *1426*, 323–334.
9. Yip, C.L.; Welch, S.K.; Klebl, F.; Gilbert, T.; Seidel, P.; Grant, F.J.; O'Hara, P.J.; MacKay, V.L. Cloning and analysis of the *Saccharomyces cerevisiae MNN9* and *MNN1* genes required for complex glycosylation of secreted proteins. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 2723–2727.
10. Corbacho, I.; Olivero, I.; Hernandez, L.M. Identification of the *MNN3* gene of *Saccharomyces cerevisiae*. *Glycobiology* **2010**, *20*, 1336–1340.
11. Sambrook, J.; Russell, D.W. Rapid isolation of yeast DNA. *Cold Spring Harbor Protoc.* **2016**, *2006*, prot4039.

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).