

Wind energy assessment during high impact winter storms in the Iberian Peninsula

Ana Gonçalves ^{1,2,*}, Margarida L. R. Liberato ^{2,3} and Raquel Nieto ¹

¹ Environmental Physics Laboratory (EPhysLab), CIM-UVigo, Universidade de Vigo, 32004 Ourense, Spain.

² Instituto Dom Luiz, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal.

³ Universidade de Trás-os-Montes e Alto Douro, UTAD, Quinta de Prados, 5000-801 Vila Real, Portugal.

* Correspondence: ana.redondo.goncalves@uvigo.es

Abstract: Electricity production through renewable energy sources, such as wind energy, is dependent of the variability of weather conditions. Thus, this work aims to assess the wind resource available and the wind energy potential (WEP) during recent December months (the three years 2017, 2018, 2019). These winters were characterized by the occurrence of high impact storms. To understand the effect of the strong winds associated with the passage of the storms during these months, 10m wind components (10-meter U and V wind components) are used. The fields were extracted at 00, 06, 12 and 18 UTC (6-hourly data), for the 2017, 2018 and 2019 December months over a geographical sector that covers the Iberian Peninsula (IP) region (35°N–45°N;10°W–4°E) and compared to climatological values for the 1981-2010 period. The obtained results show an increase of wind intensity of up to 1.2 m.s⁻¹ in IP during December 2017 and 2019 and a decrease of 1.2 m.s⁻¹ in December 2018, when compared with the respective climatology for the 1981-2010 period. This increase corresponds to an increase in the corresponding WEP. Therefore, our results agree with the values of the wind energy produced during the analyzed months for the two countries, Portugal and Spain. Finally, it is notable that the highest values of wind energy production occurred on the days of the storms passage through the IP.

Keywords: Renewable energy; Wind energy; Extreme events; High impact storms; Iberian Peninsula

1. Introduction

The growing and continuing demand for renewable and clean energy, which allow a significant reduction in greenhouse gas emissions, has increased in recent decades. However, it is necessary to understand the availability of the resources and their vulnerability to weather and climate conditions, as well as to climate change [1-2]. Wind energy is one of the fastest growing technologies to produce renewable energy, and in recent years it has become increasingly attractive and presenting a successful economic development. The wind resource has proven to be a renewable, efficient, and clean energy source, with a high potential for reducing greenhouse gas emissions [3].

In the Iberian Peninsula (IP), the two countries, Portugal and Spain, is one of the regions of the world with more installed wind power per capita and where the wind energy plays a key role in the renewable energy mix [4]. In Spain, wind production remains the main renewable energy during the year 2019, with an installed capacity of 25 799 MW and representing 20.8% of total energy production [5]. In Portugal, wind energy has a 5 457 MW of installed capacity [6] which contributes with 27.5% of total electricity generation, corresponding to 12.3 TWh [7]. Therefore, it is necessary to consider the meteorological and climatic variability when designing wind farms.

In this context, weather extreme events like strong extratropical cyclones that originate in the North Atlantic and reach Europe are one of the major natural catastrophes in mid-latitudes and the weather is primarily controlled by the passage of these systems and their associated fronts [8]. When such events, like high impact storms, are extreme and persist across several days cause serious impacts on the sector of production and consumption. These are considered a serious risk for the power production sector of energy and its facilities, and they can cause disruption, supply cuts, and affect other infrastructures depending on the energy supply [9-11]. The rise in electricity production through renewable energy sources, such as onshore and offshore wind energy, also increases the sensitivity of energy systems to weather conditions and their variability. So, it is crucial to consider the spatial and temporal variations in energy production and energy demand in the design and operation of power systems of renewable sources [12], in order to guarantee a continuous and secure energy supply in the future [13].

During the December months of 2017, 2018, and 2019, IP was affected by several high impact storms that caused major adverse impacts [14-16]: in December 2017, three high impact named storms (Ana, Bruno, and Carmen storms [14-16]); in December 2018, other three storms affected IP (Etienne, Flora [14-15]) and Oswald [17] or Deirdre [18]; and in the last considered December (2019) three additional storms (Daniel, Elsa, and Fabien storms [14-15]) affected IP with strong winds associated.

Thus, the objective of this study is to provide a first assessment of the available wind resource and wind energy potential (WEP) during the considered months of December (2017, 2018, and 2019), and to understand the influence of high impact storms in the production of electricity through wind energy technology.

2. Data and Methods

In order to analyse the wind resource availability and the wind energy potential during the three months of December (2017, 2018, 2019), the ERA5 Reanalysis 10m wind components (10-meter U and V wind components) are retrieved from the European Centre for Medium Range Weather Forecasts (ECMWF) [19]. The fields were extracted at 00, 06, 12 and 18 UTC (6-hourly data), for the 2017, 2018 and 2019 December months over a geographical sector that covers the IP region (35°N–45°N;10°W–4°E) and compared to the climatological average of 6-hourly values for the December months of the 1981-2010 period.

The availability of the wind resource is assessed for each of the three months separately. For WEP calculations, ERA5 10 m wind velocities (V_{10}) are extrapolated to 135 m, which is the considered hub height (H) of onshore wind turbines widely installed in the IP and it is used for this study [20]. The wind speed (V) varies with altitude because of frictional effects at the surface of the Earth. Therefore, we must correct for the wind speed at the presumed average hub height of the installed wind turbines [21]. To extrapolate measured wind speed to hub-height, there are several mathematical expressions. Among these are the logarithmic law and the power law [22]. The logarithmic law is used here to characterize the impact of the roughness of the earth's surface on the windspeed and it is expressed by the following equation:

$$\frac{V}{V_{10}} = \frac{\ln(\frac{H}{z})}{\ln(\frac{H_0}{z})} \quad (1),$$

where V is the windspeed at height H , V_{10} is the windspeed at height H_0 (often a reference height of 10 m), and z is the roughness length coefficient [21, 23]. The wind extrapolation is carried out assuming a logarithmic wind profile that assumes a neutrally stratified atmosphere [21]. This equation has been used in many studies of wind energy analyses [4, 24-27] with different applications. In this study, the value of 0.03 is considered for the z parameter.

The power output of a wind turbine (P_{out}), or wind potential (WP), depends on air density (the standard near-surface value of 1.225 kg.m⁻³), turbine radius, R (m), wind speed, V (m.s⁻¹), and rotor power coefficient, C_p , which is the ratio of the rotor power density (mechanical power at the turbine

shaft per unit of swept area) to the wind power density [28]. Between the turbine specific cut-in and rated wind velocity, the power output is proportional to the cubic power of wind speed:

$$P_{out} = \frac{1}{2} C_p \rho \pi R^2 V^3 \quad (2).$$

Hence, sensitivity of P_{out} to wind speed is particularly high and depends on the power coefficients of a given turbine model [29]. The wind energy potential (WEP), or gross energy output (E), for a period (t) is calculated as:

$$WEP = P_{out} \times t \quad (3).$$

For this study, the turbine ENERCON of the model 4MW E-126 EP3 is used and the characteristics are presented in Figure 1 [20, 30].




Figure 1. Power curve (in kW) of the selected wind turbine (4 MW E-126 EP3, ENERCON) for wind speed ranging from 0 to over the cut-out velocity (25 ms⁻¹). Note the rated (maximum) power of 4000 kW, wind rated speed of 13 ms⁻¹ (outlined by the vertical dotted line), rotor diameter of 127 m and cut-in velocity of 3 ms⁻¹. Adapted from: [20, 30].

3. Results and Discussion

The analysis of the wind resource and wind energy potential (WEP) for the considered December months of 2017, 2018 and 2019 is presented in Figure 2 and Figure 3, respectively.

To evaluate the wind resource availability in the December months of 2017, 2018 and 2019 and to understand how the wind resource varies spatially and temporally, the climatological average of the 10 m wind speeds for the month of December for the period 1981 to 2010 was calculated (Fig. 2. A). Then, respective anomalies are calculated for each of the considered years (Fig. 2. B-D). The mean wind speeds over IP range from 1.5 to 4.5 m.s⁻¹. So, the obtained results show an increase of wind intensity of up to 1.2 m.s⁻¹ over the IP in onshore areas, in December 2017 and 2019 (Fig. 2. B and Fig. 2. D), but also in offshore areas (Atlantic Ocean on the north of IP and on Mediterranean Sea) where it reached more than 1.8 m.s⁻¹. In Figure 2. C, for December 2018, it is verified a decrease of about 1.2 m.s⁻¹ over the IP, and even larger on the Mediterranean Sea where the difference reaches -2.4 m.s⁻¹, when compared with the respective climatology for the 1981-2010 period.

Figure 3 presents similar results obtained for the WEP. The 30-year mean WEP was calculated for the December months of the period of 1981 to 2010 (Fig. 3.A) and for each of the December months in analysis (Fig. 3. B-D). In the IP, the mean values of WEP range between 4 to 16 MWh per day and along the coastal areas and on offshore areas these can reach values of 32 MWh per day, as it is presented in Figure 3. A.




Figure 2. (A) Climatological average of ERA5 wind speed at 10m ($\text{m}\cdot\text{s}^{-1}$), for the December months of 1981 to 2010; Anomalies of wind speed at 10m ($\text{m}\cdot\text{s}^{-1}$) for December months of (B) 2017; (C) 2018; (D) 2019.

The obtained results for the anomalies show how the wind resource availability varies; in December 2017, the WEP over the IP had an increase of up to 6 MWh per day (Fig. 3. B). On offshore areas (Atlantic Ocean on the north of IP and on the Mediterranean Sea) this increase reaches 9 MWh per day (Fig. 3. B). In December 2019, there was also an increase of WEP over IP, with the values reaching 6 MWh per day in practically the entire IP. In the offshore areas, there is a decrease in WEP in the eastern region of the Mediterranean Sea and in the entire north and south coast of the IP. It is worth noting the increase in production in the region of the Gibraltar strait, where the values reached up to more than 7 MWh per day. The month of December 2018 showed the lowest values of WEP, with values across the IP lower by up to 6 MWh per day. In the offshore regions, those obtained were found to be lower up to -12 MWh per day, over the Mediterranean Sea and on the southwest coast of Iberia.

These results are in agreement with the obtained values in Santos *et al.* [29] for the mean patterns of annual and winter months (December, January and February) WEP, where the highest values range from 8 to 16 MWh per day on the northwest and north of IP region, on the southeast region of IP and on the Gibraltar strait region. Moreover, while onshore daily WEP varies between 2 and 16 MWh per day, offshore values are never below 16 MWh per day [29].




Figure 3. (A) Climatological average of Wind Energy Potential (WEP) ($\text{MWh}\cdot\text{day}^{-1}$), for the December months of 1981 to 2010; Anomalies of WEP ($\text{MWh}\cdot\text{day}^{-1}$) for December months of (B) 2017; (C) 2018; (D) 2019.

When the electricity production reports are analyzed, for December months in study, the values obtained for mainland Portugal and for Spain (peninsular region) allow to confirm the results obtained in Figures 2 and 3.

In mainland Portugal, for December 2017 there was an increase in energy production compared to previous months. This increase was more accentuated with the passage of the two storms, on 9th to 11th of December, when the wind energy ensured renewable production with values from 2000 to 6000 MW. These high values of wind energy production are related with the passage of Ana storm during these days. On December 26th it was registered the monthly peak of renewable production, 7485 MW, at a time when the country was being affected by the storm Bruno. During the period when the storm Bruno affected mainland Portugal, renewable energies, especially wind energy, represented 105% of the country's electrical needs [31]. In Spain, wind was the technology that most contributed to the production of electricity in December 2017 (24.8%), and on December 27th the daily wind energy production had a share in the generation structure with a maximum of 44.7%, during the passage of Bruno storm [32].

When analyzing December 2018 in mainland Portugal, there is a reduction in monthly wind productivity compared to previous months and compared to the values of the previous year [33]. In Spain, during the month of December of 2018, wind technology was the main source of production, representing 21% of total electricity production, which wind energy reached 4.315 GWh, registering a -25% variation compared to December 2017. The maximum production of wind energy in the peninsular Spain occurred on December 13, 2018 (with the occurrence of storm Flora), in which wind energy represented 35.9% of electricity production on that day. In addition to December 13, on

December 16 (Oswalde storm) there was a maximum of 52% coverage with wind production in the Spanish peninsular region [34].

In December 2019, the Portuguese renewable energy production was 77.2% (4 280 GWh), of which 27.4% corresponded to wind energy and 44.2% to hydroelectric production. These results reflected wind and water productivity far above the average during December, caused by the storms that occurred this month (Daniel, Elsa, and Fabien). This month, a new historical record was reached in renewable production, which corresponded to a consecutive period of 5 and a half days (uninterrupted period of 131 hours), between 18 and 23 December, when the production of renewable electricity was enough to satisfy consumption needs in mainland Portugal. In total, 331 hours of 100% renewable consumption were recorded in December 2019 [7, 35].

In December 2019, the 12th, a new historic maximum of wind power was reached in the peninsular electricity system, with a value of 18 879 MW. On December 13, the maximum daily wind energy was exceeded, reaching 397 541 MWh. The annual participation of wind energy in the peninsular production structure in 2019 was the highest recorded so far, and was the leading technology in the peninsular production structure in the months of January (25.5%), May (23.5%) , November (34.0%) and December (25.4%) [36].

Thus, the reports of energy production confirm the trends of the obtained values with theoretical values of wind speed, and on the other hand they can prove that the high impact storms had influence in the electricity production of the IP. In December 2017 and 2019 it was registered an increase of electricity production by wind energy, and in December 2018 there was a decrease [7, 31-36].

4. Conclusions

Wind energy has an important role in the renewable energy mix of IP [10]. The installed capacity and the annual participation of wind energy in the energetic system of each country is higher and this is important and good point. These countries are betting and continue to bet on wind energy to produce clean energy and reduce CO₂ emissions.

Regarding the results obtained with the theoretical values, these were in line with the real values of electricity production from wind power in both countries, that is, an increase in production in the months of December of the year 2017 and 2019, and a decrease in December 2018. In addition, it was also confirmed that the occurrence of high impact storms with associated strong winds has an influence on the resource available for energy production, since high values were obtained in the production of electricity when the passage of the storms. However, the data we have for this work did not allow us to verify whether the passage of the storms caused the cut-off of the wind turbines, due to the high values of the wind speed and strong gusts with different directions. This fact can be discussed in a future work.

Funding: This work is supported by Fundação para a Ciência e a Tecnologia – FCT through the projects PTDC/CTA-MET/29233/2017 (WEx-Atlantic) and UID/GEO/50019/2020. Partial support was also obtained from the Xunta de Galicia under the Project ED431C 2017/64-GRC Programa de Consolidación e Estructuración de Unidades de Investigación Competitivas (Grupos de Referencia Competitiva) and Consellería de Educación e Ordenación Universitaria, cofunding from the ERDF, in the framework of the Operational Program Galicia 2014–2020.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Ravestein, P., van der Schrier, G., Haarsma, R., Scheele, R., & van den Broek, M. (2018). Vulnerability of European intermittent renewable energy supply to climate change and climate variability. *Renewable and Sustainable Energy Reviews*, 97(August), 497–508. <https://doi.org/10.1016/j.rser.2018.08.057>
2. Davy, R., Gnatiuk, N., Pettersson, L., & Bobylev, L. (2018). Climate change impacts on wind energy potential in the European domain with a focus on the Black Sea. *Renewable and Sustainable Energy Reviews*, 81(June 2017), 1652–1659. <https://doi.org/10.1016/j.rser.2017.05.253>
3. El Khchine, Y., Sriti, M., & El Kadri Elyamani, N. E. (2019). Evaluation of wind energy potential and trends in Morocco. *Heliyon*, 5(6), e01830. <https://doi.org/10.1016/j.heliyon.2019.e01830>
4. Santos, F., Gómez-Gesteira, M., deCastro, M., Añel, J. A., Carvalho, D., Costoya, X., & Dias, J. M. (2018). On the accuracy of CORDEX RCMs to project future winds over the Iberian Peninsula and surrounding ocean. *Applied Energy*, 228(June), 289–300. <https://doi.org/10.1016/j.apenergy.2018.06.086>
5. RED Eléctrica de España. Available online: https://www.ree.es/sites/default/files/11_PUBLICACIONES/Documentos/Renovables-2019.pdf (accessed on 07-09-2020).
6. Direção Geral de Energia e Geologia. Available online: <https://www.dgeg.gov.pt/media/bbbbmb4/dgeg-arr-2020-07.pdf> (accessed on 07-09-2020).
7. Associação de Energias Renováveis – APREN. Available online: <https://www.apren.pt/contents/publicationsreportcarditens/boletim-energias-renovaveis-dezembro-2019-vf.pdf> (accessed on 08-09-2020).
8. Catto, J. L., Ackerley, D., Booth, J. F., Champion, A. J., Colle, B. A., Pfahl, S., Pinto, J.G., Quinting, J.F., Seiler, S (2019), “The Future of Midlatitude Cyclones”, *Current Climate Change Reports*, Vol 5(4), pp. 407–420, <https://doi.org/10.1007/s40641-019-00149-4>
9. Añel, J. A. et al (2017) *Atmosphere*, <https://doi.org/10.3390/atmos8110209>
10. Cronin, J. et al (2018) *Climatic Change*, <https://doi.org/10.1007/s10584-018-2265-4>
11. Tobin, I. et al (2018) *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/aab211>
12. Zscheischler, J. et al. (2018) *Nature Climate Change*, doi:10.1038/s41558-25-018-0156-3
13. van der Wiel, K. et al (2019) *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/ab38d3>
14. Instituto Português do Mar e da Atmosfera – IPMA (2020). Available online: <https://www.ipma.pt/en/index.html>.
15. Agencia Estatal Meteorología-AEMet, (2020), Borrascas con gran impacto: Información divulgativa y de otras temporadas. Available online: <http://www.aemet.es/es/conocerlas/borrascas>
16. Météo France (2020). Available online: <http://www.meteofrance.com/accueil>.
17. Institut für Meteorologie. Met.fu-berlin (2020). Available online: <http://www.met.fu-berlin.de/adopt-a-vortex/historie/>
18. Met Office. Available online: <https://www.metoffice.gov.uk/binaries/content/assets/metofficegovuk/pdf/weather/learn-about/uk-past-events/interesting/2018/storm-deirdre---met-office.pdf> (accessed on 09-09-2020).
19. Hersbach, H., Bell, W., Berrisford, P., Horányi, A., Sabater, J. M., Nicolas, J., Radu, R., Schepers, D., Simmons, A., Soci, C., Dee, D (2019), “Global reanalysis: goodbye ERA-Interim, hello ERA5”, *ECMWF Newsletter*, Vol 159, pp. 17–24, doi: 10.21957/vf291hehd7
20. ENERCON – Energy for the World. Available online: <https://www.all-energy.co.uk/novadocuments/457177?v=636572338170430000>(accessed on 09-09-2020).

21. Hoogwijk, M., de Vries, B., & Turkenburg, W. (2004). Assessment of the global and regional geographical, technical and economic potential of onshore wind energy. *Energy Economics*, 26(5), 889–919. <https://doi.org/10.1016/j.eneco.2004.04.016>
22. Frank, C. W., Pospichal, B., Wahl, S., Keller, J. D., Hense, A., & Crewell, S. (2020). The added value of high resolution regional reanalyses for wind power applications. *Renewable Energy*, 148, 1094–1109. <https://doi.org/10.1016/j.renene.2019.09.138>
23. Masters, G. M. (2004). Renewable and Efficient Electric Power Systems. In *Renewable and Efficient Electric Power Systems*. <https://doi.org/10.1002/0471668826>
24. Costoya, X., deCastro, M., Santos, F., Sousa, M. C., & Gómez-Gesteira, M. (2019). Projections of wind energy resources in the Caribbean for the 21st century. *Energy*, 178, 356–367. <https://doi.org/10.1016/j.energy.2019.04.121>
25. Rogers, T., Ashtine, M., Koon Koon, R., & Atherley-Ikechi, M. (2019). Onshore wind energy potential for Small Island Developing States: Findings and recommendations from Barbados. *Energy for Sustainable Development*, 52, 116–127. <https://doi.org/10.1016/j.esd.2019.08.002>
26. Ryberg, D., Caglayan, D., Schmitt, S., Linßen, J., Stolten, D., & Robinius, M. (2019). The future of European onshore wind energy potential: Detailed distribution and simulation of advanced turbine designs. *Energy*, 182, 1222–1238. <https://doi.org/10.1016/j.energy.2019.06.052>
27. Soares, P. M. M., Lima, D. C. A., Semedo, A., & Cabos, W. (2019). *Climate change impact on Northwestern African offshore wind energy resources*. *Environ. Res. Lett.* 14 (2019) 124065 <https://doi.org/10.1088/1748-9326/ab5731>
28. IEC. Wind turbines: power performance measurements of electricity producing wind turbines, vol. 12-1; 2005.
29. Santos, J. A., Rochinha, C., Liberato, M. L. R., Reyers, M., & Pinto, J. G. (2017). Projected changes in wind energy potentials over Iberia. *Renewable Energy*, 75(2015), 68–80. <https://doi.org/10.1016/j.renene.2014.09.026>
30. ENERCON – Energy for the World. Available online: <https://www.enercon.de/en/products/ep-3/e-126-ep3/>.(accessed on 09-09-2020).
31. Associação de Energias Renováveis – APREN. Available online: <https://www.apren.pt/contents/publicationsreportcarditens/12-boletim-energias-renovaveis--dezembro.pdf> (accessed on 08-09-2020).
32. Rede Eléctrica de España – REE. Available online: https://www.ree.es/sites/default/files/11_PUBLICACIONES/Documentos/InformesSistemaElectrico/2017/informacion_sis_elec_ree_2017.pdf(accessed on 08-09-2020).
33. Associação de Energias Renováveis – APREN. Available online: <https://www.apren.pt/contents/publicationsreportcarditens/12-boletim-energias-renovaveis-2018.pdf> (accessed on 08-09-2020).
34. Rede Eléctrica de España – REE. Available online: https://www.ree.es/sites/default/files/11_PUBLICACIONES/Documentos/ree_diciembre_2018.pdf (accessed on 08-09-2020).
35. Rede Eléctrica Nacional – REN. Available online: https://www.ren.pt/pt-PT/media/comunicados/detalhe/producao_renovavel_abastece_51_do_consumo_em_2019_2 (accessed on 08-09-2020).

36. Rede Eléctrica de España – REE. Available online:
https://www.ree.es/sites/default/files/11_PUBLICACIONES/Documentos/InformesSistemaElectrico/2019/inf_sis_elec_ree_2019.pdf (accessed on 08-09-2020).



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).