

1 *Conference Proceedings Paper*

## 2 **Estimation of crop production and CO<sub>2</sub> fluxes using** 3 **remote sensing: Application to a winter** 4 **wheat/sunflower rotation**

5 Gaétan Pique<sup>1,2,\*</sup>, Taeken Wijmert<sup>1</sup>, Rémy Fieuzal<sup>1</sup> and Eric Ceschia<sup>3</sup>

6 <sup>1</sup> CESBIO, Université de Toulouse, CNES/CNRS/INRAe/IRD/UPS, Toulouse, France; [wijmertt@cesbio.cnes.fr](mailto:wijmertt@cesbio.cnes.fr),  
7 [fieuzalr@cesbio.cnes.fr](mailto:fieuzalr@cesbio.cnes.fr)

8 <sup>2</sup> Agence De l'Environnement et de Maîtrise de l'Energie (ADEME), Angers Cedex 1, France;  
9 [piqueg@cesbio.cnes.fr](mailto:piqueg@cesbio.cnes.fr)

10 <sup>3</sup> INRAE, USC 1439 CESBIO, Toulouse, France; [ceschiae@cesbio.cnes.fr](mailto:ceschiae@cesbio.cnes.fr)

11 \* Correspondence: [piqueg@cesbio.cnes.fr](mailto:piqueg@cesbio.cnes.fr);

12 Received: date; Accepted: date; Published: date

13 **Abstract:** To meet the incoming world's growing food needs and climate change, the agricultural  
14 sector will be forced to adapt their practices. To do so, the contribution of agricultural fields to  
15 greenhouse gas emissions, as well as the impact, on soil, climate and productions, of certain  
16 agricultural practices have to be known. In this study, the SAFY-CO<sub>2</sub> crop model is driven by  
17 remote sensing products in order to estimate CO<sub>2</sub> fluxes on the main crop rotation observed in the  
18 study area, i.e., winter wheat followed by sunflower. Different modeling scenarios are tested,  
19 particularly for intercropping periods, the approach being validated locally thank to eddy  
20 covariance flux measurements and then applied regionally. Results showed that the model was  
21 able to reproduce crop production with high accuracy (rRMSE of 21% and 24% for winter wheat  
22 and sunflower yield, respectively) as well as daily net CO<sub>2</sub> flux (RMSE of 1.29 and 0.97 gC.m<sup>-2</sup>.d<sup>-1</sup>  
23 for winter wheat and sunflower respectively). Moreover, the tested modeling scenarios highlight  
24 the importance of taking the regrowth events into account for assessing accurate carbon budget. In  
25 a perspective of large-scale application, the model was upscaled over more than 100 plots, allowing  
26 to discuss the effect of regrowth on carbon uptake.

27 **Keywords:** crop modeling; remote sensing; CO<sub>2</sub> fluxes; croplands; regrowth  
28

---

### 29 1. Introduction

30 Agriculture is one of the main contributors to global greenhouse gas (GHG) emissions with  
31 almost 12% of the total emissions in 2017 (source: FAO). Because of the heterogeneous character of  
32 the croplands, it is challenging to accurately assess agronomic indicators such as production or CO<sub>2</sub>  
33 fluxes at plot scale over large areas. The general process-based models (Ecosys [1], Isba-Ags [2],  
34 ORCHIDEE [3], etc.) are designed to simulate carbon cycle in different ecosystem but they have  
35 difficulties to represent agricultural ecosystem because of their various climate and soil conditions.  
36 On the other hand, agronomic models (STICS [4], Cropsyst [5], CERES [6], etc.) are suitable to assess  
37 accurate CO<sub>2</sub> fluxes over croplands but they need information on management practices and  
38 cultivars that make them ill-adapted for upscaling. In this context, the simple crop model,  
39 SAFY-CO<sub>2</sub>, was developed and combined with remote sensing products (taking advantage of the  
40 regular observations of vegetation states) to estimate the vegetation development, production and  
41 the CO<sub>2</sub> fluxes over croplands.


42 The long-term objective of this research is to evaluate the impact (on production, carbon and  
43 water fluxes) of certain agricultural practices and rotations at plot scale over wide areas. The most  
44 cultivated crops must therefore be calibrated first in order to simulate crop rotations and different  
45 scenarios during off-season (bare soil, cover crops, mulching, etc.). Winter wheat and sunflower are  
46 the two main crops cultivated in south-west France and have already been validated. The authors of  
47 [7] validated SAFY-CO<sub>2</sub> for winter wheat on biomass, yield and CO<sub>2</sub> fluxes and notably estimated  
48 the daily net CO<sub>2</sub> flux (NEE for net ecosystem exchange) with good accuracy (RMSE = 1.29  
49 gC.m<sup>-2</sup>.d<sup>-1</sup>). More recently [8] validated the model for sunflower and showed that the model  
50 reproduced the NEE with also high accuracy (RMSE = 0.97 gC.m<sup>-2</sup>.d<sup>-1</sup>). Estimating NEE properly is a  
51 prerequisite for assessing carbon budget.

52 The objective of this study is to estimate crop production and more particularly CO<sub>2</sub> fluxes  
53 during a crop rotation, with particular attention to the intercrop period. The proposed approach is  
54 based on the agro-meteorological SAFY-CO<sub>2</sub> model, driven by optical satellite derived products,  
55 considering two modeling scenarios. The different variables needed for the study, as well as the  
56 main steps taken into account in the methodology are described in the section 2. The results are  
57 analyzed and discussed (sections 3 and 4), focusing first on the validation of the estimated fluxes at  
58 the plot scale, and then on estimates performed on a 14 by 13 km<sup>2</sup> area, or more than 100 plots.

## 59 2. Experiments

### 60 2.1. Study area

61 The study area was located in an agricultural region governed by a temperate climate (Figure 1).  
62 The seasonality of weather conditions allowed the cultivation of the main crops encountered in France,  
63 distinguishing “winters crops” (mainly represented by wheat) and “summers crops” (mainly  
64 represented sunflower). The relief was characterized by hilly landscapes that result in heterogeneous  
65 development of crops. Since 2005, continuous measurements of meteorological variables, CO<sub>2</sub> and  
66 water fluxes were performed on plot near Auradé (instrumental device part of ICOS network:  
67 <https://www.icos-cp.eu/>, hereafter called FR-Aur), together with a regular survey of crop biomass and  
68 agricultural practices. In this study, the analysis focused first on winter wheat grown in the 2005-2006  
69 season, followed by sunflower grown in the 2006-2007 season, considering the FR-Aur plot. Then the  
70 same rotation is studied on 111 fields and over different crop years (2013-2014 and 2014-2015).



71

72 **Figure 1.** Location of the study area in France. The altitude (m) is displayed in background


### 73 2.2. Meteorological, fluxes and satellite data

74 The daily meteorological inputs of the model (that is air temperature and global incoming  
75 radiation) were either measured at FR-Aur (for local simulations) or provided by SAFRAN reanalysis

76 [9] for simulation at larger scale. The SAFRAN meteorological data were provided all over France at  
 77 daily time step and at a spatial resolution of 8×8 km<sup>2</sup>.

78 The components needed to obtain CO<sub>2</sub> fluxes were measured using the eddy covariance method,  
 79 turbulent fluxes were then derived from EdiRe software, and post-processed (filtering, quality controls  
 80 and gap filling) in accordance with the CarboEurope-IP recommendations. Finally the gross primary  
 81 productivity (GPP) and ecosystem respiration (R<sub>ECO</sub>) were derived from the partitioning of the NEE  
 82 values of CO<sub>2</sub>. See [7] for more details on the procedure.

83 The timeline of the optical satellite images acquired during the four considered crop years is  
 84 presented in the Figure 2. Regular high spatial resolution images were provided by Formosat-2 (43,  
 85 14 and 17 images for the years 2006, 2007 and 2014 respectively), SPOT-2/4 (4, 7 and 27 images for the  
 86 years 2006, 2007 and 2015) and LANDSAT-8 (16 and 15 images for the year 2015 and 2016). Finally  
 87 the GAI were derived from surface reflectances by mean of the biophysical variables neural network  
 88 tool [10] and averaged at the plot scale.



89  
 90 **Figure 2.** Timeline of satellite images used in this study.

91 **2.2. Methods**


92 The daily time steps SAFY-CO<sub>2</sub> model simulates the temporal evolutions of vegetation variables  
 93 (GAI, biomass and yield) and CO<sub>2</sub> fluxes using climate input variables (air temperature and global  
 94 incoming radiation). The agronomic formalisms has already been presented and detailed in previous  
 95 studies ([7], [11], [12]) so the equations of the model will not be presented here. The parameters of the  
 96 model are either fix, extracted from literature or measurements, or variable and constrained by  
 97 boundaries. They are crop specific and fully detailed in [7] and [8] for winter wheat and sunflower,  
 98 respectively. On each simulated field and each year independently, the values of the 8 calibrated  
 99 parameters are determined by minimizing the quadratic difference between the simulated and satellite  
 100 derived GAI (process detailed in [7]), through a constrained version of the simplex method [13]. This  
 101 step allows the model to reproduce all types of developments observed (by satellites) on the  
 102 considered fields.

103 In the present study, the model is validated at a local scale over a winter wheat/sunflower  
 104 rotation covering two crop years (2005-2006 and 2006-2007) using CO<sub>2</sub> fluxes measurements. Then  
 105 the same rotation is simulated at a larger scale on 111 fields and over two different crop years  
 106 (2013-2014 and 2014-2015). In the two modeling exercises (i.e., local and regional scale), two  
 107 scenarios were considered, i.e., with or without simulation of regrowth events.

108 **3. Results**

109 **3.1. Local validation at FR-Aur**

110 Figure 3 presents the temporal evolutions of the net CO<sub>2</sub> flux (NEE) and its components (the  
 111 GPP and the R<sub>ECO</sub>) and Table 1 summarizes the performances of the model in estimating these three  
 112 variables for the different periods of simulation (characterized by different colors on Figure 3). Since  
 113 there is no GPP during bare soil period, GPP statistics are calculated over vegetation period (from  
 114 sowing to harvest and during off-season when regrowth are simulated).



115

116

117

**Figure 3.** Temporal evolutions of the GPP, the  $R_{ECO}$  and the NEE. Winter wheat, bare soil or regrowth and sunflower periods are displayed in yellow, brown, dashed brown and green respectively.

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

The model was able to accurately reproduce the three temporal dynamics. Indeed, over the entire simulation period (i.e., two years) the model showed very good correlations with observations ( $R^2$  of 0.93, 0.83 and 0.86 for GPP,  $R_{ECO}$  and NEE, respectively) and low errors (RMSE of 1.49, 0.70 and 1.06  $\text{gC.m}^{-2}.\text{d}^{-1}$  for GPP,  $R_{ECO}$  and NEE, respectively). Regarding the off-season period (delimited by vertical dashed lines on Figure 3), no correlations were found for the three simulated variables. This period was characterized by very heterogeneous weeds development on the field. Since the model is calibrated thanks to remote sensed GAI averaged over the entire plot, this heterogeneity is ‘smoothed’ in the optimization process and thus in the model outputs. Conversely,  $\text{CO}_2$  flux measurements are representative of a specific area, inside the plot, which change according to the wind. In these conditions, it would be a hard task to represent accurately the dynamic of the  $\text{CO}_2$  fluxes. Nevertheless, taking regrowth events into account allows to significantly improve the  $\text{CO}_2$  flux estimates. Indeed, over this period (corresponding to 102 days), the difference between simulated and measured NEE is 87% ( $104.1 \text{ gC.m}^{-2}$ ) while it is reduced to -27% ( $-31.4 \text{ gC.m}^{-2}$ ) when considering this regrowth events.

**Table 1.** Summary of model’s performances in estimating GPP,  $R_{ECO}$  and NEE for different time periods corresponding to different surface occupations.


		$R^2$	RMSE [ $\text{gC.m}^{-2}.\text{d}^{-1}$ ]	Mean bias [ $\text{gC.m}^{-2}.\text{d}^{-1}$ ]
<b>GPP</b>	2-year period	0,93	1,49	0,28
	Winter wheat season	0,94	1,48	0,38
	Regrowth period	0,03	1,46	1,15
	Sunflower season	0,92	1,50	0,09
<b><math>R_{ECO}</math></b>	2-year period	0,83	0,70	0,00
	Winter wheat season	0,88	0,66	0,07
	Bare soil period	0,05	0,93	-0,08
	Regrowth period	0,01	1,30	0,75
<b>NEE</b>	2-year period	0,86	1,06	-0,06
	Winter wheat season	0,89	1,10	0,12
	Bare soil period	0,10	1,58	-1,02
	Regrowth period	0,02	1,11	0,31
	Sunflower season	0,86	0,80	0,08

134

135

### 3.2. Model’s upscaling

136 The values of net ecosystem productivity (NEP, equal to the NEE integrated over a time period)  
137 estimated over 111 fields without consideration of regrowth event are presented in the Figure 4-A.  
138 The NEP obtained on the most observed crop rotation within the study area varies between -186.4  
139 and 298.1  $\text{gC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ . The majority of plots are therefore considered to be carbon sinks. Nevertheless,  
140 23% of the plots cultivated with these two crops behave as sources. The average NEP value  
141 considering this scenario is  $-44.1 \text{ gC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ , while that taking regrowth into account is close to  $-59.0$   
142  $\text{gC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ . This slight difference between the two scenarios can be explained by the low number of  
143 plots with regrowth events. Indeed, among the considered plots, 24 presented regrowth events  
144 (identifiable through remote sensed GAI dynamics). The Figure 4-B presents the difference of NEP  
145 between simulations without and with taking regrowth into account.



146

147 **Figure 4.** Spatial distribution of the net ecosystem productivity (NEP) simulated over 111 fields  
148 without taking regrowth events into account (A), and differences between the scenario where  
149 regrowth events are considered (B).

150 Taking regrowth into account increases the carbon sink of the considered plot from -28.0 to  
151 -139.5  $\text{gC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$ . Considering only plots where regrowth were simulated, the average NEP varies  
152 from  $-16.1 \text{ gC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$  (bare soil simulated) to  $-85.2 \text{ gC}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$  (regrowth simulated). Furthermore,  
153 among the 24 plots concerned by regrowth events, 12 behaved as a source of carbon without  
154 considering regrowth while only 4 remained a source after regrowth simulation. Indeed, because  
155 carbon assimilation period is longer when vegetation developed on a field during off-season, the  
156 NEP is lower (more negative) this means that it increases the plot carbon sink.

#### 157 4. Discussion

158 In this study the SAFY-CO<sub>2</sub> model has been adapted to simulate crop rotations. So far, only  
159 winter wheat and sunflower crops are calibrated so only rotations between these 2 crops can be  
160 simulated. A generic parametrization has also been defined for regrowth events allowing to improve  
161 NEE and thus NEP estimated which is crucial when trying to assess carbon budgets.

162 To the best of our knowledge, no crop model considers regrowth events to assess NEP and thus  
163 net ecosystem carbon budget (NECB). We demonstrated here that these events could have important  
164 impact on CO<sub>2</sub> fluxes that needs to be considered when simulating crop rotations. Indeed, the  
165 development of cover crops at large scale could have a strong mitigation impact via atmospheric  
166 carbon storage in soils and could be quantify with a tool such as SAFY-CO<sub>2</sub>.

167 So far, we are not able to identify the nature of regrowth (i.e., weeds, cover crop or spontaneous  
168 regrowth) so the same parametrization was used to simulate all regrowth events. In the near future  
169 and in order to improve regrowth simulations, the parametrization of the regrowth will have to be  
170 refined according to their nature that could be retrieved by the use of radar products. Indeed, the  
171 radar could give information on the nature of the regrowth through the geometry of the cover.

#### 172 5. Conclusions

173 In the proposed study, the SAFY-CO<sub>2</sub> model was applied to a winter wheat/sunflower rotation,  
174 offering satisfactory performances concerning the estimation of net CO<sub>2</sub> fluxes and its components.



175 Over the two simulated crop years at FR-Aur, the model estimated the net CO<sub>2</sub> flux with high  
176 correlation ( $R^2 = 0.86$ ) and low error (RMSE = 1.06 gC.m<sup>-2</sup>.d<sup>-1</sup>). The modeling scenarios highlighted  
177 the importance of taking the regrowth events into account for assessing accurate carbon budget. On  
178 the plot equipped with a flux tower, the estimates taking regrowth (weeds in this case) into account  
179 allowed to reduce the error on the NEP from 87% to -27%. On a larger scale, regrowth events  
180 increase the carbon sequestration capacity observed during a 2-year crop rotation, with values  
181 ranging from -28.0 to -139.5 gC.m<sup>-2</sup>.yr<sup>-1</sup>.

182 The approach proposed in this study constitutes a diagnostic tool, particularly promising in a  
183 context where intercrop periods tend to be vegetalized. With a view to carrying out assessments  
184 integrating a greater diversity of crops, future studies should focus on the parameterization of  
185 maize, rapeseed or soybean, as well as on the characterization of intermediate crops.

186 **Author Contributions:** “Conceptualization, G.P., T.W., R.F. and E.C.; methodology, G.P., T.W., R.F. and E.C.;  
187 software, G.P., T.W., R.F. and E.C.; validation, G.P., T.W., R.F. and E.C.; formal analysis, G.P., T.W., R.F. and  
188 E.C.; investigation, G.P., T.W., R.F. and E.C.; resources, G.P., T.W., R.F. and E.C.; data curation, G.P., T.W., R.F.  
189 and E.C.; writing—original draft preparation, G.P., T.W., R.F. and E.C.; writing—review and editing, G.P., T.W.,  
190 R.F. and E.C.; visualization, G.P., T.W., R.F. and E.C.; supervision, R.F. and E.C.; project administration, E.C.;  
191 funding acquisition, E.C. All authors have read and agreed to the published version of the manuscript.”

192 **Funding:** This research was funded by the Climate-KIC 2020 soil carbon farming project (75000142), the Agence  
193 De l’Environnement et de la Maîtrise de l’Energie (ADEME), which financed the project “Couverts  
194 Intermédiaires pour l’atténuation du Changement Climatique” (CICC) and half of Gaétan’s dissertation  
195 research, the Centre National d’Etudes Spatiales (CNES), which financed half of Gaétan’s dissertation research,  
196 and the Agence de l’Eau Adour Garonne (AEAG), which financed the Bag’ages project.

197 **Acknowledgments:** This work was made possible through the support of the Agence De l’Environnement et de  
198 la Maîtrise de l’Energie (ADEME), which financed the project “Couverts Intermédiaires pour l’atténuation du  
199 Changement Climatique” (CICC) and half of Gaétan’s dissertation research, the Centre National d’Etudes  
200 Spatiales (CNES), which financed half of Gaétan’s dissertation research, and the Agence de l’Eau Adour  
201 Garonne (AEAG), which financed the Bag’ages project. The data acquisition at FR-Aur was funded mainly by  
202 the Institut National des Sciences de l’Univers of the Centre National de la Recherche Scientifique (CNRS-INSU)  
203 through the ICOS and OSR SW observatories. Facility and staff were also funded and supported by the  
204 University Toulouse III—Paul Sabatier, the CNES and IRD (Institut de Recherche pour le Développement).

205 **Conflicts of Interest:** The authors declare no conflict of interest.

## 206 References

- 207 1. R. F. Grant *et al.*, ‘Net Biome Productivity of Irrigated and Rainfed Maize–Soybean Rotations: Modeling vs.  
208 Measurements’, *Agronomy Journal*, vol. 99, no. 6, Art. no. 6, 2007, doi: 10.2134/agronj2006.0308.
- 209 2. J.-C. Calvet *et al.*, ‘An interactive vegetation SVAT model tested against data from six contrasting sites’,  
210 *Agricultural and Forest Meteorology*, vol. 92, no. 2, Art. no. 2, Jul. 1998, doi: 10.1016/S0168-1923(98)00091-4.
- 211 3. G. Krinner *et al.*, ‘A dynamic global vegetation model for studies of the coupled atmosphere-biosphere  
212 system’, *Global Biogeochemical Cycles*, vol. 19, no. 1, Art. no. 1, 2005, doi: 10.1029/2003GB002199.
- 213 4. N. Brisson *et al.*, ‘An overview of the crop model stics’, *European Journal of Agronomy*, vol. 18, no. 3–4, Art.  
214 no. 3–4, Jan. 2003, doi: 10.1016/S1161-0301(02)00110-7.
- 215 5. C. O. Stöckle, M. Donatelli, and R. Nelson, ‘CropSyst, a cropping systems simulation model’, *European*  
216 *Journal of Agronomy*, vol. 18, no. 3–4, Art. no. 3–4, Jan. 2003, doi: 10.1016/S1161-0301(02)00109-0.
- 217 6. C. A. Jones, J. R. Kiniry, and P. T. Dyke, *CERES-Maize: a simulation model of maize growth and development*.  
218 College Station: Texas A&M University Press, 1986.
- 219 7. G. Pique *et al.*, ‘Estimation of daily CO<sub>2</sub> fluxes and of the components of the carbon budget for winter  
220 wheat by the assimilation of Sentinel 2-like remote sensing data into a crop model’, *Geoderma*, vol. 376, p.  
221 114428, Oct. 2020, doi: 10.1016/j.geoderma.2020.114428.
- 222 8. G. Pique, R. Fieuzal, P. Debaeke, A. Al Bitar, T. Tallec, and E. Ceschia, ‘Combining High-Resolution  
223 Remote Sensing Products with a Crop Model to Estimate Carbon and Water Budget Components:  
224 Application to Sunflower’, *Remote Sensing*, vol. 12, no. 18, p. 2967, Sep. 2020, doi: 10.3390/rs12182967.

- 225 9. Y. Durand, E. Brun, L. Merindol, G. Guyomarc'h, B. Lesaffre, and E. Martin, 'A meteorological estimation  
226 of relevant parameters for snow models', *Annals of Glaciology*, vol. 18, pp. 65–71, 1993, doi:  
227 10.1017/S0260305500011277.
- 228 10. F. Baret *et al.*, 'LAI, fAPAR and fCover CYCLOPES global products derived from VEGETATION', *Remote  
229 Sensing of Environment*, vol. 110, no. 3, Art. no. 3, Oct. 2007, doi: 10.1016/j.rse.2007.02.018.
- 230 11. B. Duchemin, P. Maisongrande, G. Boulet, and I. Benhadj, 'A simple algorithm for yield estimates:  
231 Evaluation for semi-arid irrigated winter wheat monitored with green leaf area index', *Environmental  
232 Modelling & Software*, vol. 23, no. 7, Art. no. 7, Jul. 2008, doi: 10.1016/j.envsoft.2007.10.003.
- 233 12. B. Duchemin *et al.*, 'Impact of Sowing Date on Yield and Water Use Efficiency of Wheat Analyzed through  
234 Spatial Modeling and FORMOSAT-2 Images', *Remote Sensing*, vol. 7, no. 5, Art. no. 5, May 2015, doi:  
235 10.3390/rs70505951.
- 236 13. J. C. Lagarias, J. A. Reeds, M. H. Wright, and P. E. Wright, 'Convergence Properties of the Nelder--Mead  
237 Simplex Method in Low Dimensions', *SIAM Journal on Optimization*, vol. 9, no. 1, Art. no. 1, Jan. 1998, doi:  
238 10.1137/S1052623496303470.  
239



© 2020 by the authors. Submitted for possible open access publication under the terms  
and conditions of the Creative Commons Attribution (CC BY) license  
(<http://creativecommons.org/licenses/by/4.0/>).