

Crown Ether Benzoxazolyl-Alanines as Fluorimetric Chemosensors for the Detection of Palladium in Aqueous Environment †

Cátia D. F. Martins, Patrícia M. R. Batista, M. Manuela M. Raposo and Susana P. G. Costa *

Centre of Chemistry, University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal;
catiadf_martins@hotmail.com (C.D.F.M.); patricia_rbatista@hotmail.com (P.M.R.B.); mfox@quimica.uminho.pt (M.M.M.R.)

* Correspondence: spc@quimica.uminho.pt

† Presented at the 24th International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2020; Available online: <https://ecsoc-24.sciforum.net/>.

Received: date; Accepted: date; Published: date

Abstract: Palladium has wide application in different contexts and, as a consequence, high levels of palladium in the environment have been reported, representing a risk to human health. Considering the interest to develop more selective and sensitive chemosensors for this analyte, two novel benzoxazolyl-alanine derivatives bearing a crown ether moiety were studied as potential fluorimetric chemosensors for palladium detection. Preliminary chemosensory studies for these unnatural amino acids in the presence of selected metal cations were performed in acetonitrile solution and in aqueous mixtures of sodium dodecyl sulfate (SDS, 20 mM, pH 7.5) solution with acetonitrile, 90:10 v/v. In acetonitrile solution, these probes had a fluorescence response for different cations but, most importantly, in SDS aqueous solution both compounds displayed a selective fluorescence response in the presence of palladium.

Keywords: fluorimetric chemosensors; metallic cations; palladium; unnatural amino acids

1. Introduction

Palladium (Pd) is a transition metal extensively applied in several fields due to its special chemical and physical properties. For example, it is used in dental restorations, chemical catalysts, jewelry, electric equipment, automobile industry, among others [1,2]. As a consequence of its wide application, high levels of palladium in the environment have been reported, representing a risk to human health [3,4]. So far, a variety of small fluorescence probes have been successfully developed for Pd²⁺ detection [5], but there is an interest to design improved water-soluble probes for recognition of this metal in biological and environmental systems [1,6].

For the sensing of metallic cations, there are reports on fluorescent sensors based on amino acids containing different heterocyclic fluorescent and/or coordination units at the side chain [7–13]. Metal cations are known to be complexed through electron donor atoms at the main/side chains in amino acids, and the insertion of heterocycles at the side chain of natural amino acids yields novel unnatural amino acids with added functionality. In particular, the inclusion of crown ethers is largely used in the design of new chemosensors due to their unique ability to coordinate the cations of alkaline metals and they are also effective complexing reagents for alkaline-earth and transition metal ions [14,15].

Bearing these facts in mind, and considering the research group's experience on the design, synthesis and characterization of fluorescent chemosensors [7–12], we report herein the evaluation of two benzoxazolyl-alanine derivatives bearing a crown ether moiety as potential fluorimetric

chemosensors for Pd²⁺ detection in aqueous media. Preliminary chemosensory studies for these unnatural amino acids in the presence of selected metal cations, with biological and environmental relevance, were performed in acetonitrile solution and in aqueous mixtures of SDS (20 mM, pH 7.5) solution with acetonitrile, 90:10 v/v.

2. Experimental Section

Methods and Materials

The synthesis and structural characterization of benzoxazolyl-alanine derivatives **1a-b** has been reported elsewhere [16]. For the photophysical characterization, UV-vis absorption spectra were obtained in acetonitrile solution (1×10^{-5} M) using a Shimadzu UV/2501PC spectrophotometer (Shimadzu Europa GmbH, Duisburg, Germany) and the fluorescence spectra were obtained using a Horiba FluoroMax-4 spectrofluorometer (HORIBA Europe GmbH, Darmstadt, Germany), using 9,10-diphenylanthracene in ethanol as fluorescence standard.

Evaluation of benzoxazolyl-alanine derivatives **1a-b** as fluorimetric chemosensors was carried out in the presence of several cations (Ag⁺, K⁺, Li⁺, Hg²⁺, Ca²⁺, Co²⁺, Pb²⁺, Mn²⁺, Fe²⁺, Zn²⁺, Ni²⁺, Cd²⁺, Cu²⁺, Pd²⁺, Fe³⁺ and Al³⁺). Solutions of compounds **1a-b** (3×10^{-5} M) and of the ions under study (1×10^{-2} M) were prepared in acetonitrile and acetonitrile/water (75:25). Solutions of the compounds **1a-b** (1×10^{-5} M) and of the ions under study (1×10^{-2} M) were prepared in aqueous mixtures of SDS (20 mM, pH 7.5) solution with acetonitrile, 90:10 v/v. Preliminary studies were carried out by addition of up to 10 equivalents of each cation to the solution of compounds **1a-b** in acetonitrile and in mixture of acetonitrile/water. A similar study was performed by addition of up to 10 and 20 equivalents of each cation to the solution of compounds **1a-b** in aqueous environments using SDS. The solutions were analysed in a CN15 viewing cabinet under UV lamp at 365 nm (Vilber Lourmat, Marne-la-Vallée, France).

3. Results and Discussion

Two benzoxazolyl-alanine derivatives bearing a crown ether moiety **1a-b** (Figure 1), previously synthesized, were characterised by UV-vis absorption and fluorescence spectroscopy. In both compounds, there is a protected benzoxazolyl-alanine core which is substituted at position 2 of the oxazole ring with a phenyl linked to a 15C5 azacrown ether moiety (**1a**) or a thiophene coupled to a 18C6 benzocrown ether (**1b**).




Figure 1. Crown ether benzoxazolyl-alanine derivatives (**1a-b**).

Solutions of crown ether benzoxazolyl-alanines **1a-b**, in acetonitrile (1×10^{-5} M), were analyzed and the wavelengths of maximum absorption and fluorescence, λ_{abs} and λ_{flu} , molar absorptivities at the absorption maximum, ϵ , relative fluorescence quantum yields, Φ_F , and Stokes' shifts were compiled in Table 1. Both compounds are highly fluorescent and the higher conjugation in

compound **1b** is in agreement with the observed bathochromic shift in absorption and fluorescence, when compared to compound **1a**.

Table 1. UV-vis absorption and fluorescence data of crown ether benzoxazolyl-alanines **1a-b**, in acetonitrile (1×10^{-5} M).

Compound	UV-vis Absorption		Fluorescence		
	λ_{abs} (nm)	$\log \epsilon$	λ_{flu} (nm)	Φ_F	Stokes' Shift
1a	334	4.04	395	0.82	61
1b	363	4.01	445	0.69	82

The novel benzoxazolyl-alanines **1a-b** were evaluated as fluorimetric chemosensors for the detection of metal cations, with biological and environmental relevance, through preliminary chemosensory studies.

Firstly, the fluorimetric behavior of compounds **1a-b** in the presence of selected cations was studied in acetonitrile, by addition of 10 equivalents of each cation. As expected, these probes had a different fluorimetric response for different cations: compound **1a** exhibited a remarkable fluorescence quenching upon interaction with Hg^{2+} , Pb^{2+} , Fe^{2+} and Pd^{2+} (Figure 2a), whereas compound **1b** interacted with Hg^{2+} and Pd^{2+} through a decrease of fluorescence and a complete quenching was seen in the presence of Fe^{3+} (Figure 3a). Considering the importance of water-soluble probes for recognition of metals in biological and environmental systems, the fluorimetric response of compounds **1a-b** to selected cations was tested in mixtures of acetonitrile/water (75:25). However, a relevant response was not observed in these conditions.




Figure 2. Preliminary chemosensing study of benzoxazolyl-alanine derivative **1a**: (a) in the presence of 10 equivalents of each cation, in acetonitrile (3×10^{-5} M); (b) in the presence of 10 equivalents of each cation, in SDS (20 mM, pH 7.5)-acetonitrile 90:10 (v/v) solution (1×10^{-5} M); (c) in the presence of 20 equivalents of each cation, in SDS (20 mM, pH 7.5)-acetonitrile 90:10 (v/v) solution (1×10^{-5} M).

Based on our previous experience, the use of an anionic surfactant such as sodium dodecylsulphate (SDS) was attempted to overcome this problem [17]. In fact, several authors reported that in aqueous environments using surfactants, selected binding sites and fluorophores can be arranged in micelles of surfactants allowing detection of metal cations in water by changes in fluorescence [18,19]. Taking this into account, solutions of compounds **1a-b** were prepared in

aqueous mixtures of SDS (20 mM, pH 7.5) solution with acetonitrile, 90:10 v/v. SDS aqueous solutions of probes **1a-b** displayed a selective fluorescence quenching in the presence of 10 equivalents of Pd²⁺ (Figures 2b and 3b). Furthermore, further addition to 20 equivalents of each ion confirmed the selectivity of both crown ether benzoxazolyl-alanine derivatives **1a-b** for Pd²⁺ (Figures 2c and 3c).



Figure 3. Preliminary chemosensing study of benzoxazolyl-alanine derivative **1b**: (a) in the presence of 10 equivalents of each cation, in acetonitrile (3×10^{-5} M); (b) in the presence of 10 equivalents of each cation, in SDS (20 mM, pH 7.5)-acetonitrile 90:10 (v/v) solution (1×10^{-5} M); (c) in the presence of 20 equivalents of each cation, in SDS (20 mM, pH 7.5)-acetonitrile 90:10 (v/v) solution (1×10^{-5} M).

4. Conclusions

In summary, two novel benzoxazolyl-alanines bearing a crown ether moiety **1a-b** were evaluated as fluorimetric chemosensors for several ions in acetonitrile and in mixtures of acetonitrile and aqueous SDS solution. As expected, these probes had a fluorimetric response for different cations in acetonitrile solutions but, most importantly, in aqueous mixtures using SDS anionic surfactant both crown ether benzoxazolyl-alanines displayed a selective fluorimetric quenching in the presence of Pd²⁺. These results clearly indicated that probes **1a-b** could be used to detect the palladium cation in environmental and biological samples, with remarkable selectivity.

Acknowledgments: The authors acknowledge Fundação para a Ciência e Tecnologia—FCT (Portugal) for funding through CQUM (UID/QUI/00686/2020).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Sun, Q.; Qiu, Y.; Chen, J.; Wu, F.S.; Luo, X.G.; Guo, Y.R.; Han, X.Y.; Wang, D.W. A Colorimetric and Fluorescence Turn-on Probe for the Detection of Palladium in Aqueous Solution and its Application *in vitro* and *in vivo*. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* **2020**, *239*, 118547, doi:10.1016/j.saa.2020.118547.
2. Ke, B.; Chen, H.; Cui, Y.; Ma, L.; Liu, Y.; Hu, X.; Bai, Y.; Du, L.; Li, M. A Bioluminescent Strategy for Imaging Palladium in Living Cells and Animals with Chemoselective Probes Based on Luciferin-Luciferase System. *Talanta* **2019**, *194*, 925–929, doi:10.1016/j.talanta.2018.11.006.
3. Iavicoli, I.; Fontana, L.; Bergamaschi, A. Palladium: Exposure, Uses, and Human Health Effects. In *Encyclopedia of Environmental Health*, 1st ed.; Nriagu, J., Ed.; Elsevier: Amsterdam, The Netherlands, 2011; pp. 307–314, doi:10.1016/B978-0-444-52272-6.00575-4.

4. Kielhorn, J.; Melber, C.; Keller, D.; Mangelsdorf, I. Palladium—A Review of Exposure and Effects to Human Health. *Int. J. Hyg. Environ. Health* **2002**, *205*, 417–432, doi:10.1078/1438-4639-00180.
5. Balamurugan, R.; Liu, J.H.; Liu, B.T. A Review of Recent Developments in Fluorescent Sensors for the Selective Detection of Palladium Ions. *Coord. Chem. Rev.* **2018**, *376*, 196–224, doi:10.1016/j.ccr.2018.07.017.
6. Zhang, J.; Zhang, L.; Zhou, Y.; Ma, T.; Niu, J. A Highly Selective Fluorescent Probe for the Detection of Palladium(II) Ion in Cells and Aqueous Media. *Microchim. Acta* **2013**, *180*, 211–217, doi:10.1007/s00604-012-0922-2.
7. Ferreira, R.C.M.; Raposo, M.M.M.; Costa, S.P.G. Heterocyclic Amino Acids as Fluorescent Reporters for Transition Metals: Synthesis and Evaluation of Novel Furyl-benzoxazol-5-yl-L-alanines. *New J. Chem.* **2018**, *42*, 3483–3492, doi:10.1039/c7nj04459c.
8. Esteves, C.I.C.; Ferreira, R.C.M.; Raposo, M.M.M.; Costa, S.P.G. New Fluoroionophores for Metal Cations Based on Benzo[d]oxazol-5-yl-alanine Bearing Pyrrole and Imidazole. *Dyes Pigments* **2018**, *151*, 211–218, doi:10.1016/j.dyepig.2017.12.040.
9. Esteves, C.I.C.; Raposo, M.M.M.; Costa, S.P.G. New 2,4,5-Triarylimidazoles Based on a Phenylalanine Core: Synthesis, Photophysical Characterization and Evaluation as Fluorimetric Chemosensors for Ion Recognition. *Dyes Pigments* **2016**, *134*, 258–268, doi:10.1016/j.dyepig.2016.07.020.
10. Esteves, C.I.C.; Raposo, M.M.M.; Costa, S.P.G. Synthesis and Evaluation of Benzothiazolyl and Benzimidazolyl Asparagines as Amino Acid Based Selective Fluorimetric Chemosensors for Cu²⁺. *Tetrahedron* **2010**, *66*, 7479–7486, doi:10.1016/j.tet.2010.07.063.
11. Martins, C.D.F.; Raposo, M.M.M.; Costa, S.P.G. Metallic Ion Sensing with a Benzothiazole-Based Fluorimetric Chemosensor. *Proceedings* **2019**, *9*, doi:10.3390/ecsoc-22-05699.
12. Esteves, C.I.C.; Batista, R.M.F.; Raposo, M.M.M.; Costa, S.P.G. Novel Functionalised Imidazo-Benzocrown Ethers Bearing a Thiophene Spacer as Fluorimetric Chemosensors for Metal Ion Detection. *Dyes Pigments* **2016**, *135*, 134–142, doi:10.1016/j.dyepig.2016.04.037.
13. Guzew, K.; Szmigiel, D.; Wróblewski, D.; Milewska, M.; Karolczak, J.; Wiczak, W. New Fluorescent Probes Based on 3-(2-Benzoxazol-5-yl)alanine Skeleton—Synthesis and Photophysical Properties. *J. Photochem. Photobiol. A* **2007**, *187*, 87–96, doi:10.1016/j.jphotochem.2006.09.019.
14. Fery-Forgues, S.; Al-Ali, F. Bis(Azacrown Ether) and Bis(Benzocrown Ether) Dyes: Butterflies, Tweezers and Rods in Cation Binding. *J. Photochem. Photobiol. C* **2004**, *5*, 139–153, doi:10.1016/j.jphotochemrev.2004.07.001.
15. Gokel, G.W.; Leevy, W.M.; Weber, M.E. Crown Ethers: Sensors for Ions and Molecular Scaffolds for Materials and Biological Models. *Chem. Rev.* **2004**, *104*, 2723–2750, doi:10.1021/cr020080k.
16. Batista, P.M.R. Synthesis and Evaluation of Novel Amino Acids Functionalized with Crown ether and Benzoxazole Units as Fluorimetric Sensors for Cations. Master's Thesis, University of Minho, Braga, Portugal, 2011.
17. Okda, H.E.; El Sayed, S.; Ferreira, R.C.M.; Gonçalves, R.C.R.; Costa, S.P.G.; Raposo, M.M.M.; Martínez-Mañez, R.; Sancenón, F. *N,N*-Diphenylanilino-Heterocyclic Aldehyde-Based Chemosensors for UV-Vis/NIR and Fluorescence Cu(II) Detection. *New J. Chem.* **2019**, *43*, 7393–7402, doi:10.1039/c9nj00880b.
18. Pallavicini, P.; Pasotti, L.; Patroni, S. Residual and Exploitable Fluorescence in Micellar Self-Assembled ON–OFF Sensors for Copper(II). *Dalton Trans.* **2007**, *48*, 5670–5677, doi:10.1039/b710765j.
19. Arduini, M.; Rampazzo, E.; Mancin, F.; Tecilla, P.; Tonellato, U. Template Assisted Self-Organized Chemosensors. *Inorg. Chim. Acta* **2007**, *360*, 721–727, doi:10.1016/j.ica.2006.06.017.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).