

# Identification of Less Harmful Pesticides against Honey Bees: Shape-Based Similarity Analysis <sup>†</sup>

Luminita Crisan <sup>\*</sup>, Ana Borota, Simona Funar-Timofei and Alina Bora <sup>\*</sup>

“Coriolan Dragulescu” Institute of Chemistry Timisoara, 24 Mihai Viteazul Av., 300223 Timisoara, Romania; ana\_borota@acad-icht.tm.edu.ro (A.B.); timofei@acad-icht.tm.edu.ro (S.F.-T.)

<sup>\*</sup> Correspondence: lumi\_crisan@acad-icht.tm.edu.ro (L.C.); lina.bora@gmail.com (A.B.); Tel.: +40-256-491-818 (L.C.); Fax: +40-256-491-824 (L.C.)

<sup>†</sup> Presented at the 24th International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2020; Available online: <https://ecsoc-24.sciforum.net/>.

Received: date; Accepted: date; Published: date

**Abstract:** The high concentration of the pesticide residues existing in vegetation, crops, and various edible products and the prolonged exposure to them, can harm human life and contribute to the disappearance of honey bees, several avian and animal species. The honey bees (*Apis mellifera*), efficient pollinators in addition to honey producers, are also considered important non-target test species for terrestrial toxicity assessment of chemicals. In this context, using thiacloprid and acetamiprid as queries, we performed a 3D similarity search to select new potential products with less harmful effects against bees. For similarity search, a small dataset of 302 compounds with pesticide activity, compiled from literature, was used. The first 10 compounds were selected and structurally analyzed according to the TanimotoCombo metrics, and compared with each of these two queries known to be effective, easily metabolized, and less toxic for bees. This approach came as a forward step in the research of pesticide ecotoxicological risk assessment, for the evaluation of their potential impact on the pollinator insects and the environment.

**Keywords:** *apis mellifera*; 3D similarity search; thiacloprid; acetamiprid

---

## 1. Introduction

Neonicotinoids are the most commonly used insecticides for pest control. The main problems associated with the use of these insecticides, alone or in combination with other factors, are related to their negative impact against many species of insects, including bees. Also, the application mode of insecticides (e.g., direct spray, soil and seed application, etc.) plays a key role in their impact against pollinators. Honeybees are considered the most successful and commercially valuable pollinators due to their pollination functions, maintenance of biodiversity in natural ecosystem as well as the commercial products delivered, such as honey, propolis, etc. This negative neonicotinoid impact on bees is very studied, because the effect at different doses is still not fully understood [1]. Honeybees can be envisaged as very vulnerable to pesticides because their genome has fewer genes compared to other insects [2]. The exposure to neonicotinoids can influence the flying and the foraging ability, reproduction, and pollination for many useful insects including honeybees [3]. In particular, they affect bees by forgetting the locations of flowers or even hives, and also by increasing the fertility of queens or bumblebee [3]. In this context, the strategy of designing new neonicotinoids by modifying existing structures may be an effective way to overcome this harmful influence against pollinators.


Of the eight marketed neonicotinoids [<https://www.reportbuyer.com/product/3952801/>], thiacloprid and acetamiprid are considered to be less toxic for honeybees. These neonicotinoids are considered a group of neurotoxins, chemically similar to nicotine, which acts specifically as nicotinic acetylcholine receptor (nAChR) antagonists [4].

Theoretical methods (QSAR, linear and nonlinear regression techniques, molecular docking, 2D and 3D similarity search, etc.) applied in cheminformatics to discover new drugs, have also been successfully employed to predict novel insecticides and pesticides with less polluting and toxic effects, to fill data gaps and to reduce toxicity testing on animals. In the current work, thiacloprid and acetamiprid were used as template molecules in 3D similarity analysis accomplished with ROCS (Rapid Overlay of Chemical Structures) [5,6] from OpenEye package (<https://www.eyesopen.com/>). The main goal is to find novel compounds, similar to template molecules, which are easy to metabolize and less toxic for bees.

## 2. Methods

From literature [7–22], a dataset of 302 compounds with known pesticide activity was selected and used further for a 3D-similarity search. The conformational space of selected compounds was carried out with the Omega tool, Openeye (OMEGA v.2.5.1.4, OpenEye Scientific Software, Santa Fe, NM, USA. [www.eyesopen.com](http://www.eyesopen.com)) [23]. A maximum of 200 conformers for each compound was generated using default options.

The lowest energy conformers for thiacloprid and acetamiprid were also generated with Omega. The BIOVIA Discovery Studio ([www.3dsbiovia.com](http://www.3dsbiovia.com)) facilities were used for structures visualization and picture delivery (Figure 1).




**Figure 1.** The 2D and 3D structure of the queries compounds.

3D similarity search was performed with ROCS (ROCS v. 3.2.1.4, OpenEye Scientific Software, Santa Fe, NM, USA. <http://www.eyesopen.com>) [5,6]. The ROCS, a shape comparison application, is very faster and useful for handling the large conformer databases. This tool offers alignment and scoring of a database engaging thirteen similarity coefficients. The resulted aligned molecules are ordered by a TanimotoCombo ranking score (as the default option).

## 3. Results and Discussions

3D overlay with ROCS has a great advantage, as it allows optimal visualization of overlapping compounds, which leads to a better understanding of their similarity. The ROCS principles are based on the Gaussian function which is widely used to represent shape and molecular volume.


In this light, several highly occupied regions corresponding to the pyridine ring and a ylidencyanamide group of both template molecules were identified (Figure 2). These regions appear to have multiple hydrogen binding abilities. The pyridine ring of thiacloprid and acetamiprid and thiazolidine ring of thiacloprid may be involved in  $\pi$ - $\pi$  or  $\pi$ - $\sigma$  hydrophobic bonds. As can be seen from Figure 2, the compounds prioritized by ROCS follow the same trend as the query compounds. This trend is in line with the high ShapeTanimoto similarity values (Figure 3), and implicitly the shapes (Figure 2) displayed by all the prioritized compounds.



**Figure 2.** Molecular shapes of queries molecules; the ROCS overlapping of the top compounds ranked by TanimotoCombo against thiacloprid (a) and acetamiprid (b); the surface around queries are rendered.

As can be observed from Figure 3, the 3D coefficients calculated with ROCS for all ten prioritized compounds, against each of the two queries, show values greater than 1.2 for TanimotoCombo, greater than 0.8 for ShapeTanimoto, and greater than 1.2 for ComboScore [24,25]. These high values indicate a very good similarity between selected compounds and acetamiprid and thiacloprid, respectively. Four out of ten prioritized ROCS compounds, highlighted with circles in Figure 3, were considered to have a good profile by comparing with each of the two queries.

2-chloro-5-(4,5-dihydroimidazol-1-ylmethyl)pyridine, **I**, (green circles) is the second compound prioritized by thiacloprid and the 7th by acetamiprid. The (2S)-1-[(6-chloropyridin-3-yl)methyl]imidazolidine-2-carbaldehyde, **II**, (blue circles) is the 3rd compound prioritized in accord with thiacloprid and the 4th by acetamiprid. 4-[[6-chloropyridin-3-yl)methyl](2,2-difluoroethyl)amino)-5H-furan-2-one, **III**, (purple circles) is the 4th compound prioritized towards thiacloprid and the 3rd towards acetamiprid. 2-chloro-5-[[7R)-7-methyl-2H,3H,5H,6H,7H-imidazo[1,2-a]pyridin-1-yl)methyl]pyridine, **IV**, (cyan circles) is the 3rd compound prioritized by thiacloprid and the 8th by acetamiprid. Based on the good qualities demonstrated by ROCS analysis, these four compounds will be subjected, in further studies, to molecular docking and molecular dynamic simulation.



**Figure 3.** The chemical structures of the top ten compounds ranked by TanimotoCombo; against thiacloprid (**left**) and acetamiprid (**right**).

The computed pharmacokinetic proprieties of the selected four compounds and the two queries are listed in Table 1. These are performed with a freely accessible web server pkCSM (<http://biosig.unimelb.edu.au/pkcsml/>). The pkCSM program affords a fast and easy method to early assessment of compounds [26]. Regarding the CNS permeability, it could be observed that all four selected compounds showed logPS values lower than -3 being considered unable to penetrate the CNS of insects.

**Table 1.** Pharmacokinetic Properties of thiacloprid, acetamiprid, and the selected compounds \*.

		Thiacloprid	Acetamiprid	I	II	III	IV
Molecule properties:	MW	252.73	222.679	195.653	225.679	288.681	263.772
	LogP	2.12088	2.06628	1.5789	0.6652	2.2428	2.7338
	#RBN	2	2	2	3	5	2
	#Acceptors	4	3	3	4	4	3
	#Donors	0	0	0	1	0	0
	Surface Area	102.944	93.827	82.063	93.244	113.179	113.057
Toxicity	Max. tolerated dose (human) MRTD (log mg/kg/day)	0.488	0.766	0.555	0.1	0.304	0.192
	hERG I inhibitor	No	No	No	No	No	No
	hERG II inhibitor	No	No	No	No	No	No
	Oral Rat Acute Toxicity (LD50)	3.085	2.906	2.675	2.793	2.969	2.864
	Oral Rat Chronic Toxicity (LOAEL) (log mg/kg_bw/day)	0.699	0.795	0.904	1.186	1.567	0.877
	Hepatotoxicity	Yes	Yes	Yes	Yes	No	Yes
	Skin Sensitisation	No	No	Yes	No	No	No
	<i>T.Pyriformis</i> toxicity pIGC50 (log ug/L)	1.132	0.986	0.869	0.026	0.807	1.025
	Minnow toxicity LC50 (log mM)	1.441	1.566	2.072	2.601	1.892	1.568
	Distribution	VDss (human) (LogL/kg)	-0.134	-0.217	0.161	0.749	-0.074
Fraction unbound (human)		0.486	0.507	0.576	0.897	0.507	0.444
BBB permeability (log BB)		0.114	0.132	0.192	-0.264	0.289	0.593
CNS permeability (logPS)		-2.922	-2.867	-3.199	-3.442	-3.688	-3.475
Excretion	Total Clearance (log mL/min/kg)	0.201	0.193	0.489	0.9	0.484	0.203
	Renal OCT2 substrate	No	No	No	No	No	No

\* MW—Molecular Weight; RBN—Rotatable Bonds; VDss—volume of distribution at steady state; BBB—blood-brain barrier; VDss is considered low if below 0.71 L/kg (log VDss < -0.15) and high if above 2.81 L/kg (log VDss > 0.45). For a given compound: a logBB > 0.3 considered to readily cross the blood-brain barrier while molecules with logBB < -1 are poorly distributed to the brains, a MRTD of less than or equal to 0.477 log(mg/kg/day) is considered low, and high if greater than 0.477 log(mg/kg/day), a LC50 values below 0.5mM (logLC50 < -0.3) are regarded as high acute toxicity, a pIGC50 (negative logarithm of the concentration required to inhibit 50% growth in log ug/L) is predicted, with a value > -0.5 log ug/L is considered toxic. Compounds with a logPS > -2 are considered to penetrate the Central Nervous System (CNS), while those with logPS < -3 are considered as unable to penetrate the CNS.

#### 4. Conclusions

In this study, thiacloprid and acetamiprid were used as queries in order to find new potential compounds with less harmful effects against bees. Four compounds (2-chloro-5-(4,5-dihydroimidazol-1-ylmethyl)pyridine, (2S)-1-[(6-chloropyridin-3-yl)methyl]imidazolidine-2-carbaldehyde, 4-[[[(6-chloropyridin-3-yl)methyl](2,2-difluoroethyl)amino]-5H-furan-2-one, and 2-chloro-5-[[[(7R)-7-methyl-2H,3H,5H,6H,7H-imidazo[1,2-a]pyridin-1-yl]methyl]pyridine) were selected as similar in shape and volume to both queries, which are known for their reduced toxic effect against bees [27,28]. This approach is a first attempt to find novel compounds with enhanced safety profile against the pollinator insects and the environment.

**Author Contributions:** L.C. and A.B. have conceived of the presented idea, designed and accomplished the computational framework, including editing; A.B. performed some computational determinations; S.F.-T. selected and provided the pesticide dataset. All authors contributed to the writing of the paper and approved the content.

**Funding:** This research received no external funding

**Acknowledgments:** The authors thank ChemAxon Ltd., OpenEye Ltd., and BIOVIA software Inc. (Discovery Studio Visualizer) for providing academic license. The authors wish to thank Schrödinger Inc for providing an academic trial license to complete the calculations for this paper. Project No 1.1 and No. 1.2 of the “Coriolan Dragulescu” Institute of Chemistry, Timisoara, Romanian Academy, financially supported the current work.

**Conflicts of Interest:** The authors indicate no potential conflict of interest.

#### References

1. Hassani, A.K.; Dacher, M.; Gary, V.; Lambin, M.; Gauthier, M.; Armengaud, C. Effects of Sublethal Doses of Acetamiprid and Thiamethoxam on the Behavior of the Honeybee (*Apis mellifera*). *Arch. Environ. Contam. Toxicol.* **2008**, *54*, 653–661.
2. Claudianos, C.; Ranson, H.; Johnson, R.M.; Biswas, S.; Schuler, M.A.; Berenbaum, M.R.; Feyereisen, R.; Oakeshott, J.G. A deficit of detoxification enzymes: Pesticide sensitivity and environmental response in the honeybee. *Insect Mol. Biol.* **2006**, *15*, 615–636.
3. Buszowski, B.; Bukowska, M.; Ligor, M.; Staneczko-Baranowska, I. A holistic study of neonicotinoids neuroactive insecticides—Properties, applications, occurrence, and analysis. *Environ. Sci. Pollut. Res.* **2019**, *26*, 34723–34740.
4. Blacquièrre, T.; Smagghe, G.; van Gestel, C.A.M.; Mommaerts, V. Neonicotinoids in bees: A review on concentrations, side-effects and risk assessment. *Ecotoxicology* **2012**, *21*, 973–992.
5. Hawkins, P.C.D.; Skillman, A.G.; Nicholls, A. Comparison of Shape-Matching and Docking as Virtual Screening Tools. *J. Med. Chem.* **2007**, *50*, 74–82.
6. Rush, T.S.; Grant, J.A.; Mosyak, L.; Nicholls, A. A shape-based 3-D scaffold hopping method and its application to a bacterial protein–protein interaction. *J. Med. Chem.* **2005**, *48*, 1489–1495.
7. Hou, S.; Zhuang, Y.; Deng, Y.; Xu, X. Photostability study of cis-configuration neonicotinoid insecticide cycloxaprid in water. *J. Environ. Sci. Health B* **2017**, *52*, 525–537.
8. Zou, M.; Tian, X.; Chen, N.; Shao, X. Nematicidal Activity of Sprio and Bridged Heterocyclic Neonicotinoid Analogues against *Meloidogyne incognita*. *Lett. Drug Des. Discov.* **2015**, *12*, 439–445.
9. Fu, Q.; Zhang, J.; Xu, X.; Wang, H.; Wang, W.; Ye, Q.; Li, Z. Diastereoselective Metabolism of a Novel Cis-Nitromethylene Neonicotinoid Paichongding in Aerobic Soils. *Environ. Sci. Technol.* **2013**, *47*, 10389–10396.
10. Xu, T.; Wang, X.; Zhang, Q.; Fan, J.; Liu, L.; Liu, M.; Zhang, H.; Li, J.; Guo, Y. Iodine-mediated oxidative cyclization for one pot synthesis of new 8-hydroxyquinaldine derivatives containing a N-phenylpyrazole moiety as pesticidal agents. *Bioorganic Med. Chem. Lett.* **2018**, *28*, 3376–3380.
11. Kong, W.; Bao, Y.; Ma, Q.; Xu, H. Synthesis and biological activities of novel pyrazolomatrine derivatives. *Bioorganic Med. Chem. Lett.* **2018**, *28*, 3338–3341.
12. Chen, C.; Chen, J.; Gu, H.; Bao, N.; Dai, H. Design, Synthesis, and Biological Activities of Novel Pyrazole Oxime Compounds Containing a Substituted Pyridyl Moiety. *Molecules* **2017**, *22*, 878.
13. Jeanmart, S.; Edmunds, A.J.F.; Lamberth, C.; Pouliot, M. Synthetic approaches to the 2010–2014 new agrochemicals. *Bioorganic Med. Chem.* **2016**, *24*, 317–341.

14. Cheng, X.; Wang, Y.; Li, W.; Li, Q.; Luo, P.; Ye, Q. Nonstereoselective foliar absorption and translocation of cycloxaprid, a novel chiral neonicotinoid, in Chinese cabbage. *Environ. Pollut.* **2019**, *252*, 1593–1598.
15. Sun, C.W.; Wang, J.; Wu, Y.; Nan, S.B.; Zhang, W.G. Novel nitenpyram analogues with tetrahydropyridone fixed cis-configuration: Synthesis, insecticidal activities, and molecular docking studies. *Heterocycles* **2013**, *87*, 1865–1880.
16. Nishiwaki, H.; Kuriyama, M.; Nagaoka, H.; Kato, A.; Akamatsu, M.; Yamauchi, S.; Shuto, Y. Synthesis of imidacloprid derivatives with a chiral alkylated imidazolidine ring and evaluation of their insecticidal activity and affinity to the nicotinic acetylcholine receptor. *Bioorganic Med. Chem.* **2012**, *20*, 6305–6312.
17. Jiang, D.; Zheng, X.; Shao, G.; Ling, Z.; Xu, H. Discovery of a Novel Series of Phenyl Pyrazole Inner Salts Based on Fipronil as Potential Dual-Target Insecticides. *J. Agric. Food Chem.* **2014**, *62*, 3577–3583.
18. Liu, S.H.; Peng, W.; Qu, Y.Y.; Xu, D.; Li, H.Y.; Song, D.L.; Duan, H.X.; Yang, X.L. Synthesis, insecticidal activity and molecular docking study of clothianidin analogues with hydrazide group. *Chin. Chem. Lett.* **2014**, *25*, 1017–1020.
19. Hua, X.; Mao, W.; Fan, Z.; Ji, X.; Li, F.; Zong, G.; Song, H.; Li, J.; Zhou, L.; Zhou, L.; et al. Novel Anthranilic Diamide Insecticides: Design, Synthesis, and Insecticidal Evaluation. *Aust. J. Chem.* **2014**, *67*, 1491–1503.
20. Shen, H.F.; Chen, X.; Liao, P.; Shao, X.S.; Li, Z.; Xu, X.Y. Design, synthesis, and insecticidal bioactivities evaluation of pyrrole- and dihydropyrrole-fused neonicotinoid analogs containing chlorothiazole ring. *Chin. Chem. Lett.* **2015**, *3245*, 1–4.
21. Liu, J.; Chen, Y.; Hua, X.; Wan, Y.; Wei, W.; Song, H.; Yu, S.; Zhang, X.; Li, Z. Design, synthesis, antifungal activities and SARs of (R)-2-Aryl-4,5-dihydrothiazole-4-carboxylic acid derivatives. *Chin. J. Chem.* **2015**, *33*, 1269–1275.
22. He, Y.; Hu, D.; Lv, M.; Jin, L.; Wu, J.; Zeng, S.; Yang, S.; Song, B. Synthesis, insecticidal, and antibacterial activities of novel neonicotinoid analogs with dihydropyridine. *Chem. Cent. J.* **2013**, *7*, 76.
23. Hawkins, P.C.D.; Nicholls, A. Conformer generation with OMEGA: Learning from the data set and the analysis of failures. *J. Chem. Inf. Model.* **2012**, *52*, 2919–2936.
24. Muchmore, S.W.; Debe, D.A.; Metz, J.T.; Brown, S.P.; Martin, Y.C.; Hajduk, P.J. *J. Chem. Inf. Model.* **2008**, *48*, 941–948.
25. Bortolato, A.; Perruccio, F.; Moro, S. *Successful Applications of In Silico Approaches for Lead/Drug Discovery*; Miteva, M.A., Ed.; Bentham Science Publishers: Sharjah, UAE, 2011.
26. Pires, D.E.; Blundell, T.L.; Ascher, D.B. pkCSM: Predicting small-molecule pharmacokinetic and toxicity properties using graph-based signatures. *J. Med. Chem.* **2015**, *58*, 4066–4072.
27. Laurino, D.; Porporato, M.; Patetta, A.; Manino, A. Toxicity of neonicotinoid insecticides to honey bees: Laboratory tests. *Bull. Insectol.* **2011**, *64*, 107–113.
28. Brandt, A.; Hohnheiser, B.; Sgolastra, F.; Bosch, J.; Meixner, M.D.; Büchler, R. Immunosuppression response to the neonicotinoid insecticide thiacloprid in females and males of the red mason bee *Osmia bicornis* L. *Sci. Rep.* **2020**, *10*, 1–10.

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).