

Polycyclic aromatic hydrocarbons in commercial herbal teas

Edward MUNTEAN^{1, 2,*} and Nicoleta MUNTEAN³

¹University of Agricultural Sciences and Veterinary Medicine Cluj Napoca, Romania

²Research & Development Station for Agriculture, Turda, Romania

³Regional Center of Public Health Cluj Napoca, National Institute of Public Health, Romania

* Corresponding author: edimuntean@yahoo.com



Abstract

Contamination of herbal teas with PAHs can occur as a result of environmental pollution as well as during processing. This research fits into current concerns directed towards establishing the PAHs' contamination level of plant resources, the focus being on the priority PAHs as established by the US Environmental Protection Agency's and by the European Union. In this study, the PAHs' content of several commercial herbal teas brands was assessed by reversed-phase high performance liquid chromatography (HPLC) using an Agilent 1100 system with fluorescence and photodiode array detectors. The obtained data revealed that the maximum overall PAH's contamination in the studied herbal teas was recorded for dandelion, mulberry and St.John's wort, all of these exposing high surface area of leaves; from the 15 tracked PAHs, only three were present in all samples (acenaphthene, fluorene and benzo(b)fluoranthene), while anthracene and benzo(g,h,i)perylene were quantified in only two samples; the maximum levels of individual PAHs were recorded in dandelion.

Keywords: medicinal herbal teas, health risk, polycyclic aromatic hydrocarbons, contamination

Introduction/ context

- The presence of polycyclic aromatic hydrocarbons (PAHs) in medicinal plants may result in health risk to consumers, which are generally hoping that a natural remedy may be helpful in the healing process.
- Contamination of herbal teas with PAHs can occur as a result of environmental pollution as well as during processing (e.g. in some drying or preservation stages that allow combustion products to come into contact with medicinal plants).
- This research fits into current concerns directed towards establishing the PAHs' contamination level of plant resources, the focus being on the priority PAHs as established by the US Environmental Protection Agency's and by the European Union: benzo[a]-anthracene, benzo[b]fluoranthene, indeno[1,2,3-cd]pyrene, benzo[k]-fluoranthene, benzo[a]pyrene, dibenzo-[a,h]anthracene, naphthalene, acenaphthene, acenaphthylene, fluorene, phenanthrene, anthracene, fluoranthene, pyrene, chrysene and benzo(g,h,i)perylene.

Research objective

Assessment of PAHs' content from several commercial herbal teas brands using high performance liquid chromatography (HPLC): basil (*Basilici Folium*), chamomile (*Chamomillae Folium*), mulberry (*Mori Folium*), plantain (*Plantaginis Folium*), dandelion (*Taraxaci Herba*), mint (*Menthae Herba*), St. John's wort (*Hyperici Herba*), black locust (*Acacia Flos*), common elder (*Sambuci Flos*), lime (*Tiliae Flos*), marigold (*Calendula Flos*) and fennel (*Foeniculi Fructus*).

Materials and methods

Plant material: 12 herbal tea brands were purchased from local hypermarkets.

Extraction: ~5 g tea samples were grounded, weighed then extracted with 20 mL hexane: acetone (1:1) for 30 minutes by ultrasonication, at room temperature ; the supernatant was decanted while the residue was re-extracted with a new volume of solvent mixture in the same conditions, the procedure being repeated twice more. The obtained extracts were combined, passed through an anhydrous Na_2SO_4 layer and evaporated to dryness in a rotary evaporator; the residue was dissolved in 2 mL acetonitrile, then the final solution was filtered through 0.47 mm PTFE membrane filter and injected in the HPLC system. Three replicates from each tea brand followed the same procedure.

HPLC analysis: using an Agilent 1100 system with fluorescence detection in which baseline separations for the above-mentioned PAHs was accomplished using an Envirosep PP column in a total run time less than 30 minutes (fig.1).




Figure 1. HPLC chromatogram of PAHs – standards' mixture

Results and Discussion

- The maximum overall PAH's contamination in the studied herbal teas was recorded for dandelion (131.15 $\mu\text{g}/\text{kg}$), mulberry (27.78 $\mu\text{g}/\text{kg}$) and St.John's wort (18.28 $\mu\text{g}/\text{kg}$), all of these exposing high surface area of leaves to environmental contamination with PAHs (table 1).
- From the 15 quantified PAHs, only three were present in all samples (acenaphthene, fluorene and benzo(b)fluoranthene), while anthracene and benzo(g,h,i)perylene were quantified in only two samples.
- The maximum levels of individual PAHs were recorded in dandelion: 30.79 $\mu\text{g}/\text{kg}$ naphthalene, 28.91 $\mu\text{g}/\text{kg}$ pyrene, 23.79 $\mu\text{g}/\text{kg}$ phenanthrene and 10.28 $\mu\text{g}/\text{kg}$ benzo(b)fluoranthene. Dandelion herbal tea exhibit also the largest PAH's range (14 PAHs).
- The lowest PAHs concentrations were recorded in basil and fennel (1,96 and 2.20 $\mu\text{g}/\text{kg}$), these being the least contaminated also in terms of PAH's range (only 7 from the studied PAHs).
- Carcinogenic PAHs ranged from 0.45 $\mu\text{g}/\text{kg}$ (lime) to 36.54 $\mu\text{g}/\text{kg}$ (dandelion), with St.John's wort revealing the maximum concentration of benzo(g,h,i) perylene from all the samples.

Results and Discussion

Table 1. Mean values of recorded PAHs' concentrations in the studied herbal teas [$\mu\text{g}/\text{kg}$]

	Basil	Fennel	Lime	Platain	Black locust	Chamomile	Mint	Marygold	Elder	St. John's wort	Mulberry	Dandelion
Naphtalene	N.D.	N.D.	N.D.	N.D.	0.04	N.D.	1.40	N.D.	7.48	N.D.	N.D.	30.79
Acenaphtene	0.43	0.81	0.52	0.44	0.63	2.29	5.25	2.51	0.07	4.67	8.02	0.91
Fluorene	0.06	0.86	0.21	0.46	0.54	0.26	0.48	0.46	0.56	0.47	2.71	2.14
Phenanthrene	N.D.	N.D.	N.D.	N.D.	N.D.	2.63	N.D.	2.21	3.43	N.D.	7.25	23.79
Anthracene	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	0.21	N.D.	N.D.	4.14
Fluoranthene	N.D.	N.D.	N.D.	N.D.	0.55	N.D.	N.D.	N.D.	2.49	2.52	7.33	3.94
Pyrene	N.D.	N.D.	1.30	0.85	N.D.	N.D.	0.27	N.D.	1.29	N.D.	0.13	28.91
Benz(a)anthracene	0.13	0.32	0.11	0.13	0.21	0.22	0.28	4.98	N.D.	0.10	0.25	N.D.
Chrysene	0.92	0.07	0.08	N.D.	2.40	3.34	2.04	0.59	0.04	0.22	1.51	1.16
Benzo(b)fluoranthene	0.24	0.06	0.16	0.39	0.13	0.19	0.23	0.51	0.15	0.15	0.30	10.28
Benzo(k)fluoranthene	0.05	N.D.	0.07	0.30	0.05	0.04	0.04	0.01	0.14	0.08	0.16	6.11
Benzo(a)pyrene	N.D.	0.03	0.03	0.19	0.06	0.04	0.06	0.14	0.10	0.14	0.11	8.89
Dibenzo(a,h) anthracene	0.13	0.05	N.D.	0.68	0.11	N.D.	N.D.	0.09	0.10	0.14	N.D.	0.15
Benzo(g,h,i)perylene	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	6.73	N.D.	2.10
Indeno(1,2,3-c,d)pyrene	N.D.	N.D.	N.D.	N.D.	0.15	N.D.	N.D.	N.D.	N.D.	3.07	N.D.	7.85
Total	1.96	2.20	2.48	3.43	4.87	9.01	10.06	11.50	16.07	18.28	27.78	131.15
Total carcinogenic	1.47	0.53	0.45	1.69	3.12	3.83	2.66	6.33	0.54	10.62	2.34	36.54

N.D = not detected

Results and Discussion

- Based on the obtained data, a PCA model was developed using 14 variables (PAHs' concentrations), in which: the first two principal components explain 76.33% variance (figures 2 and 3), "mulberry" is an outlier, "dandelion" is outside the model border.
- PCA revealed a close correlation between the recorded concentrations of naphthalene, anthracene, pyrene, benzo(a)pyrene, benzo(b)fluoranthene and benzo(k)fluoranthene. (figure 3).




Figure 2. PCA scores




Figure 3. PCA biplot

Conclusions

- 15 PAHs were determined in 12 herbal teas' brands, each product showing a particular contamination pattern
- PAHs' contamination is more important for *herba* (dandelion, mulberry, St. John's wort) > *flos* (common elder, black locust, lime) > *semen* (fennel) –with some exceptions.
- The content of carcinogenic PAHs' are higher than 50% from the total PAHs' content only for basil, black locust, marigold and St. John's wort.
- The obtained data are comparable with the published ones [1, 2], but showing much smaller PAHs' concentrations.

Many thanks for your attention!



**Please send your questions to
[edimuntean@yahoo.com!](mailto:edimuntean@yahoo.com)**

**IECPS
2020**