

Type of the Paper (Proceedings, etc.)

OPPORTUNITIES FOR IMPROVING FEED USE EFFICIENCY FOR SUSTAINABLE DAIRY PRODUCTION IN PAKISTAN

Muhammad Tariq

Department of Livestock Management, University of Agriculture Faisalabad Sub-Campus Toba Tek
Singh

*Email: tariqlm@uaf.edu.pk

Abstract: Buffalo and cattle are our main dairy animals making 30% of the total livestock. Out of total milk produced in the country, buffalo contributes about 68 %, followed by cattle (27%) rest (5%) by sheep, goat and camel. They are kept both in rural as well peri (urban) dairy production systems. These dairy animals mostly strive on low-quality feed stuffs including roughages and crop-residues with poor nutritive value resulting in poor production and reproduction performance. Recent investigations and published data show that there is also issue of unjustified feeding without considering the production and physiological stages of dairy animals resulting in overfeeding of non-productive and under-feeding of productive-animals leading to poor feed use efficiency. There is also no separate feeding system for different classes of the dairy animals. Furthermore, in the prevailing dairy production scenario there is absence of efficient feeding system especially devised for growing heifers (kept as replacement herd) keeping in view their feeding requirements that could reduce their age at puberty with significant reduction in the cost of feeding. Similarly, introduction of milk replacer and early weaning of calves is very effective way without any adverse effects on growth when given free access to good quality calf starter and it could add to dairy economics. Fodder scarcity during certain time of the year winter months (November to January) and other in summer (May-June) in Pakistan is another big constraint toward sustainable dairying. It is concluded that more efficient feed utilization in these dairy production systems could be achieved by developing innovative approaches and solutions (hay and silage making) to fight these scarcity periods and developments in dairy nutrition; like establishment of nutrient requirements for dairy animals for local dairy breeds, adopting group feeding practices of dairy animals according to stage of lactation and production status, proper feeding systems for growing heifers and effective milk replacer feeding for calves. These practical and innovative steps could effectively lead toward sustained dairy production in Pakistan.

Keywords: Dairy Production; Feeding Management; Resource Use Efficiency; Feeding System; Nutrition

1. Introduction

The dairy sector has seen a major intensification during the past fifty years worldwide, with the increase in scales and efficiency of production driven by the demand from a growing human population and increasing incomes of parts of the population [1];[10]. The increase in milk output was achieved through advances in animal nutrition and breeding, feed use efficiency, health management, housing, and automation strategies, along with supporting policies [2]. In Pakistan's peri-urban commercial dairy farms, feed accounts for more than seventy percent of the operational costs [3];[10]., because animals are stall-fed year-round on wheat straw, purchased green fodder, and concentrate feeds which farmers obtain from the markets. Seasonal variations in quantity and

quality of the roughage feeds are a major concern to farmers, especially during the scarcity periods [4], and high feed costs negatively affect the profitability of the dairy enterprises. In consequence inadequate nutrition of dairy animals is considered one of the major limitations to dairy production [5], and may even lead to morbidity and mortality of high milk producers [6] Low gross margins of milk sales due to the high feed costs [7] often push farmers to unethical practices such as adulteration of milk in order to improve their economic situation [6].

In spite of the substantial contribution of livestock to the national economy, per head productivity of dairy animals under present farm conditions is relatively low in Pakistan. Among dairy buffaloes, 98% are producing less than 10 liters of milk per day [8]. High productive and reproductive efficiency of livestock can only be achieved if animals receive the required quantities of feedstuffs providing all nutrients in the needed proportion [9]. A sound intervention strategy to increase income from dairy animals should focus on two fronts: firstly, lowering feed costs, and secondly, increasing individual animal productivity [3]. Both require adequate nutritional management and a high efficiency of feed utilization. Given the paucity of information regarding the nutritional status of lactating animals in Pakistan's commercial and semi-commercial dairy units, specific measures that help dairy farmers to adequately address the previous issues are difficult to devise. This article therefore discussed the feeding practices for lactating dairy animals, in order to identify shortcomings and potential improvements.

2. Existing Feeding Practices Dairy Animals

In commercial and semi-commercial farms most of the farmers use green fodder crops, wheat straw and concentrate feeds (mostly industrial by-products: cottonseed cake, maize oil cake, cereal by-products). Usually green fodder is bought daily from fodder markets; only few farmers grew their own fodder; wheat straw is also mostly bought on a daily basis from retailers dispersed in town. In general, dairy household are chopping the green roughages to 2-3 cm length and mixed these with concentrate feeds and finely chopped wheat straw to make a total mixed ration [10].

3. General Aspect of Dairy Production and Feeding Management

3.1. Different Feed Types

Nutrition is a major factor affecting animal health, milk production and reproduction, and thus overall productivity of dairy animals. Maize and berseem are the preferred green feeds of commercial and semi-commercial dairy farmers as reported by [10]; [3]. Wheat straw is an integral part of the basal dairy diet. All lactating animals are daily supplied with a home-mix of concentrates (4-5 kg per animal) without taking into account the individual milk production [11]. Type and quantity of concentrate feeds varied within a farm and between farms, and depended on season, availability, market price and financial capacity of the farmer. Therefore concentrate feeds, being major sources of protein and energy, are often not fed at the required level due to unavailability and high costs [12], especially for small scale farmers. Furthermore, the quality of concentrate feeds usually not matching that of the roughages since farmers have little or no knowledge of feed quality.

3.2 Feeding of Lactating Buffalo and Cattle

The undifferentiating feeding of lactating buffaloes and cattle in a herd is apparently practiced for ease of management and to save labor, but this practice leads to nutrient undersupply in high

yielding animals and to nutrient oversupply in animals at the end of lactation or of low milk production potential. Both phenomena affect the health and reproductive performance of the animals [3]. Apart from not getting enough fodder, the rations are found to be deficient in energy and protein, some only supplying about 50% of the protein demand of all classes of the dairy animals; conversely, the fiber concentration of the rations is found to be higher than the recommended level. Only very few of the dairies are feeding mineral mixtures to their lactating animals [10].




Figure: 1 Year round fodder situation and availability of different fodders during different months of the year [13]

3.3. Seasonal Impacts on Feeding Management

Different seasons during a year have a significant effect on the offer of feed resources as well as on their nutritional value as well as the availability of different types of green fodders as shown in Figure 1. According to various sources [14]; [15]; [16], there are two periods of fodder scarcity in Punjab: the peak of winter (December and January) and the hot summer months (June and July). In these periods undernourishment and low performance of animals is very likely due to scarcity of fodder in terms of both quantity and quality. Such scarcity can be overcome by proper feeding management and hay and silage making; the latter is recently receiving increasing attention in Pakistan [17].




Figure 2: Shows how silage fits in dairy production system?

4. Economic Situation of these Dairy Farms and Management of Available Resources

Due to differences in the available resources and economic situation of the farms, the average offer of feed dry matter, crude protein and fibrous fractions (NDF) are significantly different between different production systems (rural, semi-commercial and commercial) with similarly differences of values for dry matter offered. Since mixed crop and livestock farmers also cultivated wheat and fodder crops such as maize, sorghum and berseem, they have access to good quality roughages and wheat straw in sufficient quantities and at low cost. Although this was not the case for the (mostly) landless semi-commercial and commercial farmers, the latter disposed of sufficient money and bargaining power (economies of scales effect of these comparatively large and fully commercial dairy units) to purchase high amounts of feed at reduced prices on the fodder markets [10]. The published studies show that the total number of animals per dairy herd, and of lactating animals, respectively, was negatively correlated with the offered amount of green fodder, and positively correlated with the offered amount of concentrates [18].

In another study previously conducted by the author observed that from the amounts of feed offered daily, about 90% is consumed and 10% is lost through leftovers and spillage [10]. Since group feeding of lactating buffaloes and cattle ignores species, stage of lactation and milk yield, the calculational approach assumes that each animal in a herd consumes the same amount of feed; this is undoubtedly an incorrect assumption which cannot be adjusted for here. The findings of other studies on small scale dairy production in Pakistan [10] and India [19] show that in these dairy systems buffaloes (major dairy animal) are always more deficient in energy and total digestible nutrients' supply than cattle.

The trend towards energetically oversupplying cattle and undersupplying buffaloes may be explained by the fact that cattle are kept for their higher milk yields when ignoring milk fat content. By mixing cattle with buffalo milk, farmers are able to sell a higher total volume of milk without the need to adulterate the product, thus receiving higher total revenue for their daily production without having to fear sanctions from their middlemen or clients.

5. Conclusions

It is concluded that more efficient feed utilization in these dairy production systems could be achieved by developing innovative approaches and solutions (hay and silage making) to fight these scarcity periods and developments in dairy nutrition; like establishment of nutrient requirements for dairy animals for local dairy breeds, adopting group feeding practices of dairy animals according to stage of lactation and production status, proper feeding systems for growing heifers and effective milk replacer feeding for calves. These practical and innovative steps could effectively lead toward sustained dairy production in Pakistan.

6. References

1. FAO, 2011b. Dairy development in Pakistan. Food and Agriculture Organization of the United Nations. Rome, Italy.
2. FAO, 2011a. Dairy development in Argentina. Food and Agriculture Organization of the United Nations. Rome, Italy.
3. Habib, G., Hameed, A., Akmal M., 2007. Current feeding management of peri-urban dairy buffaloes and scope for improvement. *Pakistan Veterinary Journal*, 27(1), 35-41.
4. Gillah, K.A., Kifaro, G.C., Madsen, J., 2012. Urban and peri urban dairy farming in East Africa: A review on production levels, constraints and opportunities. *Livestock Research for Rural Development*. Volume 24, Article #198. Retrieved May 6, 2013, from <http://www.lrrd.org/lrrd24/11/gill24198.htm>
5. Olafadehan, O.A., 2007. Potentials for improving dairy cattle production in the agro-pastoral farming system in the derived savanna of Oyo State, Nigeria. PhD Thesis, Department of Animal Science, University of Ibadan, Nigeria.
6. Jalil, H., Rehman, H.U., Maqbool, H.S., Hussain, S.S., 2009. Analysis of milk production system in peri-urban areas of Lahore (Pakistan): A case study. *Pakistan Economic and Social Review*, 47(2), 229-242.
7. Garcia, O., Mahmood K., Hemme, T., 2003. A review of milk production in Pakistan with particular emphasis on small scale producers. Pro-poor Livestock Policy Initiative Working Paper No. 3, FAO, Rome.
8. Khan, R., Qureshi, M.S., Mushtaq, A., Ghufanullah and Naveed, A., 2012. Effect of quality and frequency of drinking water on productivity and fertility of dairy buffaloes. *The Journal of Animal and Plant Sciences*, 22(2), 96-101.
9. NRC, 2001. Nutrient requirements of dairy animals; 7th Ed. National Research Council, National Academy Press, Washington, DC, USA.
10. Tariq, M. Opportunities for improving resource use efficiency in peri-urban dairy herds in Faisalabad, Punjab, Pakistan. Ph.D Thesis. University of Kassel Germany. Sep. 2013
11. Khan, U., Lund, J., Javaid, S., Ul-Hasan, Z., 2008. Economic analysis of milk production in different cattle colonies of Karachi. *Pakistan Journal of Agricultural Sciences*, 45(2), 403-409
12. Nouala, F.S., Akinbamijo, O.O., Adewumi, A., Hoffmann, E., Muetzel, S., Becker, K., 2006. The influence of *Moringaoleifera* leaves as substitute to conventional concentrate on the *in vitro* gas production and digestibility of groundnut hay. *Livestock Research for Rural Development*, Volume 18, Article #121. Retrieved Nov. 30 2020, from <http://www.lrrd.org/lrrd18/9/noua18121.htm>
13. Akmal, M. Scope & opportunities of green fodder for dairy development. Presented at International Dairy Conference in University of Agriculture, Faisalabad, Pakistan. 2008.
14. Teufel, N., Gall, C.F., 1999. Characteristics of village milk production in the Punjab of Pakistan. In: *Deutscher Tropentag 1999 Berlin: Sustainable Technology Development in Animal Agriculture*. Accessed in Nov. 30, 2020, from http://ftp.gwdg.de/pub/tropentag/proceedings/1999/referate/STD_A3.pdf.

15. Afzal, M., Anwar, M., Mirza, M.A., 2007. Some factors affecting milk yield and lactation length in Nili Ravi buffaloes. *Pakistan Veterinary Journal*, 27(3), 113-117.
16. Ali, T., 2007. Pakistan: A case study of milk production and marketing by small and medium scale contract farmers of Haleeb Foods Ltd, Sustainable Markets Group, International Institute for Environment and Development (IIED), London.
17. Teufel, N. personal communication, May, 2013
18. Hagmann, J., 2010. Opportunities and constraints of peri-urban buffalo and dairy cattle systems in Faisalabad, Punjab, Pakistan. ICDD Working Paper No (2) University of Kassel, Germany. Available at http://www.uni-kassel.de/einrichtungen/fileadmin/datas/einrichtungen/icdd/Publications/ICDD_Working_Paper_No.2_Hagmann.pdf
19. Singh, V.K, Singh, P., Verma, A.K., Mehra, U.R., 2008. On farm assessment of nutritional status of lactating cattle and buffaloes in urban, peri-urban and rural areas of Middle Gangetic Plains. *Volume 20, Article #130*. Retrieved Nov. 30, 2020, from <http://www.lrrd.org/lrrd20/8/singh20130.htm>



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).