



Analysis of the risk of infection by COVID-19 taking into account the social determinants of health in Bogotá between February and August 2020

Nathalia Solís-Oñate ^{1, *}, Sthefany Villa-Díaz ², Luis Camilo Blanco-Becerra³ and Ronal Jackson Sierra-Parada ⁴

¹ Faculty of Environmental Engineering, Santo Tomás University; nathaliasolis@usantotomas.edu.co

² Faculty of Environmental Engineering, Santo Tomás University; sthefanyvilla@usantotomas.edu.co

³ Master's Degree in Public Health, Santo Tomás University; luis.blanco@usantotomas.edu.co

⁴ Faculty of Environmental Engineering, Santo Tomás University; ronalsierra@usantotomas.edu.co

* Correspondence: nathaliasolis@usantotomas.edu.co or sthefanyvilla@usantotomas.edu.co

Abstract: This spatial epidemiology study was carried out to evaluate the risk of contracting COVID-19 taking into account the Social Determinants of Health (SDH). The study demonstrated a strong relationship between the data from the model created in the risk analysis and the infections by locality registered during the months of February to August 2020. Kennedy, Bosa and Engativá were the suburbs that presented a higher risk of contagion of COVID-19 compared to the rest of the city. Although for the study of Bogotá the same variables and weights were considered for the estimation of risk, the study found that no area of the city was excluded from presenting the virus. However, areas where there was a higher risk of contracting the virus depended on the characteristics of the local population, the number of reproduction of the virus by suburbs and the applicable poverty rates. With the results obtained and the model implemented, is possible to verify what was established in the 90s regarding the term Syndemia is verified, due to the synergy between two diseases, in this case COVID-19 and base comorbidities present in the population, added to the Social Determinants of Health, which allow a holistic approach to management and prevention measures against contagion, starting in areas of extreme poverty.

Keywords: Social Determinants of Health; risk; vulnerability analysis; COVID-19; SARS-CoV-2; syndemic; transmission

1. Introduction

In the city of Wuhan, China, cases of people with acute respiratory conditions, caused by the new SARS-CoV 2 (Severe Acute Respiratory Syndrome Coronavirus 2) were reported in late 2019, resulting in an emerging disease named COVID-19.

The epicenter of the pandemic between May and June 2020 focuses on the Americas, as 32,437,597 cases are known in this continent alone, in other words, 43% of the cases recorded worldwide and with respect to deaths, 48% of the more than one million deaths are attributed to this area [1]. For Colombia, WHO reports reported a total of 1,482,072 cumulative cases and 40,019 deaths on December 20 [1].

As a result of the pandemic generated by this new virus, social and health inequities in society, including the Social Determinants of Health (SDH), became more pronounced provide vital information since it shows that the most vulnerable people do not have the ease to access timely or good quality health care services due to lack of economic income [2]. In addition to being directly related to the term Sindemia [3] due to the synergy between base comorbidities and in this case the COVID-19 together with the SDH [4] and which has gained strength after the beginning of the pandemic as it fits perfectly into the current situation.

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Based on the above and using air quality data, quality in health service, food, socio-economic strata and base diseases as SDH and then as variables for vulnerability, contagion and death by COVID-19 and effective number of reproduction of the virus by location, the COVID-19 threat, the risk calculation and its respective analysis is made in order to determine the risk to contagion by COVID-19 in the city of Bogota taking into account the SDH in the period from February to August 2020.

2. Materials and methods

A spatial epidemiological study was carried out to show the distribution of the contagions against COVID-19 that occur in the city of Bogotá, Colombia, between February and August 2020 taking into account the SDH. In this study a model is created based on the following formula: Risk= Vulnerability * Menace, this is modified for this study by adding the variables of SDH, effective number of reproduction and population density. Air quality variables (PM10 and 2.5, O3 and NO2) are taken as environmental SDH, socio-economic strata and quality in health service as social SDH, behavioral SDH take the food of the population of the capital and for biological SDH the diseases base that presents the residents of the city of Bogotá D.C. The threat is taken as a constant variable so it is present throughout the city.

2.1 Vulnerability calculation

Vulnerability is taken as SDH according to the Canadian classification, in this case biological determinants are taken as pre-existing diseases of the population, a standard diet of 2000 calories is used as variable for behavioral determinants, while the socio-economic stratum and quality in the health service were taken into account for the social SDH, in terms of environmental SDH the annual averages of the data of PM10 and 2.5, O3 and NO2 for the city of Bogota were used, Finally, information on COVID-19 infections and deaths by localities between February and August is used.

With respect to the calculation of vulnerability, an equal weight is assigned, that is, each determinant was assigned a total of 25 points, in order to obtain 100 points in total.

2.2. Menace

This is taken as the presence of the virus in this case the COVID-19, so in this model the variable is constant, since it is present throughout the city. A value of 10 points was assigned, taking into account the WHO classification of infectious microorganisms for their risk.

2.3. Calculation of Risk

For this calculation the formula of risk and disaster assessment is taken into account, an adaptation is made to the model of this study which is determined and established by the authors, the equation is the following:

$$R = (5(V) * 0,2(M)) * PD * Rt, \quad (1)$$

where, R: risk, 5: vulnerability weighting value, V: vulnerability, 0.2: menace weighting value, M: menace, PD: population density, Rt: effective number of reproduction. This calculation is carried out per block with the aim of obtaining the lowest possible error at the moment of obtaining the arithmetic mean for each locality, then the classification of the risk according to color and type of the averages by locality obtained is carried out.

3. Analysis and results

In Figure 1, it is evident that the results obtained in the map (a) from the model executed has a close relationship with the map showing the contagions of the city (b) recorded during the study period, since the western part presents a high and very high risk, with

localities such as Suba, Ciudad Bolívar and Kennedy, which present a very high risk, in addition to showing that the localities located in the Central-East, such as La Candelaria, Santa Fe, Teusaquillo and Chapinero, reveal a low and very low risk respectively, which resembles the map of contagions. On the other hand, localities such as Engativá, Fontibón and Bosa, with high risk according to the calculations made, resemble the contagions present, since they are found with significantly high risks.

Localities such as Kennedy, Bosa and Ciudad Bolívar, are among the most vulnerable, since the predominant stratum is 2, which is related to the amount of income, for what the country is stratified is characterized by low to medium-low economic income, therefore, they sometimes do not have access to quality education, adequate food, among others. In addition, Suba, Engativá, Kennedy and Usaquén are the localities with the greatest presence of adults older than 65 years, which increases the risk of infection with SARS-CoV-2 and complications derived from COVID-19 [5], since they are extremely vulnerable population due to their biological age and the comorbidities presented by these people [6] [7].




Figure 1. Calculated risk map by locality and COVID-19 infection of the city of Bogotá map. (a) map graphically describing the risk calculated by the model created for this article. (b) map graphically describing the risk calculated by the model created for this article. source: authors.

4. Discussion

The SDH have a great influence since they largely determine the level of risk that the population may present to infectious diseases, in this case, the COVID-19, which has a direct relationship with the results given by the model carried out in this study, which shows that the localities located in the western area of the city, are those that present a higher high and very high risk, in addition to the south-eastern part. The western part of

the city is known for the low quality of the air, since there are located areas of industry with large number of companies [8] [9], being a risk factor for cardiorespiratory diseases which increases the possibilities of contagion to COVID-19 [10] [11]. They also have a population that lives from informal work [12], that is, their livelihood is given by the activities they carry out every day, which leads to the difficult access to education and health services of excellent quality [13], among others.

The aforementioned is related to the term syndemia, in consequence, of the presence of COVID-19 together with base comorbidities that the population had long ago along with social and health inequities [4], which are represented with the SDH, so it is appropriate to define what is currently happening as syndemic and in this way should be addressed in order to reduce the disparities that the same society raises.

One of the strengths of this study is the versatility it has to be able to extrapolate the model to different cities, as well as the ability to add more variables, such as lifestyle, amount of exercise done per week, food consumption per person daily or weekly, the amount of income and number of people contributing to the household, air pollutants such as CO₂ and black carbon and the time of exposure to them, among others; additional benefits of implementing spatial epidemiological studies, including graphical demonstration [14] by mapping of risk distribution in the case of this study, in terms of weaknesses, errors can occur in the data taken from the DANE, since its basis are surveys carried out by ordinary people.

In conclusion, the result of the model created in this study can be used by governmental entities adding more variables in order to make it more robust and in this way attack in a more efficient and precise way situations like the one currently experienced with the COVID-19 and social and health disparities.

References

1. World Health Organization, «COVID-19 Weekly Epidemiological Update,» 2020.
2. M. Pinilla, A. R. Varela y C. González, «Los pobre, los más afectados por la pandemia,» Universidad de los Andes, 07 abril 2020.
3. C. M. Jaramillo, «Al día con la COVID-19,» Medellín, 2020.
4. R. Horton, «Offline: COVID-19 is not a pandemic,» *The Lancet*, vol. 396, n° 10255, 26 September 2020.
5. L. Sauer, «Jonh Hopkins Medicine,» Jonh Hopkins Medicine, 2020. [En línea]. Available: <https://www.hopkinsmedicine.org/health/conditions-and-diseases/coronavirus>. [Último acceso: 10 December 2020].
6. Center for Diseases Control and Prevention, «Older adults».
7. A. Calderón-Larrañaga, S. Dekhtyar, D. L. Vetrano, T. Bellander y L. Fratiglioni, «COVID-19: risk accumulation among biologically and socially vulnerable older populations,» *Ageing Research Reviews*, vol. 63, November 2020.
8. L. A. Rodríguez-Camargo, R. J. Sierra-Parada y .. Luis Camilo Blanco-Becerra, «Análisis espacial de las concentraciones de PM_{2,5} en Bogotá según los valores de las guías de la calidad del aire de la Organización Mundial de la Salud para enfermedades cardiopulmonares, 2014-2015,» *Biomédica*, vol. 20, n° 1, 2020.
9. M. y. E. A. Instituto de Hidrología, «Calidad del aire,» Bogotá.
10. M. Urrutia-Pereira, C. A. Mello-da-Silva y D. Solé, «COVID-19 and air pollution: A dangerous association?,» *Allergologia et Immunopathologia*, vol. 48, n° 5, pp. 496-499, 2020.
11. E. C. f. L. A. a. t. Caribbean, «The social challenge in times of COVID-19,» ECLAC, 2020.
12. N. Frölich, «Coronavirus en Colombia: la cuarentena está causando hambre y protestas,» *Deutsche Welle*, 2020.
13. S. R. Acosta, «La desigualdad de ingreso y salud en Colombia,» *Revista de la Facultad Latinoamericana de Ciencias Sociales*, vol. 24, n° 48, p. Junio, 2016.
14. D. Baker y M. J. Nieuwenhuijsen, *Environmental Epidemiology study methods and application*, 1 ed., vol. 1, New York: Oxford University Press, 2008, pp. 189-197.